

MOUSE SETS

MITCH RUDOMINER

ABSTRACT. In this paper we explore a connection between descriptive set theory and inner model theory. From descriptive set theory, we will take a countable, definable set of reals, A . We will then show that $A = \mathbb{R} \cap \mathcal{M}$, where $\mathcal{M}$ is a canonical model from inner model theory. In technical terms, $\mathcal{M}$ is a “mouse”. Consequently, we say that A is a mouse set. For a concrete example of the type of set A we are working with, let $\text{OD}_n^{\omega_1}$ be the set of reals which are Σ_n definable over the model $L_{\omega_1}(\mathbb{R})$, from an ordinal parameter. In this paper we will show that for all $n \geq 1$, $\text{OD}_n^{\omega_1}$ is a mouse set. Our work extends some similar results due to D.A. Martin, J.R. Steel, and H. Woodin. Several interesting questions in this area remain open.

CONTENTS

0. Introduction	2
1. Prerequisites	6
2. Some Lightface Pointclasses in $L(\mathbb{R})$	15
3. Big Premice	32
4. A Correctness Theorem	59
5. The Ranked Iteration Game	81
6. Petite Premice	91
7. A Comparison Theorem	100
8. Conclusions	125
References	131

Key words and phrases. large cardinals, descriptive set theory, inner model theory.

0. INTRODUCTION

In this paper we will extend some work by D. A. Martin, J. R. Steel, and H. Woodin. We begin by describing this work.

If ξ is a countable ordinal and $x \in \mathbb{R}$ and $n \geq 2$, then we say that $x \in \Delta_n^1(\xi)$ if for every real $w \in \text{WO}$ such that $|w| = \xi$, $x \in \Delta_n^1(w)$. Set

$$A_n = \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in \Delta_n^1(\xi) \}.$$

In short, A_n is the set of reals which are Δ_n^1 in a countable ordinal. Under the hypothesis of Projective Determinacy (PD), the sets A_n are of much interest to descriptive set theorists. If $n \geq 2$ is even then $A_n = C_n$, the largest countable Σ_n^1 set. If $n \geq 3$ is odd then $A_n = Q_n$, the largest countable Π_n^1 set which is closed downward under Δ_n^1 degrees. The sets A_n have been studied extensively by descriptive set theorists. See for example [Ke1], and [KeMaSo].

The work of Martin, Steel, and Woodin shows that there is a connection between the sets A_n , and certain inner models of the form $L[\vec{E}]$, where $\vec{E}$ is a sequence of extenders. In the paper “Projectively Wellordered Inner Models” [St5], Martin, Steel, and Woodin prove the following theorem:

Theorem 0.1 (Martin, Steel, Woodin). *Let $n \geq 1$ and suppose that there are n Woodin cardinals with a measurable cardinal above them. Let $\mathcal{M}_n$ be the canonical $L[\vec{E}]$ model with n Woodin cardinals. Then*

$$\mathbb{R} \cap \mathcal{M}_n = A_{n+2}.$$

The above theorem is also true with $n = 0$. The “canonical $L[\vec{E}]$ model with 0 Woodin cardinals” is just the model L . If there is a measurable cardinal, then $A_2 = \mathbb{R} \cap L$. Actually, this result is true under the weaker hypothesis that $\mathbb{R} \cap L$ is countable.

Now let A^* be the collection of reals which are ordinal definable in $L(\mathbb{R})$. To emphasize the parallel with the sets A_n above, note that assuming that every game in $L(\mathbb{R})$ is determined ($\text{AD}^{L(\mathbb{R})}$), we have:

$$A^* = \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in (\Delta_1^2)^{L(\mathbb{R})}(\xi) \},$$

and A^* is the largest countable $(\Sigma_1^2)^{L(\mathbb{R})}$ set of reals.

In the paper “Inner Models with Many Woodin Cardinals” [St2], Steel and Woodin prove the following theorem:

Theorem 0.2 (Steel, Woodin). *Suppose that there are ω Woodin cardinals with a measurable cardinal above them. Let $\mathcal{M}_\omega$ be the canonical $L[\vec{E}]$ model with ω Woodin cardinals. Then*

$$\mathbb{R} \cap \mathcal{M}_\omega = A^*.$$

Theorems 0.1 and 0.2 are obviously similar to each other. The two theorems establish a very deep connection between descriptive set theory and inner model theory. Notice that between Theorem 0.1 and Theorem 0.2 there is some room. Our goal in writing this paper was to prove a theorem, similar to the these two theorems, for pointclasses Δ between the Δ_n^1 pointclasses and the pointclass $(\Delta_1^2)^{L(\mathbb{R})}$.

More specifically, let us say that a set of reals A is a **mouse set** iff $A = \mathbb{R} \cap \mathcal{M}$ for some canonical $L[\vec{E}]$ model $\mathcal{M}$. (To be more precise, we will take $\mathcal{M}$ to be a countable, iterable *mouse*. A mouse is a model of the form $L_\alpha[\vec{E}]$, for some ordinal α . Theorems 0.1 and 0.2 above can be restated in terms of mice.) Given a pointclass Δ let us set:

$$A_\Delta = \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in \Delta(\xi) \}.$$

Our goal in writing this paper was to find some pointclasses Δ such that $A_n \subset A_\Delta$ for all n , and $A_\Delta \subset A^*$, and then to prove that A_Δ is a mouse set.

To accomplish this task, first we need to have some such pointclasses Δ . In Section 2 we define some lightface pointclasses in $L(\mathbb{R})$ which are similar to the pointclasses of the analytical hierarchy. For ordinals α such that $1 \leq \alpha \leq (\delta_1^2)^{L(\mathbb{R})}$, we define a sequence of pointclasses:

$$\langle \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)}, \Delta_{(\alpha,n)} \mid n \in \omega \rangle.$$

For $\alpha = 1$, we will have that $\Sigma_{(1,n)} = \Sigma_n^1$. So our setting includes the setting of the analytical hierarchy. For $\alpha = (\delta_1^2)^{L(\mathbb{R})}$, we will have that $\Sigma_{(\alpha,0)} = (\Sigma_1^2)^{L(\mathbb{R})}$. So our setting also includes the setting of Theorem 0.2 above.

Then we define:

$$A_{(\alpha,n)} = \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in \Delta_{(\alpha,n)}(\xi) \}.$$

We leave the full technical definition of $A_{(\alpha,n)}$ for later. For the sake of this introduction, it is enough to understand that $A_{(\alpha,n)}$ is the set of reals which are ordinal definable in $L(\mathbb{R})$, “at the $(\alpha, n)^{\text{th}}$ level.” Assuming $\text{AD}^{L(\mathbb{R})}$, these sets are very much like the analytical sets A_n defined above. That is, for even n , $A_{(\alpha,n)} = C_{(\alpha,n)}$, the largest countable $\Sigma_{(\alpha,n)}$ set. For odd n , $A_{(\alpha,n)} = Q_{(\alpha,n)}$, the largest countable $\Pi_{(\alpha,n)}$ set closed downward under $\Delta_{(\alpha,n)}$ degrees.

Because of this analogy, we conjecture that, assuming there are ω Woodin cardinals, $A_{(\alpha,n)}$ is a mouse set, for all (α, n) . To establish this, we need to have some canonical $L[\vec{E}]$ models $\mathcal{M}$. In Section 3, we study mice $\mathcal{M}$ which are “bigger” than the models $\mathcal{M}_n$ mentioned in Theorem 0.1, but “smaller” than the model $\mathcal{M}_\omega$ mentioned in Theorem 0.2. But how can $\mathcal{M}$ have “more than” n Woodin cardinals, for each n , but “less than” ω Woodin cardinals? The answer of course is that “more than” and “less than” here refer to consistency strength. Our mice $\mathcal{M}$ will have the property that for each n , $\mathcal{M}$ satisfies that there is a transitive set model of $\text{ZFC} + \exists n \text{ Woodin Cardinals}$.

In this paper we are not able to fully prove our conjecture that $A_{(\alpha,n)}$ is a mouse set, for all (α, n) . We are able to show this for some (α, n) though. Our argument has several parts, and each part works for a different set of (α, n) . In the end we manage to prove that $A_{(\alpha,n)}$ is a mouse set for all (α, n) such that $\alpha \leq \omega_1^{\omega_1}$, and either $\text{cof}(\alpha) > \omega$, or $\text{cof}(\alpha) \leq \omega$ and $n = 0$. There is nothing special about the ordinal $\omega_1^{\omega_1}$ here. The main property of $\omega_1^{\omega_1}$ which we use is the fact that all ordinals $\alpha < \omega_1^{\omega_1}$ are simply definable from countable ordinals. It is easy to see how to extend our results somewhat beyond $\omega_1^{\omega_1}$. Eliminating the restriction in our work to ordinals $\alpha < \omega_1^{\omega_1}$ does not appear to be a very difficult problem. On the other hand, the restriction to $n = 0$ in the case that $\text{cof}(\alpha) \leq \omega$ seems to be more difficult to overcome. In fact, at the very first step, we are not even able to show that the set $A_{(2,1)}$ is a mouse set. We believe that the full conjecture, for all (α, n) , is a very interesting open problem. In this paper we develop some ideas which should be useful towards solving this problem.

To explain the roles of the various sections of this paper, we will describe our strategy for proving the conjecture that $A_{(\alpha,n)}$ is a mouse set. Our strategy is a natural generalization of the strategy used by Martin, Steel, and Woodin in [St5]. The strategy begins with defining, for each ordinal α and each $n \in \omega$, the notion of a mouse $\mathcal{M}$ being (α, n) -big. If a mouse is (α, n) -big then it satisfies a certain large cardinal hypothesis which is stronger than the hypothesis “ $\exists k$ Woodin cardinals”, for all $k \in \omega$, but weaker than the hypothesis “ $\exists \omega$ Woodin cardinals.” If a mouse is not (α, n) -big, we call it (α, n) -small. The definition of (α, n) -big is given in Section 3. Our definition there only works for $\alpha \leq \omega_1^{\omega_1}$. The next step in our strategy is to prove a theorem which states that if $\mathcal{M}$ is a countable, iterable mouse and $\mathcal{M}$ is (α, n) -big, then $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$. We prove such a theorem in Section 4.

The next step in our strategy is to define, for each ordinal α and each $n \in \omega$, the notion of a mouse $\mathcal{M}$ being $\Pi_{(\alpha,n)}$ -iterable. This notion is an approximation of full iterability, with the added advantage that it is simply definable: The set of (reals coding) countable mice $\mathcal{M}$ which are $\Pi_{(\alpha,n)}$ -iterable is a $\Pi_{(\alpha,n)}$ set. We give the definition of $\Pi_{(\alpha,n)}$ -iterable in Section 5. Our definition there only works for $\alpha < \kappa^{\mathbb{R}}$, where $\kappa^{\mathbb{R}}$ is the least $\mathbb{R}$ -admissible ordinal. Next, our strategy is to prove a theorem which states that if $\mathcal{M}$ and $\mathcal{N}$ are countable mice, and $\mathcal{N}$ is $\Pi_{(\alpha,n)}$ -iterable, and $\mathcal{M}$ is fully iterable and $(\alpha, n+1)$ -small, then $\mathcal{M}$ and $\mathcal{N}$ can be compared.

Unfortunately we are not able to prove this theorem. This accounts for our inability to prove the above-mentioned conjecture, in the case $\text{cof}(\alpha) \leq \omega$ and $n > 0$. We can prove an approximation of this theorem though. In Section 6, we define the notion of a mouse $\mathcal{M}$ being (α, n) -petite. This is a condition which is slightly more restrictive than (α, n) -small. We are able to prove the above-mentioned comparison theorem with (α, n) -petite in place of (α, n) -small. We do this in Section 7, for $\alpha \leq \omega_1^{\omega_1}$.

In the case $\text{cof}(\alpha) > \omega$, (α, n) -petite is synonymous with (α, n) -small. So in this case we get a proof of our conjecture. The proof goes as follows: Let $\mathcal{M}$ be a countable, iterable mouse such that $\mathcal{M}$ is (α, n) -big but every proper initial segment of $\mathcal{M}$ is (α, n) -small. The results of Section 4 give us that $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$, and the results of Section

7 give us that $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha,n)}$. Thus $A_{(\alpha,n)} = \mathbb{R} \cap \mathcal{M}$ and so $A_{(\alpha,n)}$ is a mouse set. This argument is given in Section 8.

In the case $\text{cof}(\alpha) \leq \omega$, our definition of (α, n) -petite is not synonymous with (α, n) -small. So in this case we do not get a proof of our conjecture. However we do manage to eke out a proof of the conjecture in the case $n = 0$. That is, we prove that for all $\alpha \leq \omega_1^{\omega_1}$, $A_{(\alpha,0)}$ is a mouse set. Actually, the fact that $A_{(\alpha,0)}$ is a mouse set is perhaps our most “quotable” result, since $A_{(\alpha,0)}$ is a very natural, and easily described set. $A_{(\alpha,0)}$ is the set of reals which are ordinal definable in some level of $L(\mathbb{R})$ below the α^{th} level. For example, with $\alpha = \omega_1$

$$A_{(\omega_1,0)} = \{ x \in \mathbb{R} \mid (\exists \beta < \omega_1) \ x \text{ is ordinal definable in } L_\beta(\mathbb{R}) \}.$$

So in this paper we prove that, assuming there are ω Woodin cardinals, $A_{(\omega_1,0)}$ is a mouse set.

This paper is based in part on my PhD dissertation. I would like to thank my thesis advisor, Professor John Steel, for his guidance.

1. PREREQUISITES

In this section we briefly discuss some material which is a prerequisite for understanding this paper. First we discuss some material from the paper “Scales in $L(\mathbb{R})$ ” by J. R. Steel [St1]. Then we discuss some material from the papers “Fine Structure and Iteration Trees” by W. Mitchell and J. R. Steel [MiSt], and “Inner Models With Many Woodin Cardinals” by J. R. Steel [St2]. We present this material here purely as a service to the reader.

Scales In $L(\mathbb{R})$

All of the definitions and results in this sub-section are taken word-for-word from [St1]. As we shall heavily use this material we reprint it here for the reader’s convenience. We work in the theory $\text{ZF} + \text{DC}$.

Throughout this paper we will use the symbol $\mathbb{R}$ to mean ${}^\omega\omega$ and the word *real* to refer to an element of $\mathbb{R}$.

A *pointclass* is a class of subsets of $\mathbb{R}$ closed under recursive substitution. If Γ is a pointclass then

$$\begin{aligned} \neg\Gamma &= \{ \mathbb{R} - A \mid A \in \Gamma \} = \text{the dual of } \Gamma \\ \exists^{\mathbb{R}}\Gamma &= \{ \exists^{\mathbb{R}}A \mid A \in \Gamma \}; \text{ where } \exists^{\mathbb{R}}A = \{ x \mid \exists y (\langle x, y \rangle \in A) \} \\ \text{and } \forall^{\mathbb{R}}\Gamma &= \{ \forall^{\mathbb{R}}A \mid A \in \Gamma \}; \text{ where } \forall^{\mathbb{R}}A = \{ x \mid \forall y (\langle x, y \rangle \in A) \}. \end{aligned}$$

By $\text{Det}(\Gamma)$ we mean the assertion that all games whose payoff set is in Γ are determined.

Because it is convenient, we shall use the Jensen J-hierarchy for $L(\mathbb{R})$. Let $\text{rud}(M)$ be the closure of $M \cup \{M\}$ under the rudimentary functions. Let HF be the set of hereditarily finite sets. Now let

$$\begin{aligned} J_1(\mathbb{R}) &= \text{HF} \cup \mathbb{R} \\ J_{\alpha+1}(\mathbb{R}) &= \text{rud}(J_\alpha(\mathbb{R})); \quad \text{for } \alpha > 0 \\ \text{and } J_\lambda(\mathbb{R}) &= \bigcup_{\alpha < \lambda} J_\alpha(\mathbb{R}); \quad \text{for } \lambda \text{ a limit ordinal.} \end{aligned}$$

Of course $L(\mathbb{R}) = \bigcup_{\alpha \in \text{OR}} J_\alpha(\mathbb{R})$. For all $\alpha \geq 1$, $J_\alpha(\mathbb{R})$ is transitive. For $1 \leq \alpha < \beta$, $J_\alpha(\mathbb{R}) \in J_\beta(\mathbb{R})$. For $\alpha > 1$, $\text{rank } J_\alpha(\mathbb{R}) = \text{OR} \cap J_\alpha(\mathbb{R}) = \omega\alpha$. Also

$$\begin{aligned} J_{\alpha+1}(\mathbb{R}) \cap \mathcal{P}(J_\alpha(\mathbb{R})) &= \\ \{ x \subseteq J_\alpha(\mathbb{R}) \mid x \text{ is definable over } J_\alpha(\mathbb{R}) \text{ from parameters in } J_\alpha(\mathbb{R}) \}. \end{aligned}$$

It is useful to refine this hierarchy. There is a single rudimentary function S such that for all transitive sets M , $S(M)$ is transitive, $M \in S(M)$, and $\text{rud}(M) = \bigcup_{n \in \omega} S^n(M)$. Now let

$$\begin{aligned} S_\omega(\mathbb{R}) &= J_1(\mathbb{R}) \\ S_{\alpha+1}(\mathbb{R}) &= S(S_\alpha(\mathbb{R})); \quad \text{for } \alpha > \omega \\ \text{and } S_\lambda(\mathbb{R}) &= \bigcup_{\alpha < \lambda} S_\alpha(\mathbb{R}); \quad \text{for } \lambda \text{ a limit ordinal.} \end{aligned}$$

Then for all $\alpha \geq 1$, $J_\alpha(\mathbb{R}) = S_{\omega\alpha}(\mathbb{R})$.

If $\alpha \geq 2$ and $X \subseteq J_\alpha(\mathbb{R})$ then $\Sigma_n(J_\alpha(\mathbb{R}), X)$ is the class of all relations on $J_\alpha(\mathbb{R})$ definable over $(J_\alpha(\mathbb{R}); \in)$ by a Σ_n formula, from parameters in $X \cup \{\mathbb{R}\}$. Thus we are always allowed $\mathbb{R}$ itself (but not necessarily its elements) as a parameter in definitions over $J_\alpha(\mathbb{R})$. $\Sigma_\omega(J_\alpha(\mathbb{R}), X) =_{\text{def}} \bigcup_{n \in \omega} \Sigma_n(J_\alpha(\mathbb{R}), X)$. We write “ $\Sigma_n(J_\alpha(\mathbb{R}))$ ” for “ $\Sigma_n(J_\alpha(\mathbb{R}), \emptyset)$,” and “ $\Sigma_n(J_\alpha(\mathbb{R}))$ ” for “ $\Sigma_n(J_\alpha(\mathbb{R}), J_\alpha(\mathbb{R}))$.” Similar conventions apply to the Π_n and Δ_n notations.

Our first lemma states that for all $\alpha \geq 1$ and all $A \subseteq \mathbb{R}$ if $A \in \Sigma_1(J_{\alpha+1}(\mathbb{R}))$ then $A = \bigcup_{n \in \omega} A_n$ with each $A_n \in \Sigma_\omega(J_\alpha(\mathbb{R}))$.

Lemma 1.1. *There is a recursive function which assigns to each Σ_1 formula φ with 2 free variables, a sequence of formulae $\langle \varphi_n \mid n \in \omega \rangle$ so that for all $\alpha \geq 1$ and all $x \in \mathbb{R}$, $J_{\alpha+1}(\mathbb{R}) \models \varphi[x, \mathbb{R}]$ iff $(\exists n \in \omega) J_\alpha(\mathbb{R}) \models \varphi_n[x, \mathbb{R}]$.*

Proof. This is essentially proved in Lemma 1.11 of [St1]. We have

$$\begin{aligned} J_{\alpha+1}(\mathbb{R}) &\models \varphi[x, \mathbb{R}] \\ \Leftrightarrow (\exists n \in \omega) S_{\omega\alpha+n}(\mathbb{R}) &\models \varphi[x, \mathbb{R}] \\ \Leftrightarrow (\exists n \in \omega) S^n(J_\alpha(\mathbb{R})) &\models \varphi[x, \mathbb{R}]. \end{aligned}$$

Now rudimentary functions are simple and so there are Σ_0 formulae $\langle \theta_n \mid n \in \omega \rangle$ such that for all transitive sets M and all $a, b \in M$

$$\theta_n(M, a, b) \Leftrightarrow S^n(M) \models \varphi[a, b].$$

Finally let φ_n be formulae such that for all transitive sets M and all $a, b \in M$

$$\theta_n(M, a, b) \Leftrightarrow M \models \varphi_n[a, b]. \quad \square$$

The following is Lemma 1.1 of [St1].

Lemma 1.2. *The sequences $\langle J_\beta(\mathbb{R}) \mid \beta < \alpha \rangle$ and $\langle S_\beta(\mathbb{R}) \mid \beta < \alpha \rangle$ are uniformly $\Sigma_1(J_\alpha(\mathbb{R}))$ for $\alpha > 1$.*

If $1 \leq \alpha < \beta$ and $X \subseteq J_\alpha(\mathbb{R})$, then $J_\alpha(\mathbb{R}) \prec_n^X J_\beta(\mathbb{R})$ means that for all $\vec{a} \in (X \cup \{\mathbb{R}\})^{<\omega}$, and all Σ_n formulae φ , $J_\alpha(\mathbb{R}) \models \varphi[\vec{a}]$ iff $J_\beta(\mathbb{R}) \models \varphi[\vec{a}]$. We write “ $J_\alpha(\mathbb{R}) \prec_n J_\beta(\mathbb{R})$ ” for “ $J_\alpha(\mathbb{R}) \prec_n^{J_\alpha(\mathbb{R})} J_\beta(\mathbb{R})$.”

Let θ be the least ordinal not the surjective image of $\mathbb{R}$, and δ_1^2 the least ordinal not the image of $\mathbb{R}$ under a surjection f such that $\{ \langle x, y \rangle \mid f(x) \leq f(y) \}$ is Δ_1^2 . Then $\theta^{L(\mathbb{R})} =$ the least α such that $\mathcal{P}(\mathbb{R}) \cap L(\mathbb{R}) \subseteq J_\alpha(\mathbb{R})$. The following is Lemma 1.2 of [St1].

Lemma 1.3. *Let σ be least such that $J_\sigma(\mathbb{R}) \prec_1^{\mathbb{R}} L(\mathbb{R})$. Then*

- (a) $(\Sigma_1^2)^{L(\mathbb{R})} = \Sigma_1(J_\sigma(\mathbb{R})) \cap \mathcal{P}(\mathbb{R})$
- (b) $(\Delta_1^2)^{L(\mathbb{R})} = J_\sigma(\mathbb{R}) \cap \mathcal{P}(\mathbb{R})$
- (c) $(\delta_1^2)^{L(\mathbb{R})} = \sigma$

Definition 1.4. *Let $\alpha, \beta \in \text{OR}$ with $\alpha \leq \beta$. The interval $[\alpha, \beta]$ is a Σ_1 -gap iff*

- (i) $J_\alpha(\mathbb{R}) \prec_1^{\mathbb{R}} J_\beta(\mathbb{R})$
- (ii) $(\forall \alpha' < \alpha) [J_{\alpha'}(\mathbb{R}) \not\prec_1^{\mathbb{R}} J_\alpha(\mathbb{R})]$
- (iii) $(\forall \beta' > \beta) [J_\beta(\mathbb{R}) \not\prec_1^{\mathbb{R}} J_{\beta'}(\mathbb{R})]$

That is, a Σ_1 -gap is a maximal interval of ordinals in which no new Σ_1 facts about elements of $\mathbb{R} \cup \{\mathbb{R}\}$ are verified. If $[\alpha, \beta]$ is a Σ_1 -gap, we say that α begins the gap and β ends it. Notice that we allow $\alpha = \beta$. We shall also allow $[(\delta_1^2)^{L(\mathbb{R})}, \theta^{L(\mathbb{R})}]$ as a Σ_1 -gap. Notice that if $[\alpha, \beta]$ is a Σ_1 -gap and $\alpha < \beta$ then $J_\alpha(\mathbb{R})$ is admissible. The following is Lemma 2.3 of [St1].

Lemma 1.5. *The Σ_1 -gaps partition $\theta^{L(\mathbb{R})}$.*

The following is Lemma 1.11 of [St1].

Lemma 1.6. *Suppose that $\alpha > 1$ and α begins a Σ_1 -gap. Then*

- (a) *There is a $\Sigma_1(J_\alpha(\mathbb{R}))$ **partial** map $h: \mathbb{R} \xrightarrow{\text{onto}} J_\alpha(\mathbb{R})$.*
- (b) *If $\beta < \alpha$ then there is a **total** map $f: \mathbb{R} \xrightarrow{\text{onto}} J_\beta(\mathbb{R})$ such that $f \in J_\alpha(\mathbb{R})$.*

We conclude with the main theorem from [St1]. The following is Theorem 2.1 from that paper.

Lemma 1.7. *If $\alpha > 1$ and $\text{Det}(J_\alpha(\mathbb{R}))$ holds then the pointclass $\mathcal{P}(\mathbb{R}) \cap \Sigma_1(J_\alpha(\mathbb{R}))$ has the scale property.*

Fine Structure and Iteration Trees

In this subsection we briefly review some of the terms we will use from inner model theory. We would like to give the reader an intuitive understanding of what a “mouse”

is. At the end of this subsection we state the Condensation Theorem, and then we prove a corollary to the Condensation Theorem which will be important to us in the sequel.

As we did for $L(\mathbb{R})$, we also use the Jensen J -hierarchy for models of the form $L[\vec{E}]$, where $\vec{E}$ is a sequence of extenders. Recall that one of the properties of a J -hierarchy implies that for any ordinal α , $\text{OR} \cap J_\alpha^{\vec{E}} = \omega\alpha$.

A *potential premouse* is a model of the form $\mathcal{M} = (M; \in, E^{\mathcal{M}}, F^{\mathcal{M}})$, where M is a transitive set, and $E^{\mathcal{M}}, F^{\mathcal{M}}$ are amenable predicates over M , and for some sequence $\vec{E} = \langle E_\alpha \mid \alpha \in \text{dom}(\vec{E}) \rangle$ we have that:

- (i) $\vec{E}$ is a *good* sequence of extenders, and the domain of $\vec{E}$ is some set of ordinals, and
- (ii) for all $\gamma \in \text{dom}(\vec{E})$, $\gamma = \omega\gamma$, and E_γ is an extender of length γ over the model $J_\gamma^{\vec{E} \restriction \gamma}$, and
- (iii) $M = J_\alpha^{\vec{E} \restriction \alpha}$, for some α , and
- (iv) $E^{\mathcal{M}}$ codes $\vec{E} \restriction \alpha$, and
- (v) $F^{\mathcal{M}}$ codes E_α , if $\alpha \in \text{dom}(\vec{E})$; otherwise $F^{\mathcal{M}} = \emptyset$.

For the definition of a *good* sequence of extenders, see [MiSt]. If $F^{\mathcal{M}} = \emptyset$ then we say that $\mathcal{M}$ is *passive*. Otherwise we say that $\mathcal{M}$ is *active*.

If $\mathcal{M} = (J_\alpha^{\vec{E} \restriction \alpha}; \in, E^{\mathcal{M}}, F^{\mathcal{M}})$ is a potential premouse then, for $\beta \leq \alpha$, $\mathcal{J}_\beta^{\mathcal{M}}$ is the premouse $\mathcal{N} = (J_\beta^{\vec{E} \restriction \beta}; \in, E^{\mathcal{N}}, F^{\mathcal{N}})$, where $E^{\mathcal{N}}$ codes $\vec{E} \restriction \beta$ and $F^{\mathcal{N}}$ codes E_β , if $\beta \in \text{dom}(\vec{E})$, or $F^{\mathcal{N}} = \emptyset$ otherwise. For $\beta \leq \alpha$, $\mathcal{J}_\beta^{\mathcal{M}}$ is called an *initial segment* of $\mathcal{M}$. We write $\mathcal{N} \trianglelefteq \mathcal{M}$ to mean that $\mathcal{N}$ is an initial segment of $\mathcal{M}$. $\mathcal{N} \triangleleft \mathcal{M}$ means that $\mathcal{N}$ is a proper initial segment of $\mathcal{M}$.

We must briefly consider the fine structure of definability over potential premice. A potential premouse $\mathcal{M}$ is *n-sound*, iff $\mathcal{M}$ is the Σ_n hull in $\mathcal{M}$ of $\rho_n(\mathcal{M}) \cup \{p_n(\mathcal{M})\}$, where $\rho_n(\mathcal{M})$ is the Σ_n *projectum* of $\mathcal{M}$, and $p_n(\mathcal{M})$ is the n -th *standard parameter* of $\mathcal{M}$. $\mathcal{M}$ is *sound* iff for all $n \in \omega$, $\mathcal{M}$ is n -sound. (I am being a little sloppy here. See [MiSt] for the real definition of sound. In particular, when talking about definability over potential premice, we do not actually use the Levy hierarchy of Σ_n definability, but rather a quasi-Levy hierarchy called $r\Sigma_n$ definability.)

A *premouse* is a potential premouse, all of whose proper initial segments are sound. From now on, we shall never need to talk about potential premice, only premice. Potential premice are only defined in order to be able to define premice.

Notation 1.8. Let $\mathcal{M} = (M; \in, E^{\mathcal{M}}, F^{\mathcal{M}})$ be a premouse. Then we will use the symbol $\mathcal{M}$ to refer both to the *model* $\mathcal{M}$, and to the *set* M . From the context it will be apparent to which object we are referring.

A premouse $\mathcal{M}$ is *meek*, iff there is no initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that $\mathcal{N}$ is active and, letting κ be the critical point of last extender on the $\mathcal{N}$ sequence, $\mathcal{N}$ satisfies that there are cofinally many Woodin cardinals below κ . See [St2]. Meekness is a property which ensures that a premouse does not satisfy too large of a large cardinal hypothesis. For the sake of this paper, it is enough to know that if no initial segment of $\mathcal{M}$ has ω Woodin cardinals, then $\mathcal{M}$ is meek.

A premouse $\mathcal{M}$ is *iterable* iff one can build iteration trees on $\mathcal{M}$ without encountering illfoundedness. There is more than one way to make this notion precise. One particularly useful way is via the notion of realizability. A premouse $\mathcal{M}$ is *realizable* iff $\mathcal{M}$ can be embedded into another premouse $\bar{\mathcal{M}}$, where $\bar{\mathcal{M}}$ has the property that the extenders on the $\bar{\mathcal{M}}$ sequence all come from extenders over V . See [St2] for the precise definition of realizable. In [St2] it is shown that if a countable, meek premouse $\mathcal{M}$ is realizable, then $\mathcal{M}$ is iterable enough to do whatever we want with it. In particular, if $\mathcal{M}$ and $\mathcal{N}$ are two countable, meek, realizable premice, then $\mathcal{M}$ and $\mathcal{N}$ can be compared.

A *mouse* is an iterable premouse. Ironically, since there is not a single, universally useful notion of iterable, the term mouse is never used. In our setting, we could say that a mouse is a realizable, meek premouse.

A premouse is an “ L -like” model. If $\mathcal{M}$ is a premouse then $\mathcal{M}$ is a model of GCH, and in fact, $\mathcal{M}$ is strongly acceptable. Thus if κ is a cardinal of $\mathcal{M}$ and $x \in J_{\kappa}^{\mathcal{M}}$, then $\mathcal{P}(x) \cap \mathcal{M} \subset J_{\kappa}^{\mathcal{M}}$. Another way in which a premouse $\mathcal{M}$ is like L is that $\mathcal{M}$ satisfies the following Condensation Theorem.

Theorem 1.9 (The Condensation Theorem). *Let $\mathcal{H}$ and $\mathcal{M}$ be countable, realizable, meek premice. Let $n \geq 0$ and suppose that $\mathcal{H}$ and $\mathcal{M}$ are both $n + 1$ -sound. Suppose there is a*

map $\pi : \mathcal{H} \rightarrow \mathcal{M}$ such that π is $r\Sigma_{n+1}$ elementary, and $\text{crit}(\pi) \geq \rho_{n+1}(\mathcal{H})$. Then either

(a) $\mathcal{H}$ is a proper initial segment of $\mathcal{M}$

or

(b) There is an extender E on the $\mathcal{M}$ sequence such that $\text{lh}(E) = \rho_{n+1}(\mathcal{H})$, and $\mathcal{H}$ is a proper initial segment of $\text{Ult}_0(\mathcal{M}, E)$.

The Condensation Theorem, which is due to Mitchell and Steel, is proved in the union of [MiSt] and [St2]. See in particular Theorem 8.2 of [MiSt].

We conclude this section with two corollaries to the Condensation Theorem. These corollaries concern the $\leq$ -least initial segment $\mathcal{N}$ of some given premouse $\mathcal{M}$ such that $\mathcal{N}$ satisfies some given fact about some given parameter. Our corollaries imply, for instance, that the $\leq$ -least premouse $\mathcal{N}$ which satisfies a given *sentence* must project to ω . We will use these corollaries several times in the sequel.

Before stating our corollaries we will need one lemma. The following lemma is implicit in section 2 of [MiSt]. See [MiSt] for a definition of the fine structural terms used in this lemma.

Lemma 1.10. *Let $\mathcal{M}$ be an n -sound premouse. Let $\mathcal{H} = \mathcal{H}_{n+1}^{\mathcal{M}}(X)$ where $X \subseteq \mathcal{M}$ and $u_n(\mathcal{M}) \in X$. Let $\pi : \mathcal{H} \rightarrow \mathcal{M}$ be the inverse of the collapse. Then π is a weak n -embedding and π is $r\Sigma_{n+1}$ elementary.*

In the above lemma, the fact that π is a weak n -embedding means that:

- (a) π is $r\Sigma_{n+1}$ elementary.
- (b) For $1 \leq k \leq n-1$, $\pi(\rho_k^{\mathcal{H}}) = \rho_k^{\mathcal{M}}$.
- (c) $\rho_n^{\mathcal{H}}$ = the least α such that $\pi(\alpha) \geq \rho_n^{\mathcal{M}}$.
- (d) $\mathcal{H}$ is n -sound, and
- (e) $\pi(p_k(\mathcal{H})) = p_k(\mathcal{M})$ for $k \leq n$.

Now we state our first corollary to the Condensation Theorem.

Corollary 1.11. *Let $\mathcal{M}$ be a countable, realizable, meek premouse. Let δ be a cardinal of $\mathcal{M}$, and let $q \in J_\delta^{\mathcal{M}}$. Let $n \geq 0$ and let φ be an $r\Sigma_{n+1}$ formula with 1 free variable. Suppose there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\delta \in \mathcal{N}$ and $\mathcal{N} \models \varphi[q]$. Let $\mathcal{N}$ be the $\leq$ -least such initial segment. Then $\rho_{n+1}(\mathcal{N}) = \delta$.*

Proof. Let $\mathcal{H} = \mathcal{H}_{n+1}^{\mathcal{N}}(\delta \cup \{u_n(\mathcal{N})\})$ and let $\pi : \mathcal{H} \rightarrow \mathcal{N}$ be the inverse of the collapse map. By Lemma 1.10 π is a weak n -embedding and π is $r\Sigma_{n+1}$ elementary. Thus $\mathcal{H}$ is n -sound and $\pi(u_n(\mathcal{H})) = u_n(\mathcal{N})$. So $\mathcal{H} = \mathcal{H}_{n+1}^{\mathcal{H}}(\delta \cup \{u_n(\mathcal{H})\})$. As δ is a cardinal of $\mathcal{M}$, $\rho_{n+1}^{\mathcal{H}} = \delta$, and so $p_{n+1}(\mathcal{H}) = \langle \emptyset, u_n(\mathcal{H}) \rangle$ and $\mathcal{H}$ is $n+1$ -sound, and $\rho_{n+1}(\mathcal{H}) = \delta \leq \text{crit } \pi$. By the Condensation Theorem $\mathcal{H} \trianglelefteq \mathcal{N}$. (Note that alternative (b) of the Condensation Theorem cannot hold since $\delta = \rho_{n+1}(\mathcal{H})$ is a cardinal of $\mathcal{M}$, and so $\mathcal{J}_\delta^{\mathcal{M}}$ is passive.) Now consider two cases.

Case 1. $\delta \in H$.

Then as $\mathcal{H} \models \varphi[q]$, we have $\mathcal{H} = \mathcal{N}$ and so we are done.

Case 2. $\mathcal{H} = \mathcal{J}_\delta^{\mathcal{N}}$.

In this case we must have $u_n(\mathcal{N}) = \emptyset$. Let $\mathcal{H}' = \mathcal{H}_{n+1}^{\mathcal{N}}(\delta \cup \{\delta\})$ and let $\pi' : \mathcal{H}' \rightarrow \mathcal{N}$ be the inverse of the collapse map. By Lemma 1.10, π' is a weak n -embedding and π' is $r\Sigma_{n+1}$ elementary. Thus $\mathcal{H}'$ is n -sound and $u_n(\mathcal{H}') = \emptyset$. As $\mathcal{H}' = \mathcal{H}_{n+1}^{\mathcal{H}'}(\delta \cup \{\delta\})$ and δ is a cardinal of $\mathcal{M}$, $\rho_{n+1}^{\mathcal{H}'} = \delta$.

Claim. $\langle \delta \rangle$ is $(n+1)$ -solid over $(\mathcal{H}', \emptyset)$.

Proof of Claim. $Th_{n+1}^{\mathcal{H}'}(\delta) = Th_{n+1}^{\mathcal{N}}(\delta) = Th_{n+1}^{\mathcal{H}}(\delta)$. But $\mathcal{H} = \mathcal{J}_\delta^{\mathcal{N}} = \mathcal{J}_\delta^{\mathcal{H}'}$. So $Th_{n+1}^{\mathcal{H}'}(\delta) \in \mathcal{H}'$. $\diamond$

Thus $p_{n+1}(\mathcal{H}') = \langle \delta, \emptyset \rangle$ and $\mathcal{H}'$ is $(n+1)$ -sound and $\rho_{n+1}(\mathcal{H}') = \delta \leq \text{crit } \pi'$. So we may apply the Condensation Theorem and conclude that $\mathcal{H}' \trianglelefteq \mathcal{N}$. As $\mathcal{H}' \models \varphi[q]$, $\mathcal{H}' = \mathcal{N}$. $\square$

Let $\mathcal{M}$, δ , q , and φ be as in the previous corollary. Now let $\mathcal{N}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ such that $\mathcal{N} \models \varphi[q]$. By analogy with the behavior of the model L , we would expect that $\text{OR}^{\mathcal{N}} < \delta$. Using the previous Corollary we can see only that $\text{OR}^{\mathcal{N}} < (\delta^+)^{\mathcal{M}}$. It is almost certainly true that $\text{OR}^{\mathcal{N}} < \delta$, but we are unable to see how to show this, simply by quoting the *statement* of the Condensation Theorem. A proof that $\text{OR}^{\mathcal{N}} < \delta$ can probably be given using an argument similar to the *proof* of the Condensation Theorem, that is, by using a comparison argument. It would take us too far afield to do such a comparison argument here, and besides we do not need the result in full generality. Below we give a short argument which only uses the *statement* of the

Condensation Theorem, in the special case that $\delta = \omega_1^{\mathcal{M}}$. What follows is our second corollary to the Condensation Theorem.

Corollary 1.12. *Let $\mathcal{M}$ be a countable, realizable, meek premouse. Suppose $\mathcal{M} \models \text{“}\omega_1 \text{ exists,“}$ and let $\delta = \omega_1^{\mathcal{M}}$. Let $q \in J_\delta^{\mathcal{M}}$. Let φ be any formula with 1 free variable. Suppose there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N} \models \varphi[q]$. Let $\mathcal{N}$ be the $\trianglelefteq$ -least such initial segment. Then $\text{OR}^{\mathcal{N}} < \delta$.*

Proof. Suppose that $\text{OR}^{\mathcal{N}} \geq \delta$, and we will derive a contradiction. If $\text{OR}^{\mathcal{N}} > \delta$, then by the previous corollary $\rho_\omega(\mathcal{N}) = \delta$. If $\text{OR}^{\mathcal{N}} = \delta$ then trivially $\rho_\omega(\mathcal{N}) = \delta$. Let $n \in \omega$ be large enough so that $\rho_n(\mathcal{N}) = \delta$, and also so that φ is an $r\Sigma_n$ formula.

Fix $\xi < \delta$ such that $q \in J_\xi^{\mathcal{M}}$ and $\rho_\omega(\mathcal{J}_\xi^{\mathcal{M}}) = \omega$. Let $\mathcal{H} = \mathcal{H}_{n+5}^{\mathcal{N}}(\xi + 1 \cup \{\delta, u_{n+5}(\mathcal{N})\})$, and let $\pi : \mathcal{H} \rightarrow \mathcal{N}$ be the inverse of the collapse map. Let $\kappa = \text{crit}(\pi)$. A standard argument show that $\text{ran}(\pi) \cap \delta$ is transitive, and so $\pi(\kappa) = \delta$. By Lemma 1.10 π is a weak $(n+4)$ -embedding and π is $r\Sigma_{n+5}$ elementary. Since $\pi(q) = q$, we have that $\mathcal{H} \models \varphi[q]$. Also, $\mathcal{H}$ is n -sound and $\rho_n(\mathcal{H}) = \kappa$. Thus we may apply the Condensation Theorem to conclude that either

(a) $\mathcal{H}$ is a proper initial segment of $\mathcal{M}$

or

(b) There is an extender E on the $\mathcal{M}$ sequence such that $\text{lh}(E) = \kappa$, and $\mathcal{H}$ is a proper initial segment of $\text{Ult}_0(\mathcal{M}, E)$.

In case (a) we have that $\text{OR}^{\mathcal{H}} < \delta$, and this contradicts our assumption that $\mathcal{N}$ was the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ such that $\mathcal{N} \models \varphi[q]$. So suppose that case (b) occurs.

Fix an extender E as in (b). Let $\mathcal{P} = \mathcal{J}_\kappa^{\mathcal{M}}$. Let $i = i_E^{\mathcal{P}}$. Thus $i : \mathcal{P} \rightarrow \text{Ult}_0(\mathcal{P}, E)$. Now $\mathcal{H}$ is a proper initial segment of $\text{Ult}_0(\mathcal{P}, E)$, and so $\text{Ult}_0(\mathcal{P}, E)$ satisfies the statement: “There is an initial segment $\mathcal{Q}$ of me such that $\mathcal{Q} \models \varphi[q]$.” Let $\mu = \text{crit}(E) = \text{crit}(i)$. Since $\rho_\omega(\mathcal{J}_\xi^{\mathcal{M}}) = \omega$ and μ is a cardinal in $\mathcal{P}$, we have that $\mu \geq \xi$. Thus $i(q) = q$. It follows that $\mathcal{P}$ satisfies the statement: “There is an initial segment $\mathcal{Q}$ of me such that $\mathcal{Q} \models \varphi[q]$.” Again, this contradicts our assumption that $\mathcal{N}$ was the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ such that $\mathcal{N} \models \varphi[q]$. $\square$

2. SOME LIGHTFACE POINTCLASSES IN $L(\mathbb{R})$

In this section we work in the theory $\text{ZF} + \text{DC}$. For each ordinal $\alpha \geq 2$, we now define an analytical-hierarchy-like sequence of lightface pointclasses.

Definition 2.1. For $\alpha \geq 2$, let

$$\Sigma_{(\alpha,0)} = \mathcal{P}(\mathbb{R}) \cap \Sigma_1(J_\alpha(\mathbb{R}))$$

$$\text{and for } n \geq 0 \text{ let } \Pi_{(\alpha,n)} = \neg \Sigma_{(\alpha,n)}$$

$$\Delta_{(\alpha,n)} = \Sigma_{(\alpha,n)} \cap \Pi_{(\alpha,n)}$$

$$\Sigma_{(\alpha,n+1)} = \exists^{\mathbb{R}} \Pi_{(\alpha,n)}$$

So a set of reals A is in the pointclass $\Sigma_{(\alpha,n)}$, iff A is definable over $J_\alpha(\mathbb{R})$ using a formula φ with a certain restricted syntax. We would like to give a name to this syntax. We will say that a formula φ is $\Sigma_{(\bullet,n)}$, iff φ is the type of formula which is used to define a $\Sigma_{(\alpha,n)}$ set. More specifically, let $\mathcal{L}_{\mathbb{R}}$ be the language with the constant symbol $\dot{\mathbb{R}}$ in addition to the binary relation symbol $\in$. A formula in $\mathcal{L}_{\mathbb{R}}$ is interpreted in the model $(J_\alpha(\mathbb{R}); \in)$ by interpreting $\dot{\mathbb{R}}$ as $\mathbb{R}$. A formula φ is $\Sigma_{(\bullet,0)}$ iff φ is a Σ_1 formula of $\mathcal{L}_{\mathbb{R}}$. A formula φ is $\Pi_{(\bullet,0)}$ iff φ is a Π_1 formula of $\mathcal{L}_{\mathbb{R}}$. If $n \geq 1$, then φ is $\Sigma_{(\bullet,n)}$ iff there is a $\Pi_{(\bullet,n-1)}$ formula ψ such that $\varphi = (\exists v \in \dot{\mathbb{R}}) \psi$, and φ is $\Pi_{(\bullet,n)}$ iff there is a $\Sigma_{(\bullet,n-1)}$ formula ψ such that $\varphi = (\forall v \in \dot{\mathbb{R}}) \psi$. Thus, a set of reals A is in $\Sigma_{(\alpha,n)}$ just in case

$$A = \{ x \in \mathbb{R} \mid J_\alpha(\mathbb{R}) \models \varphi[x] \}$$

for some $\Sigma_{(\bullet,n)}$ formula φ .

If $x \in \mathbb{R}$ then the relativized pointclasses $\Sigma_{(\alpha,n)}(x)$, $\Pi_{(\alpha,n)}(x)$, $\Delta_{(\alpha,n)}(x)$ are defined as usual. Set $\tilde{\Sigma}_{(\alpha,n)} = \bigcup_{x \in \mathbb{R}} \Sigma_{(\alpha,n)}(x)$. Similarly for $\tilde{\Pi}_{(\alpha,n)}$. Set $\tilde{\Delta}_{(\alpha,n)} = \tilde{\Sigma}_{(\alpha,n)} \cap \tilde{\Pi}_{(\alpha,n)}$. For completeness we also set $\Sigma_{(1,n)} = \Sigma_n^1$, etc.

Since a $\Sigma_{(\bullet,n)}$ formula is, in particular, a Σ_{n+1} formula, $\Sigma_{(\alpha,n)} \subseteq \Sigma_{n+1}(J_\alpha(\mathbb{R}))$. But, in general, $\Sigma_{(\alpha,n)} \neq \Sigma_{n+1}(J_\alpha(\mathbb{R}))$. The reason we are interested in the pointclasses $\Sigma_{(\alpha,n)}$ instead of just $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$, is that the property of being a *scaled* pointclass is propagated by the quantifiers $\exists^{\mathbb{R}}$ and $\forall^{\mathbb{R}}$. It is a little annoying that our definition yields that $\Sigma_{(\alpha,n)} \subseteq \Sigma_{n+1}(J_\alpha(\mathbb{R}))$ instead of $\Sigma_{(\alpha,n)} \subseteq \Sigma_n(J_\alpha(\mathbb{R}))$. The reason we made our definition

the way we did is so that, as in the analytical hierarchy, it is the *even* Σ pointclasses which are scaled. This emphasizes the analogy with the pointclasses of the analytical hierarchy.

We are interested in the pointclasses $\Sigma_{(\alpha,n)}$ only for certain ordinals α . If $[\alpha, \beta]$ is a Σ_1 -gap and $\alpha < \gamma \leq \beta$, then for all $x \in \mathbb{R}$ and all $n \in \omega$, $\Sigma_{(\gamma,n)}(x) = \Sigma_{(\alpha,n)}(x)$. So we are only interested in the pointclass $\Sigma_{(\alpha,n)}$ in the case that α begins a Σ_1 -gap. Furthermore, if α begins a Σ_1 -gap and $J_\alpha(\mathbb{R})$ is admissible, then $\Sigma_1(J_\alpha(\mathbb{R}))$ is closed under the real quantifiers $\forall^{\mathbb{R}}$ and $\exists^{\mathbb{R}}$. So $\forall x \in \mathbb{R}$

$$\Sigma_{(\alpha,0)}(x) = \Pi_{(\alpha,1)}(x) = \Sigma_{(\alpha,2)}(x) = \dots$$

$$\text{and} \quad \Pi_{(\alpha,0)}(x) = \Sigma_{(\alpha,1)}(x) = \Pi_{(\alpha,2)}(x) = \dots$$

So the collection of pointclasses $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$ is not particularly interesting. This leads us to the following.

Definition 2.2. An ordinal α is *projective-like* iff α begins a Σ_1 -gap and $J_\alpha(\mathbb{R})$ is not admissible.

Remark 2.3. α projective-like $\Rightarrow [\alpha, \alpha]$ is a Σ_1 -gap.

We are only interested in the pointclasses $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$ in the case that α is projective-like. In Proposition 2.10 below we show that if α is projective-like then the collection of pointclasses $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$ does *not* collapse to two pointclasses as it does in the case that α is admissible. The importance of being projective-like is given by the following lemma.

Lemma 2.4. Let $\alpha \geq 2$ be projective-like. Then $\exists x \in \mathbb{R}$ such that there is a total function $f : \mathbb{R} \rightarrow \omega\alpha = \text{OR} \cap J_\alpha(\mathbb{R})$ with $\text{ran}(f)$ cofinal in $\omega\alpha$ and with $f \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$.

Proof. By definition of non-admissible $\exists \beta < \alpha$ and $\exists g : J_\beta(\mathbb{R}) \xrightarrow{\text{cof}} \omega\alpha$ such that $g \in \Sigma_1(J_\alpha(\mathbb{R}))$. By Lemma 1.6(b) $\exists h \in J_\alpha(\mathbb{R})(h : \mathbb{R} \xrightarrow{\text{onto}} J_\beta(\mathbb{R}))$. Let $f = g \circ h$. Then $\text{ran}(f)$ is cofinal in $\omega\alpha$ and $f \in \Sigma_1(J_\alpha(\mathbb{R}))$. By lemma 1.6(a) we may take the parameter in the definition of f over $J_\alpha(\mathbb{R})$ to be a real. $\square$

Definition 2.5. Let $\alpha \geq 2$ be projective-like. A real x as in Lemma 2.4 above is called a *good parameter* for $J_\alpha(\mathbb{R})$.

Let $\alpha \geq 2$ be projective-like. The collection of pointclasses $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$ has many of the properties of the analytical hierarchy $\{ \Sigma_n^1, \Pi_n^1 \mid n \in \omega \}$. However there is one very important property of the analytical hierarchy which may fail for the collection $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$. This latter collection *may not form a hierarchy*. That is

Warning. For some projective-like α it is not the case that $\Sigma_{(\alpha,n)} \subseteq \Delta_{(\alpha,n+1)}$.

Of course it is true that $\Sigma_{(\alpha,n)} \subseteq \Pi_{(\alpha,n+1)} \subseteq \Sigma_{(\alpha,n+2)}$. Also we do have the following, which explains why we are interested in good parameters: Let $\alpha \geq 2$ be projective-like. Let x be a good parameter for $J_\alpha(\mathbb{R})$. Then $(\forall n) \quad \Sigma_{(\alpha,n)}(x) \subseteq \Delta_{(\alpha,n+1)}(x)$. That is, relative to a good parameter, our pointclasses do form a hierarchy. In fact, we have the following.

Lemma 2.6. *Let $\alpha \geq 2$ be projective-like, and let $n \geq 0$. Let x be a good parameter for $J_\alpha(\mathbb{R})$. Then $\Sigma_{(\alpha,n)}(x) = \mathcal{P}(\mathbb{R}) \cap \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x\})$.*

Proof. This is essentially what is shown in Lemma 2.5 of [St1]. So this proof is due to John Steel. We argue by induction on n . For $n = 0$ the lemma is true by definition of $\Sigma_{(\alpha,0)}$. Now let $n \geq 1$. Even without the parameter x we already have that that $\Sigma_{(\alpha,n)} \subseteq \Sigma_{n+1}(J_\alpha(\mathbb{R}))$. So certainly we have $\Sigma_{(\alpha,n)}(x) \subseteq \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x\})$. For the other direction, by induction, it suffices to see that $\Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x\}) = \exists^\mathbb{R} \Pi_n(J_\alpha(\mathbb{R}), \{x\})$. Let $h: \mathbb{R} \xrightarrow{\text{onto}} J_\alpha(\mathbb{R})$ be a partial $\Sigma_1(J_\alpha(\mathbb{R}))$ map. There is such an h since α begins a Σ_1 -gap. Let R be any $\Pi_n(J_\alpha(\mathbb{R}))$ relation. Then

$$(\exists v \in J_\alpha(\mathbb{R})) R(v)$$

iff

$$(\exists y \in \mathbb{R}) [y \in \text{dom } h \wedge (\forall v \in J_\alpha(\mathbb{R})) (v = h(y) \implies R(v))].$$

If $n \geq 2$ then “ $y \in \text{dom } h$ ” is $\Pi_n(J_\alpha(\mathbb{R}))$ and we are done. If $n = 1$ then we must see that the relation “ $y \in \text{dom } h$ ” is $\Sigma_{(\alpha,1)}(x)$. We have that the relation “ $y \in \text{dom } h$ ” is $\Sigma_{(\alpha,0)}$. So it suffices to prove the following:

Claim. $\Sigma_{(\alpha,0)} \subseteq \Sigma_{(\alpha,1)}(x)$.

Proof of Claim. Let $f : \mathbb{R} \xrightarrow{\text{cof}} \omega\alpha$ be a total $\Sigma_1(J_\alpha(\mathbb{R}), \{x\})$ map. Let R be any $\Sigma_{(\alpha,0)}$ set. So there is a Σ_1 formula φ of $\mathcal{L}_{\mathbb{R}}$ so that $R(y) \Leftrightarrow J_\alpha(\mathbb{R}) \models \varphi[y]$

$$\Leftrightarrow (\exists \beta < \omega\alpha) S_\beta(\mathbb{R}) \models \varphi[y]$$

$$\Leftrightarrow (\exists z \in \mathbb{R}) S_{f(z)}(\mathbb{R}) \models \varphi[y]$$

$$\Leftrightarrow (\exists z \in \mathbb{R})(\forall \beta < \omega\alpha)(\forall S \in J_\alpha(\mathbb{R}))[(\beta = f(z) \wedge S = S_\beta(\mathbb{R})) \implies S \models \varphi[y]].$$

So $R \in \exists^{\mathbb{R}}\Pi_1(J_\alpha(\mathbb{R}), \{x\}) = \Sigma_{(\alpha,1)}(x)$. ◇

This completes the proof of the lemma. □

Remark 2.7. In particular, if $\alpha \geq 2$ is projective like, then $\Sigma_{(\alpha,n)}$ is simply the pointclass $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$.

If 0 is a good parameter for $J_\alpha(\mathbb{R})$, then for all n , $\Sigma_{(\alpha,n)} \subseteq \Delta_{(\alpha,n+1)}$, and so we would expect the $\Sigma_{(\alpha,n)}$ pointclasses to behave exactly like the analytical pointclasses. More generally, if x is a good parameter for $J_\alpha(\mathbb{R})$, then we would expect the pointclasses $\Sigma_{(\alpha,n)}(x)$ to behave exactly like the analytical pointclasses. But we want to consider the *lightface* pointclasses $\Sigma_{(\alpha,n)}$, even when 0 is not a good parameter for $J_\alpha(\mathbb{R})$. In this situation, even though $\Sigma_{(\alpha,n)} \subseteq \Delta_{(\alpha,n+1)}$ may fail, many of the proofs of various well-known properties of the analytical hierarchy go through in our setting. Below we record some of these properties. We begin with some trivial observations.

Lemma 2.8. *Let $\alpha \geq 2$ be projective-like.*

- (1) *Every analytical pointset is in $\Delta_{(\alpha,0)}$.*
- (2) *For all $n \geq 0$, $\Sigma_{(\alpha,n)}$ and $\Pi_{(\alpha,n)}$ are **adequate**. That is, they are closed under recursive substitution, $\wedge$, $\vee$, $\exists^{\leq}$, and $\forall^{\leq}$. (See [Mo] pg. 158.)*
- (3) *For all $n \geq 0$, $\Sigma_{(\alpha,n)}$ and $\Pi_{(\alpha,n)}$ are ω -parameterized.*
- (4) *For all $n \geq 0$, $\Sigma_{(\alpha,n)}$ is closed under $\exists^{\mathbb{R}}$ and $\Pi_{(\alpha,n)}$ is closed under $\forall^{\mathbb{R}}$.*
- (5) *For all $n \geq 1$, $\Sigma_{(\alpha,n)}$ and $\Pi_{(\alpha,n)}$ are closed under both integer quantifiers: $\exists^\omega$, $\forall^\omega$.*

Remark 2.9. Notice that (5) may fail for $n = 0$ if α is a successor ordinal or $\text{cof}(\alpha) = \omega$.

Proof. These are all easy. For example, (3) holds for $n = 0$ because the Σ_1 -satisfaction relation over $J_\alpha(\mathbb{R})$ is $\Sigma_1(J_\alpha(\mathbb{R}))$. Then (3) holds for $n > 0$ by induction. $\square$

Now we can see that if α is projective-like then the collection of pointclasses $\{ \Sigma_{(\alpha,n)}, \Pi_{(\alpha,n)} \mid n \in \omega \}$ does not collapse to two pointclasses as it does in the case that α is admissible.

Proposition 2.10. *Let $\alpha \geq 2$ be projective-like. Then $(\forall n \in \omega) \quad \Sigma_{(\alpha,n+1)} \neq \Pi_{(\alpha,n)}$.*

Proof. Suppose for some n that $\Sigma_{(\alpha,n+1)} = \Pi_{(\alpha,n)}$. Then for all $x \in \mathbb{R}$, $\Sigma_{(\alpha,n+1)}(x) = \Pi_{(\alpha,n)}(x)$. But then for x a good parameter for $J_\alpha(\mathbb{R})$, $\Sigma_{(\alpha,n)}(x) \subseteq \Pi_{(\alpha,n)}(x)$. But this is impossible as there is a universal $\Sigma_{(\alpha,n)}$ set. $\square$

Lemma 2.11 (Second Periodicity Theorem). *Let $\alpha \geq 2$ be projective-like. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \in \omega$. Then $\Sigma_{(\alpha,2n)}$ and $\Pi_{(\alpha,2n+1)}$ have the scale property.*

Proof. This follows from Lemma 1.7, Lemma 2.8, and Theorems 6C.2 and 6C.3 of [Mo]. $\square$

Remark 2.12. In particular, under the hypotheses of Lemma 2.11, $\Pi_{(\alpha,2n+1)}$ and $\Sigma_{(\alpha,2n+2)}$ are *Spector* pointclasses. (See section 4C of [Mo].)

One very important consequence of the scale property is the basis property. The next lemma says that our even Σ pointclasses have the basis property.

Lemma 2.13 (The Basis Property). *Let $\alpha \geq 2$ be projective-like. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \geq 2$ be even. Suppose $A \subset \mathbb{R}$ and $A \in \Sigma_{(\alpha,n)}$ and A is not empty. Then there is an x in A such that $\{x\} \in \Sigma_{(\alpha,n)}$.*

Proof. This follows from 4E.7 in [Mo]. $\square$

Definition 2.14. Let $\alpha \geq 2$ be projective-like, and let $n \geq 0$. Let $x, y \in \mathbb{R}$. Then we say that $x \in \Delta_{(\alpha,n)}(y)$ iff $\{x\} \in \Delta_{(\alpha,n)}(y)$, equivalently iff $\{x\} \in \Sigma_{(\alpha,n)}(y)$. We say that $x \in \Delta_{(\alpha,n)}$ iff $x \in \Delta_{(\alpha,n)}(0)$. Let $A \subseteq \mathbb{R}$. Then we say that A is *closed downward under* $\Delta_{(\alpha,n)}$ iff whenever $x \in A$ and $y \in \Delta_{(\alpha,n)}(x)$ we have that $y \in A$.

Remark 2.15. If $n \geq 1$ then $x \in \Delta_{(\alpha,n)}(y)$ iff $\{ (i, j) \mid x(i) = j \} \in \Delta_{(\alpha,n)}(y)$. For $n = 0$, since $\Sigma_{(\alpha,0)}$ is not closed under $\forall^\omega$ for some α , this latter condition is not equivalent to $x \in \Delta_{(\alpha,0)}(y)$.

Another consequence of the scale property is the bounded quantification property. This is explained in the next lemma.

Lemma 2.16 (Bounded Quantification). *Let $\alpha \geq 2$ be projective-like. Let $n \geq 1$ be odd. Assume $\text{Det}(\Pi_{(\alpha,n)})$. Let $Q(x, y, z) \in \Pi_{(\alpha,n)}$. Set*

$$P(x, y) \Leftrightarrow \exists z \in \Delta_{(\alpha,n)}(y) Q(x, y, z).$$

Then P is $\Pi_{(\alpha,n)}$.

Proof. This follows from 4D.3 of [Mo]. □

Another consequence of the scale property is the following result due to Kechris. Recall that WO is the set of reals which code wellorders of ω , and hence code countable ordinals. If $w \in \text{WO}$ then $|w|$ is the countable ordinal coded by w .

Lemma 2.17 (Boundedness Theorem for Subsets of ω_1). *Let $\alpha \geq 2$ be projective-like. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \geq 1$ be odd. Let $A \subset \text{WO}$ be $\Sigma_{(\alpha,n)}$ and suppose that $\sup \{ |x| : x \in A \} < \omega_1$. Then there exists a $w \in \text{WO}$ such that $w \in \Delta_{(\alpha,n)}$ and such that $|x| < |w|$ for all $x \in A$.*

Proof. The proof of Theorem 1.1 from [Ke2] goes through here. □

The study of the $\Sigma_{(\alpha,n)}$ pointclasses, and in particular the comparison of these pointclasses with the pointclasses of the analytical hierarchy, is an interesting endeavor in its own right. We pursue this study in the papers [Ru1] and [Ru2]. In this paper however, descriptive set theory is not our primary focus. So we will discuss only those topics from descriptive set theory which will be needed in this paper. In addition to the several results we mentioned above, there are two other topics which we will discuss: the topic of countable sets of ordinal definable reals, and the topic of the *correctness* of a given set of reals.

Countable Sets of Ordinal Definable Reals

As Remark 2.15 indicates, when considering the property:

$$x \in \Delta_{(\alpha,n)}(y)$$

the case $n = 0$ is somewhat different than the case $n \geq 1$. Notice that the next two definitions are only for $n \geq 1$.

Definition 2.18. Let $x \in \mathbb{R}$, $\xi < \omega_1$. Let $n \geq 1$. Then we say that $x \in \Delta_{(\alpha,n)}(\xi)$ iff $\forall w \in \text{WO}$ such that $|w| = \xi$, $x \in \Delta_{(\alpha,n)}(w)$.

Definition 2.19. Let $\alpha \geq 2$ be projective-like. Let $n \geq 1$. Then

$$A_{(\alpha,n)} =_{\text{def}} \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in \Delta_{(\alpha,n)}(\xi) \}.$$

Assuming $\text{Det}(J_{\alpha+1}(\mathbb{R}))$, the sets $A_{(\alpha,n)}$ for $n \geq 1$ have much descriptive set-theoretic interest. For even $n \geq 2$, $A_{(\alpha,n)} = C_{(\alpha,n)}$, where $C_{(\alpha,n)}$ is the largest countable $\Sigma_{(\alpha,n)}$ set. For odd $n \geq 1$, $A_{(\alpha,n)} = Q_{(\alpha,n)}$, where $Q_{(\alpha,n)}$ is the largest countable $\Pi_{(\alpha,n)}$ set which is closed down under $\Delta_{(\alpha,n)}$. Even though $\Sigma_{(\alpha,n)} \subset \Delta_{(\alpha,n+1)}$ may fail, the sets $C_{(\alpha,n)}$ and $Q_{(\alpha,n)}$ still share most of the properties of the analogous sets C_n and Q_n in the theory of the analytical hierarchy. For example, for $n \geq 1$ there is a $\Delta_{(\alpha,n)}$ -good wellorder of $A_{(\alpha,n)}$. We will not go into any more detail here about the descriptive set-theoretic properties of the sets $A_{(\alpha,n)}$, for $n \geq 1$. But see [Ru1] and [Ru2] where these sets are studied in detail.

We wish now to discuss the basic notion of a real x being *ordinal definable* over the model $J_\alpha(\mathbb{R})$. For $n \geq 1$, the set of reals which are Σ_n ordinal definable over $J_\alpha(\mathbb{R})$ has a wellordering which is definable over $J_\alpha(\mathbb{R})$. So assuming $\text{Det}(J_{\alpha+1}(\mathbb{R}))$, this set is countable. Consequently, every real which is ordinal definable over $J_\alpha(\mathbb{R})$ is in fact definable over $J_\alpha(\mathbb{R})$ from a single countable ordinal. This leads us to the following definition.

Definition 2.20. Let $n \geq 1$. $\text{OD}_n^{J_\alpha(\mathbb{R})}$ is the set of reals x such that for some Σ_n formula φ in the language $\mathcal{L}_{\mathbb{R}}$, and some countable ordinal ξ , and every real $w \in \text{WO}$ such that

$|w| = \xi$, we have that x is the unique real x' such that $J_\alpha(\mathbb{R}) \models \varphi[x', w]$. Set

$$\text{OD}^{J_\alpha(\mathbb{R})} = \bigcup_n \text{OD}_n^{J_\alpha(\mathbb{R})}.$$

Let $n \geq 1$, and let us consider the following question. What is the relationship between $A_{(\alpha, n)}$ and $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$? Since $\Delta_{(\alpha, n)} \subseteq \Delta_{n+1}(J_\alpha(\mathbb{R}))$, we expect that $A_{(\alpha, n)} \subseteq \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. Notice however that this fact is not completely immediate, for the following technical reason. Let $x \in A_{(\alpha, n)}$. Then there is some $\xi < \omega_1$ such that $x \in \Delta_{(\alpha, n)}(w)$ for every $w \in \text{WO}$ such that $|w| = \xi$. But the $\Delta_{(\alpha, n)}$ definition of x is allowed to depend on w , so it is not obvious that x is actually definable from ξ . However, assuming $\text{Det}(J_{\alpha+1}(\mathbb{R}))$, we have that $A_{(\alpha, n)} = C_{(\alpha, n)}$ if n is even, and $A_{(\alpha, n)} = Q_{(\alpha, n)}$ if n is odd. This implies that $A_{(\alpha, n)}$ is countable, and $A_{(\alpha, n)}$ has a $\Delta_{(\alpha, n)}$ -good wellordering. This in turn implies that there is a single fixed $\Sigma_{(\bullet, n)}$ formula φ , such that for all $x \in A_{(\alpha, n)}$, there is a countable ordinal ξ such that whenever $w \in \text{WO}$ and $|w| = \xi$, x is the unique real x' such that $J_\alpha(\mathbb{R}) \models \varphi[x', w]$. Since φ is, in particular, a Σ_{n+1} formula, we have that $x \in \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. That is, assuming $\text{Det}(J_{\alpha+1}(\mathbb{R}))$, we have that for $n \geq 1$, $A_{(\alpha, n)} \subseteq \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. We ask the following two questions.

Question. Assume $\text{AD}^{L(\mathbb{R})}$. Let $\alpha \geq 2$ be projective-like. Let $n \geq 1$. Is it true that $A_{(\alpha, n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$?

A courser question is the following.

Question. Assume $\text{AD}^{L(\mathbb{R})}$. Let $\alpha \geq 2$ be projective-like. Is it true that $\bigcup_n A_{(\alpha, n)} = \text{OD}^{J_\alpha(\mathbb{R})}$?

That is to say, if x is ordinal definable over $J_\alpha(\mathbb{R})$, does it follow that x is definable using only quantification over the reals, over $\Sigma_1(J_\alpha(\mathbb{R}))$? We do not know the answer to these questions in full generality.

We do have some partial results along these lines. Notice that if $\beta < \alpha$, then $\text{OD}^{J_\beta(\mathbb{R})} \subseteq \text{OD}^{J_\alpha(\mathbb{R})}$. Notice also that $\text{OD}^{J_\beta(\mathbb{R})} = \text{OD}^{J_\alpha(\mathbb{R})}$ is possible. In fact, assuming $\text{AD}^{L(\mathbb{R})}$, there are only countably many α such that $\text{OD}^{J_\alpha(\mathbb{R})} \neq \bigcup_{\beta < \alpha} \text{OD}^{J_\beta(\mathbb{R})}$. Now, given $A \subseteq \text{OD}^{J_\alpha(\mathbb{R})}$, let us say that A is *non-trivial* iff there is an $x \in A$ such that $x \notin \text{OD}^{J_\beta(\mathbb{R})}$ for any

$\beta < \alpha$. In [Ru1] and [Ru2] we show that if $A_{(\alpha,n)}$ is non-trivial, then $A_{(\alpha,n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. So the first question can be rephrased as: if $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$ is non-trivial, must $A_{(\alpha,n)}$ also be non-trivial. The second, courser question can be rephrased as: if $\text{OD}^{J_\alpha(\mathbb{R})}$ is non-trivial, must $A_{(\alpha,n)}$ also be non-trivial for some n . In [Ru1] and [Ru2] we show that the answer to the second question is yes, in the case that α has uncountable cofinality. We are not able to prove anything more about these questions—using techniques from descriptive set theory. Somewhat surprisingly, we do get some additional information using inner model theory.

In this paper we show that the answer to the second question is yes, for $\alpha \leq \omega_1^{\omega_1}$. Also, in the case that $\text{cof}(\alpha) > \omega$, we show that the answer to the first question is yes. This is done as follows. For $\text{cof}(\alpha) > \omega$ we show that there is a mouse $\mathcal{M}$ such that $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$, and $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha,n)}$. Thus $A_{(\alpha,n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. For $\text{cof}(\alpha) \leq \omega$ we have a gap in our proof. We only get that there is a mouse $\mathcal{M}$ such that $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$, and $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha,n+1)}$. Thus $A_{(\alpha,n)} \subseteq \text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq A_{(\alpha,n+1)}$, and so $\bigcup_n A_{(\alpha,n)} = \text{OD}^{J_\alpha(\mathbb{R})}$.

There is one last point we need to discuss. We would like to extend the definition of $A_{(\alpha,n)}$ to the case $n = 0$.

Definition 2.21. Let $\alpha \geq 2$. Then $A_{(\alpha,0)} = \bigcup_{\beta < \alpha} \text{OD}^{J_\beta(\mathbb{R})}$.

Assuming determinacy, we have the following alternate characterization of $A_{(\alpha,0)}$.

Proposition 2.22. Assume $\text{Det}(\Sigma_1(J_\alpha(\mathbb{R})))$. Then $A_{(\alpha,0)} = \text{OD}_1^{J_\alpha(\mathbb{R})}$.

Proof. Notice that $A_{(\alpha,0)}$ is a $\Sigma_1(J_\alpha(\mathbb{R}))$ set and there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ wellorder of $A_{(\alpha,0)}$. So assuming $\text{Det}(\Sigma_1(J_\alpha(\mathbb{R})))$, $A_{(\alpha,0)}$ is countable. Then, since there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ wellorder of $A_{(\alpha,0)}$, we have that $A_{(\alpha,0)} \subseteq \text{OD}_1^{J_\alpha(\mathbb{R})}$. Conversely, suppose that $x \in \text{OD}_1^{J_\alpha(\mathbb{R})}$. Then it is easy to see that x is ordinal definable in $J_\beta(\mathbb{R})$ for some $\beta < \alpha$. (If α is a successor ordinal, then use Lemma 1.1.) $\square$

So $A_{(\alpha,0)}$ is a $\Sigma_{(\alpha,0)}$ set, and assuming determinacy, it is countable. In analogy with even $n \geq 2$ we might expect that $A_{(\alpha,0)}$ is the largest countable $\Sigma_{(\alpha,0)}$ set. It is pretty easy to see that this is true in the case that α is a limit ordinal. If α is a successor ordinal,

we do not know if this is true in general. In [Ru1] and [Ru2] we study this question and get some partial results.

Now that we have defined $A_{(\alpha,n)}$ for all $n \geq 0$, we are ready to state the main conjecture which motivated this paper.

Definition 2.23. Let $A \subset \mathbb{R}$. Then we say that A is a *mouse set* iff $A = \mathbb{R} \cap \mathcal{M}$ for some countable, realizable, meek premouse $\mathcal{M}$.

Conjecture 2.24. Let $\alpha \geq 2$ be projective-like. Then $A_{(\alpha,n)}$ is a mouse set for all $n \geq 0$.

As we mentioned in the introduction, in this paper we will prove the conjecture in the case $\alpha \leq \omega_1^{\omega_1}$, and either $\text{cof}(\alpha) > \omega$, or $\text{cof}(\alpha) \leq \omega$ and $n = 0$. In Section 3 we begin the process of proving the conjecture by giving the definition of an (α, n) -big premouse. Then in Section 4 we give one half of the proof that $A_{(\alpha,n)}$ is a mouse set: we show that if $\mathcal{M}$ is a countable, iterable premouse, and $\mathcal{M}$ is (α, n) -big, then $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$. Actually, we prove the ostensibly stronger result that $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$. We will then use this ostensibly stronger result to prove that $A_{(\alpha,n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. (See the discussion above.) The proof in Section 4 that $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$ will use that fact that $\mathbb{R} \cap \mathcal{M}$ has a certain *correctness* property. Below we explain this correctness property. It turns out that it is related to our notion of good parameters.

Good Parameters and Correctness

Recall that if α is projective-like, then a good parameter for $J_\alpha(\mathbb{R})$ is a real x such that there is a total function $f : \mathbb{R} \xrightarrow{\text{cof}} \omega\alpha$ which is $\Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. In this section we study good parameters and their relationship to the correctness of a given set of reals. We work in the theory $\text{ZF} + \text{DC}$.

We start with an alternative characterization of good parameters.

Definition 2.25. Let $x \in \mathbb{R}, \alpha \geq 2, n \geq 0$. Then we say that x is an *n -name for α* iff there is a formula θ such that α is the least ordinal so that $J_\alpha(\mathbb{R}) \models \theta[x]$, and θ is a $\Sigma_{(\bullet,n)}$ formula if n is even, or a $\Pi_{(\bullet,n)}$ formula if n is odd.

Proposition 2.26. *Let $\alpha \geq 2$ be projective-like. Let $x \in \mathbb{R}$. Then the following are equivalent:*

- (1) *x is a good parameter for $J_\alpha(\mathbb{R})$.*
- (2) *x is a 1-name for α*

Proof. (1) $\Rightarrow$ (2). Let $g : \mathbb{R} \xrightarrow{\text{cof}} \omega\alpha$ be a total function which is $\Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. Let φ be a Σ_1 formula so that $g(y) = \beta$ iff $J_\alpha(\mathbb{R}) \models \varphi[x, y, \beta]$. Then α is least so that $J_\alpha(\mathbb{R}) \models (\forall y \in \mathbb{R})(\exists \beta)\varphi[x, y, \beta]$.

(2) $\Rightarrow$ (1). Let φ be a Σ_1 formula such that α is least so that $J_\alpha(\mathbb{R}) \models (\forall y \in \mathbb{R})\varphi[x, y]$. Let $g : \mathbb{R} \rightarrow \omega\alpha$ be defined by

$$g(y) = (\mu \beta)[S_\beta(\mathbb{R}) \models \varphi[x, (y)_0]] + (y)_1(0).$$

Then $g \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$ and $\text{ran}(g)$ is cofinal in $\omega\alpha$. □

Good parameters play an important role in the *correctness* of sets of reals. We turn to this topic now. We begin with a definition. If $A \subseteq \mathbb{R}$ is any set of reals, and $\alpha \geq 1$ is any ordinal, then $J_\alpha(A)$ is defined similarly to the way we defined $J_\alpha(\mathbb{R})$ on page 7.

Definition 2.27. Let $\alpha \geq 1, n \geq 1, A \subseteq \mathbb{R}, A \neq \emptyset$. Then we say that A is $\Sigma_n(J_\alpha(\mathbb{R}))$ -*correct* iff there is an $\bar{\alpha} \geq 1$ such that

- (1) $\mathbb{R} \cap J_{\bar{\alpha}}(A) = A$ and
- (2) $\exists j : (J_{\bar{\alpha}}(A); \in, A) \rightarrow (J_\alpha(\mathbb{R}); \in, \mathbb{R})$ such that j is Σ_n elementary.

If Γ is a pointclass and A a pointset then we say that A is a *basis* for Γ iff $B \cap A \neq \emptyset$ for all $B \in \Gamma$.

Lemma 2.28. *Suppose α begins a Σ_1 -gap. Let A be a nonempty set of reals. Then the following are equivalent:*

- (a) *A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.*
- (b) *Letting*

$$H = \{ a \in J_\alpha(\mathbb{R}) \mid (\exists x \in A) \{a\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\}) \}$$

we have that $H \prec_1 J_\alpha(\mathbb{R})$ and $\mathbb{R} \cap H = A$.

- (c) *A is closed under join, and for all $x \in A$, A is a basis for $\Sigma_1(J_\alpha(\mathbb{R}), \{x\})$.*

Proof. Clearly $(b) \Rightarrow (a) \Rightarrow (c)$. We will show that $(c) \Rightarrow (b)$. Let $f: \mathbb{R} \xrightarrow{\text{onto}} J_\alpha(\mathbb{R})$ be a partial $\Sigma_1(J_\alpha(\mathbb{R}))$ map. Let $X = f''A$. Then $X \subseteq H$. But (c) easily implies that $X \prec_1 J_\alpha(\mathbb{R})$. Thus $X = H$ and so $H \prec_1 J_\alpha(\mathbb{R})$. But (c) also easily implies that $\mathbb{R} \cap X = A$. $\square$

Remark 2.29. If α begins a Σ_1 -gap and A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct then the $\bar{\alpha}$ and the j of definition 2.27 are unique. [Proof: Given any such $\bar{\alpha}$ and j let $X = \text{ran } j$. Then $X \prec_1 J_\alpha(\mathbb{R})$ so $H \subseteq X$, where H is as in the lemma. But let $f: \mathbb{R} \xrightarrow{\text{onto}} J_\alpha(\mathbb{R})$ be a partial $\Sigma_1(J_\alpha(\mathbb{R}))$ map. Let $a \in X$. Since $J_\alpha(\mathbb{R}) \models “(\exists x \in \mathbb{R}) f(x) = a”$, there is an x in A such that $f(x) = a$. But then $a \in H$. So $X = H$.] We will use the notation j_A to refer to this unique j .

Corollary 2.30. *Suppose α begins a Σ_1 -gap. Let $n \geq 2$. Let A be a non-empty set of reals. Then the following are equivalent:*

- (a) A is $\Sigma_n(J_\alpha(\mathbb{R}))$ -correct.
- (b) A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct and j_A is Σ_n elementary.
- (c) A is closed under join and $(\forall x \in A) A$ is a basis for $\Sigma_n(J_\alpha(\mathbb{R}), \{x\})$.

Proof. Clearly $(b) \Rightarrow (a) \Rightarrow (c)$. Then $(c) \Rightarrow (b)$ follows from the proof of the previous lemma. $\square$

In the analytical hierarchy, if $n \geq 2$ is even and a non-empty set of reals A is closed under join and closed downward under Δ_n^1 , then A is Σ_n^1 -correct. This is because, for even $n \geq 2$, Σ_n^1 has the basis property. In our setting, for even $n \geq 2$, $\Sigma_{(\alpha, n)}$ has the basis property. (See Lemma 2.13.) However, the next two lemmas show that in our setting a good parameter is also required for correctness. The first lemma handles the case $n = 0$, and the second lemma handles the case $n \geq 2$ even.

Lemma 2.31. *Suppose α begins a Σ_1 -gap. Assume $\text{Det}(J_\alpha(\mathbb{R}))$. Let A be a non-empty set of reals. Suppose A is closed under join and closed downward under $\Delta_{(\alpha, 0)}$.*

- (1) *If α is a limit ordinal then A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.*

- (2) If $\alpha = \beta + 1$ and there is no $x \in A$ such that x is a 0-name for α , then A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.
- (3) If $\alpha = \beta + 1$ and there is an $x \in A$ such that x is a 0-name for α , and β is projective-like, then the following are equivalent:
- (a) A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.
 - (b) A is $\Sigma_\omega(J_\beta(\mathbb{R}))$ -correct.
 - (c) $(\exists z \in A) z$ is a good parameter for $J_\beta(\mathbb{R})$.

Proof.

(1) and (2). Let $x \in A$ and let φ be a Σ_1 formula. Suppose that $J_\alpha(\mathbb{R}) \models (\exists y \in \mathbb{R}) \varphi[x, y]$. Let γ be the least ordinal such that $J_\gamma(\mathbb{R}) \models (\exists y \in \mathbb{R}) \varphi[x, y]$. In either case (1) or case (2) we have that $\gamma < \alpha$. Notice then that $\{\gamma\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. Also γ is projective-like. Let $B = \{y \in \mathbb{R} \mid J_\gamma(\mathbb{R}) \models \varphi[x, y]\}$. Then $B \in \Sigma_{(\gamma, 0)}(x) \subseteq \Sigma_{(\gamma, 2)}(x)$. By the Basis Property for $\Sigma_{(\gamma, 2)}$, $(\exists y \in B)$ with $y \in \Delta_{(\gamma, 2)}(x)$. Since $\{\gamma\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$ we have that $\{y\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. So $y \in A$. This shows that $(\forall x \in A) A$ is a basis for $\Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. So A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.

(3). Let α, β, x be as in (3). Notice that $\{\beta\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$.

(a) $\Rightarrow$ (b). Since $\{\beta\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$, we have that $\Sigma_\omega(J_\beta(\mathbb{R})) \subseteq \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$. So if A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct, then $(\forall w \in A) A$ is a basis for $\Sigma_1(J_\alpha(\mathbb{R}), \{x, w\})$, so $(\forall w \in A) A$ is a basis for $\Sigma_\omega(J_\beta(\mathbb{R}), \{w\})$.

(b) $\Rightarrow$ (c). This follows from the fact that

$$J_\beta(\mathbb{R}) \models (\exists z)(z \text{ is a good parameter for } J_\beta(\mathbb{R})).$$

(c) $\Rightarrow$ (a). Let z be as in (c). Let $w \in A$. Let B be a non-empty set of reals with $B \in \Sigma_1(J_\alpha(\mathbb{R}), \{w\})$. Then there is a $B' \subseteq B$, $B' \neq \emptyset$, $B' \in \Sigma_\omega(J_\beta(\mathbb{R}), \{w\})$. Let $n \in \omega$ with $B' \in \Sigma_n(J_\beta(\mathbb{R}), \{w\})$, and n odd. Then $B' \in \Sigma_{(\beta, n-1)}(\langle z, w \rangle)$. As $n - 1$ is even, by the Basis Property for $\Sigma_{(\beta, n-1)}$, $(\exists y \in B') y \in \Delta_{(\beta, n-1)}(\langle z, w \rangle)$. As $\{\beta\} \in \Sigma_1(J_\alpha(\mathbb{R}), \{x\})$, we have that $y \in \Sigma_1(J_\alpha(\mathbb{R}), \{x, z, w\})$. So $y \in A$. This shows that $(\forall w \in A) A$ is a basis for $\Sigma_1(J_\alpha(\mathbb{R}), \{w\})$. So A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct. $\square$

Remark 2.32. Notice that the one case not covered by Lemma 2.31 is the case in which $\alpha = \beta + 1$, and there is an $x \in A$ such that x is a 0-name for α , but β is not projective-like. For example, let $\beta = \kappa^{\mathbb{R}} =$ the least ordinal κ such that $J_\kappa(\mathbb{R})$ is admissible, and let $\alpha = \beta + 1$. Then 0 is a 0-name for α and β is not projective-like. In this case there is a set of reals A such that A is closed under join and closed downward under $\Delta_{(\alpha,0)}$, and A is not $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct. Since we are not interested in this case in this paper, we do not pursue this here.

Lemma 2.33. *Let $\alpha \geq 2$ be projective-like. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \geq 2$ be even. Let A be a non-empty set of reals, closed under join and closed downward under $\Delta_{(\alpha,n)}$. Then the following are equivalent.*

- (a) A is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -correct.
- (b) $(\exists x \in A) x$ is a good parameter for $J_\alpha(\mathbb{R})$.

Proof.

(a) $\Rightarrow$ (b). This follows from the fact that the set of good parameters for $J_\alpha(\mathbb{R})$ is $\Pi_2(J_\alpha(\mathbb{R}))$.

(b) $\Rightarrow$ (a). Let x be as in (b). Now let $z \in A$ and let B be a non-empty set of reals with $B \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{z\})$. Then $B \in \Sigma_{(\alpha,n)}(\langle x, z \rangle)$. By the Basis Property for $\Sigma_{(\alpha,n)}$, $(\exists y \in B) y \in \Delta_{(\alpha,n)}(\langle x, z \rangle)$. So $B \cap A \neq \emptyset$. $\square$

Suppose that $A \subset \mathbb{R}$ and A is $\Sigma_n(J_\alpha(\mathbb{R}))$ -correct. Then there is some ordinal $\bar{\alpha} \geq 1$ such that

- (1) $\mathbb{R} \cap J_{\bar{\alpha}}(A) = A$ and
- (2) $\exists j : (J_{\bar{\alpha}}(A); \in, A) \rightarrow (J_\alpha(\mathbb{R}); \in, \mathbb{R})$ such that j is Σ_n elementary.

But how can we find the ordinal $\bar{\alpha}$ given the set A ? It turns out that this question will be important to us when we apply our results about correctness in Section 4. In this paper we will only be dealing with ordinals $\alpha < \omega_1^{\omega_1}$. In this special case, the problem of finding $\bar{\alpha}$ is simpler. This is because every ordinal $\alpha < \omega_1^{\omega_1}$ is coded by a finite set of countable ordinals, and every countable ordinal is coded by a real. We turn to a discussion of this topic now.

A Coding Scheme

We will need a scheme for coding an ordinal $\alpha < \omega_1^{\omega_1}$ using countable ordinals. Such an α is easily coded by a finite function on the countable ordinals.

Definition 2.34. Let $\mathfrak{C}$ be the set of all finite partial functions from ω_1 to $\omega_1 - \{0\}$. $\mathfrak{C}$ will be our set of codes for ordinals $\alpha < \omega_1^{\omega_1}$. If $s \in \mathfrak{C}$ let $\bar{s}$ denote the total function $\bar{s} : \omega_1 \rightarrow \omega_1$ extending s such that $\bar{s}(\xi) = 0$ if $\xi \notin \text{dom}(s)$. For $s, t \in \mathfrak{C}$ say that $s < t$ if $\bar{s}(\xi) < \bar{t}(\xi)$, where ξ is greatest so that $\bar{s}(\xi) \neq \bar{t}(\xi)$. It is easy to see that $(\mathfrak{C}, <)$ is a wellorder of order type $\omega_1^{\omega_1}$. For $s \in \mathfrak{C}$ let $|s|$ = the rank of s in $(\mathfrak{C}, <)$. For $\alpha < \omega_1^{\omega_1}$ let $\text{code}(\alpha)$ = the unique $s \in \mathfrak{C}$ such that $|s| = \alpha$. One more piece of notation: For $s \in \mathfrak{C}$ let $\max(s)$ be the largest countable ordinal in $\text{dom}(s) \cup \text{ran}(s)$.

Below we record some trivial facts about our coding scheme which we will need to use later.

Lemma 2.35.

- (a) Let P be the relation on $\mathfrak{C}$ defined by $P(s, t) \Leftrightarrow |s| < |t|$. Then P is Σ_0 definable.
- (b) Let R be the relation on $\mathfrak{C}$ defined by $R(s) \Leftrightarrow \text{cof}(|s|) > \omega$. Then R is Σ_0 definable.
- (c) Let $\alpha < \omega_1^{\omega_1}$. For all $\xi < \max(\text{code}(\alpha))$ there is an $\alpha_0 < \alpha$ such that for all α' with $\alpha_0 \leq \alpha' < \alpha$, $\xi \leq \max(\text{code}(\alpha'))$.
- (d) $\max(\text{code}(\alpha + 1)) \leq \max(\text{code}(\alpha)) + 1$.
- (e) Let $\alpha < \omega_1^{\omega_1}$ be a limit ordinal of cofinality ω . Then there are cofinally many $\alpha' < \alpha$ such that $\max(\text{code}(\alpha')) \leq \max(\text{code}(\alpha))$.

Proof.

(a) $P(s, t) \Leftrightarrow |s| < |t| \Leftrightarrow s < t \Leftrightarrow \xi \notin \text{dom}(s) \vee s(\xi) < t(\xi)$, where ξ is the largest ordinal in $\text{dom}(s) \cup \text{dom}(t)$ such that either $\xi \notin \text{dom}(s) \cap \text{dom}(t)$ or $s(\xi) \neq t(\xi)$.

(b) $R(s) \Leftrightarrow \xi$ is a successor ordinal $\wedge s(\xi)$ is a successor ordinal, where ξ is the least ordinal in $\text{dom}(s)$.

(c) Let $s = \text{code}(\alpha)$. Fix $\xi < \max(s)$. We may assume that $\xi > 0$. Let η be least such that $\eta \in \text{dom}(s)$.

Case 1. $\xi < \eta$.

Let $s_0 = s - \{(\eta, s(\eta))\} \cup \{(\xi, 1)\}$. Let $\alpha_0 = |s_0|$. This works.

Case 2. $\eta \leq \xi < s(\eta)$.

Let $s_0 = s - \{(\eta, s(\eta))\} \cup \{(\eta, \xi)\}$. Let $\alpha_0 = |s_0|$. This works.

Case 3. $\eta \leq \xi, s(\eta) \leq \xi$.

Let $s_0 = s - \{(\eta, s(\eta))\}$. Let $\alpha_0 = |s_0|$. This works.

(d) This is a particular case of (c).

(e) Let $s \in \mathfrak{C}$ be a code. Then $\alpha = |s|$ is a limit ordinal of cofinality ω iff ξ is a limit ordinal or $s(\xi)$ is a limit ordinal, where ξ is the least ordinal in $\text{dom}(s)$. If $s(\xi)$ is a limit ordinal then $\{ |s - \{(\xi, s(\xi))\} \cup \{(\xi, \eta)\}| : \eta < s(\xi) \}$ is a cofinal set of ordinals $\alpha' < \alpha$ such that $\max(\text{code}(\alpha')) \leq \max(\text{code}(\alpha))$. If ξ is a limit ordinal and $s(\xi) = \eta + 1$ then $\{ |s - \{(\xi, s(\xi))\} \cup \{(\xi, \eta)\} \cup \{(\gamma, 1)\}| : \gamma < \xi \}$ works. $\square$

We conclude this section with a lemma about correctness, in the special case that $\alpha < \omega_1^{\omega_1}$. This is the only result about correctness which we will need in the sequel.

If M is a transitive model of ZF and $s \in \mathfrak{C}$ and $\max(s) < \omega_1^M$, then we may interpret the term “ $|s|$ ” in M . Clearly $|s|^M$ is just the rank of s in the wellorder $(\mathfrak{C}^M, <)$, where $\mathfrak{C}^M$ is $\{ s \in \mathfrak{C} \mid \max(s) < \omega_1^M \}$. We will use this notion in the next lemma.

Lemma 2.36. *Let $\alpha < \omega_1^{\omega_1}$, and let $n \geq 0$ be even. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $A \subseteq \mathbb{R}$ be such that A is closed downward under $\Delta_{(\alpha, n)}$, and $A = \mathbb{R} \cap L(A)$. Let $s = \text{code}(\alpha)$ and suppose that $\max(s) < \omega_1^{L(A)}$. Let $\bar{\alpha} = |s|^{L(A)}$. Then there exists a Σ_{n+1} elementary embedding*

$$j : J_{\bar{\alpha}}(A) \rightarrow J_{\alpha}(\mathbb{R})$$

Proof. We begin by showing that the conclusion of the lemma is true for *some* ordinal $\bar{\alpha}$. Then we will show that it is true for our particular $\bar{\alpha}$. The fact that the conclusion of the lemma is true for some $\bar{\alpha}$ is just the fact that A is $\Sigma_{n+1}(J_{\alpha}(\mathbb{R}))$ -correct. So we begin by showing this. Since we are assuming that n is even and A is closed under join and closed downward under $\Delta_{(\alpha, n)}$, Lemmas 2.31 and 2.33 above indicate that we must show that there is some good parameter in A . We will show that there is one—namely any real coding s .

Recall that WO is the set of reals coding countable ordinals. WO is a Π_1^1 set. If $w \in \text{WO}$ then $|w|$ is the countable ordinal coded by w . (Do not be confused by the fact that if $s \in \mathfrak{C}$ then we are also using the notation $|s|$ to refer to the ordinal $\alpha < \omega_1^{\omega_1}$ such that $\text{code}(\alpha) = s$.) Let $\mathfrak{C}^*$ be the set of finite partial functions h from WO to WO so that for no w in the range of h is it true that $|w| = 0$. So $\mathfrak{C}^*$ is essentially a set of reals, and $\mathfrak{C}^*$ is a collection of codes for the elements of $\mathfrak{C}$. Let $\pi : \mathfrak{C}^* \rightarrow \mathfrak{C}$ be the obvious decoding function. For $x, y \in \mathfrak{C}^*$, say that $x <^* y \Leftrightarrow \pi(x) < \pi(y)$. Thus $(\mathfrak{C}^*, <^*)$ is a pre-wellorder of ordertype $\omega_1^{\omega_1}$. Notice that $(\mathfrak{C}^*, <^*)$ is lightface definable over $J_1(\mathbb{R})$. Let $\text{Rnk} : \mathfrak{C}^* \rightarrow \omega_1^{\omega_1}$ be the rank function. So $\text{Rnk}(x) = |\pi(x)|$. For $\xi < \omega_1^{\omega_1}$ let us use the abbreviation $\text{Rnk} \upharpoonright \xi$ to mean $\text{Rnk} \upharpoonright \{x \mid \text{Rnk}(x) < \xi\}$.

Claim 1. *Let $\beta \geq 2$. Then*

- (a) $\text{Rnk} \upharpoonright \text{OR}^{J_\beta(\mathbb{R})}$ is $\Sigma_1(J_\beta(\mathbb{R}))$.
- (b) For all $\xi \in J_\beta(\mathbb{R})$, $\text{Rnk} \upharpoonright \xi \in J_\beta(\mathbb{R})$.

Proof of Claim 1. This is a standard induction on β . ◇

Since $\max(s) < \omega_1^{L(A)}$, there is an $x_0 \in L(A)$ such that $x_0 \in \mathfrak{C}^*$ and $\pi(x_0) = s$. Thus $\text{Rnk}(x_0) = \alpha$. By a simple recursive coding, we may think of x_0 as a real. Thus $x_0 \in A$. Fix such a real x_0 .

Claim 2. x_0 is a good parameter for $J_\alpha(\mathbb{R})$.

Proof of Claim 2. α is least such that $J_\alpha(\mathbb{R})$ satisfies:

$$(\forall y \in \mathfrak{C}^*) [y <^* x_0 \Rightarrow (\exists \beta, \exists X)(\beta = \text{Rnk}(y) \wedge X = J_\beta(\mathbb{R}))].$$

Thus x_0 is a 1-name for α and so x_0 is a good parameter for $J_\alpha(\mathbb{R})$. ◇

We want to show that A is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -correct. First suppose that $n \geq 2$. Then we have that A is closed downward under $\Delta_{(\alpha, n)}$, and A is closed under join, and there is a good parameter for $J_\alpha(\mathbb{R})$ in A . It follows from lemma 2.33 that A is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -correct. Next suppose that $n = 0$. We have that A is closed downward under $\Delta_{(\alpha, 0)}$ and A is closed under join. If α is a limit ordinal, then it follows from part (a) of Lemma 2.31 that A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct. Suppose then that α is a successor ordinal. Since $\max(\text{code}(\alpha)) < \omega_1^{L(A)}$, it is also true that $\max(\text{code}(\alpha - 1)) < \omega_1^{L(A)}$. Thus there is

an $x \in L(A)$ such that $x \in \mathfrak{C}^*$ and $\text{Rnk}(x) = \alpha - 1$. As in the claim above, x is a good parameter for $J_{\alpha-1}(\mathbb{R})$. It follows from parts (b) and (c) of Lemma 2.31 that A is $\Sigma_1(J_\alpha(\mathbb{R}))$ -correct.

So we now have that in all cases A is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -correct. Thus there is a Σ_{n+1} embedding:

$$j : J_\gamma(A) \rightarrow J_\alpha(\mathbb{R}).$$

We only need to see that $\gamma = \bar{\alpha}$. Recall that $s = \text{code}(\alpha)$, and $\bar{\alpha} = |s|^{L(A)}$, and $x_0 \in A$, and $\pi(x_0) = s$.

We split into three cases.

Case 1. $\alpha < \omega_1$

Then $s = \{(0, \alpha)\}$. Thus $\bar{\alpha} = \alpha$. Also, x_0 is essential an element of WO , with $|x_0| = \alpha$. Notice that if $w \in \text{WO} \cap A$, and $|w| = \xi \in J_\gamma(A)$, then $|w|^{J_\gamma(A)} = \xi$. Furthermore, the relation $\xi = |w|$ is $\Sigma_1(J_\gamma(A))$. Since $x_0 \in A$, and j is at least Σ_1 elementary, it is easy to see that $\gamma = \alpha$.

Case 2. $\alpha = \omega_1$

Then $s = \{(1, 1)\}$ and so $\bar{\alpha} = \omega_1^{L(A)}$. Since j is at least Σ_1 elementary, it is easy to see that $\gamma = (\omega_1)^{J_\gamma(A)} = \omega_1^{L(A)}$.

Case 3. $\alpha > \omega_1$

Then $\gamma > \omega_1^{J_\gamma(A)} = \omega_1^{L(A)}$. As $\max(s) < \omega_1^{L(A)}$, $s \in J_\gamma(A)$. Notice that $j(s) = s$. First suppose towards a contradiction that $\gamma > \bar{\alpha}$. In particular $\bar{\alpha} \in J_\gamma(A)$. Then $\bar{\alpha} = |s|^{J_\gamma(A)}$, and so $j(\bar{\alpha}) = \alpha$. But this is a contradiction as $J_\gamma(A)$ satisfies that $J_{\bar{\alpha}}(\mathbb{R})$ exists. So $\gamma \leq \bar{\alpha}$. Now suppose towards a contradiction that $\gamma < \bar{\alpha}$. Then there is a $t \in \mathfrak{C}$ with $\max(t) < \omega_1^{J_\gamma(A)}$ such that $|t|^{J_\gamma(A)} = \gamma$. Notice that $j(t) = t$. Let $\delta = |t|$ in V . Thus $j(\gamma) = \delta$. But this contradicts the fact that $J_\alpha(\mathbb{R})$ satisfies that $J_\delta(\mathbb{R})$ exists. $\square$

3. BIG PREMICE

In this section we finally get to inner model theory. This section marks the beginning of the main portion of this paper.

In the previous section we defined the sets $A_{(\alpha,n)}$ and we stated the conjecture that $A_{(\alpha,n)}$ is a mouse set. We now begin the process of proving this conjecture. In this section we will define the notion of a premouse $\mathcal{M}$ being (α, n) -big. In the next section we will show that if $\mathcal{M}$ is a countable, iterable premouse, and $\mathcal{M}$ is (α, n) -big, then $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$. This will give us one half of the proof that $A_{(\alpha,n)}$ is a mouse set.

Our approach here is to generalize some work of Martin and Steel. We begin by describing this work. The following is (equivalent to) a definition from [St5]. We repeat it here for the readers convenience.

Definition 3.1. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $n \geq 0$. Then $\mathcal{M}$ is *n-big above* β iff there is an *active* initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that, letting κ be the critical point of the last extender from the $\mathcal{N}$ -sequence, we have that $\beta < \kappa$ and there are n ordinals $\delta_1, \dots, \delta_n$, with $\beta < \delta_1 < \dots < \delta_n < \kappa$, such that each δ_i is a Woodin cardinal of $\mathcal{N}$.

$\mathcal{M}$ is *n-small* above β iff $\mathcal{M}$ is not *n-big* above β . We will say that $\mathcal{M}$ is *n-big* iff $\mathcal{M}$ is *n-big* above 0, equivalently iff $\mathcal{M}$ is *n-big* above ω . Similarly with *n-small*.

In order to parallel our definition of $A_{(\alpha,n)}$ for $\alpha \geq 2$, we make the following definition with $\alpha = 1$. For $n \geq 0$ we set

$$A_{(1,n)} = \{ x \in \mathbb{R} \mid (\exists \xi < \omega_1) x \in \Delta_{n+2}^1(\xi) \}.$$

So $A_{(1,n)}$ is the same set that we were calling A_{n+2} in the introduction. We have changed the indexing here in order to better express the analogy between $\alpha = 1$ and $\alpha \geq 2$. The following theorem and its proof are implicit in [St5].

Theorem 3.2 (Martin, Steel, Woodin). *Let $\mathcal{M}$ be a countable, iterable premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $n \geq 0$, and suppose that $\mathcal{M}$ is *n-big* above β . Let $\mathbb{P} \in \mathcal{M}$ be a partial order, with $(\text{card}(\mathbb{P})^+)^{\mathcal{M}} \leq \beta$. Suppose that G is $\mathcal{M}$ -generic over $\mathbb{P}$. Then for all $x \in \mathbb{R} \cap \mathcal{M}[G]$, $A_{(1,n)}(x) \subseteq \mathbb{R} \cap \mathcal{M}[G]$. In particular, if $\mathcal{M}$ is *n-big* then $A_{(1,n)} \subseteq \mathbb{R} \cap \mathcal{M}$.*

Our goal in this section is to give a definition which extends Definition 3.1 above. In order to emphasize this, let us give a synonym for the term *n-big*. For $n \geq 0$ we will say that a premouse is $(1, n)$ -big above β iff it is *n-big* above β . Now let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $\alpha \geq 2$, and $n \geq 0$. We wish to define the notion: $\mathcal{M}$ is (α, n) -big above

β . To motivate the definition, we list five criteria that we would like for the definition to satisfy. The definition of n -big (Definition 3.1 above) satisfies the analogous criteria.

- (1) If a countable, iterable premouse $\mathcal{M}$ is (α, n) -big above β , and if $\mathbb{P} \in \mathcal{M}$ is a partial order with $(\text{card}(\mathbb{P})^+)^{\mathcal{M}} \leq \beta$, and if G is $\mathcal{M}$ -generic over $\mathbb{P}$, then for all $x \in \mathbb{R} \cap \mathcal{M}[G]$, $A_{(\alpha, n)}(x)$ should be contained in $\mathcal{M}[G]$.
- (2) If $\mathcal{N}$ is (α, n) -big above β and $\mathcal{N} \leq \mathcal{M}$ then $\mathcal{M}$ should be (α, n) -big above β .
- (3) If $\mathcal{M}$ is (α, n) -big above β and $\beta' < \beta$ then $\mathcal{M}$ should be (α, n) -big above β' .
- (4) If $\mathcal{M}$ is (α, n) -big above β and $(\alpha', n') <_{\text{lex}} (\alpha, n)$ then $\mathcal{M}$ should be (α', n') -big above β .
- (5) The property of being (α, n) -big should be preserved by non-dropping iterations. That is, if $\mathcal{M}$ is (α, n) -big above β , and if we iterate $\mathcal{M}$ without dropping to yield $i : \mathcal{M} \rightarrow \mathcal{M}'$, then $\mathcal{M}'$ should be (α, n) -big above $i(\beta)$.

The fifth criterion is motivated by the proof of Theorem 3.2 above, which we hope to extend. This criterion turns out to be the most problematic for us. In order to achieve criterion (5), we will aim for the following stronger criterion:

- (5') If $\mathcal{M}$ is (α, n) -big above β then $\mathcal{M}$ should have a code for α , and the statement “I am (α, n) -big above β ” should be definable in $\mathcal{M}$.

Right away we see that there is a conflict between criterion (5') and criterion (4). For example, suppose $\mathcal{M}$ is a countable premouse and $\mathcal{M}$ is $(\omega_1, 0)$ -big above β . By (4), $\mathcal{M}$ should be $(\alpha, 0)$ -big above β for every countable ordinal α . Then by (5'), $\mathcal{M}$ should have a code for every countable ordinal α . But this is impossible as $\mathcal{M}$ is countable.

Our solution to this conflict is the following. We will define an auxiliary notion: $\mathcal{M}$ is *explicitly* (α, n) -big above β . For this auxiliary notion we will drop criterion (4). So a premouse can be explicitly $(\omega_1, 0)$ -big without being explicitly $(\alpha, 0)$ -big for every countable α . This auxiliary notion will satisfy criteria (1), (2), (3), and (5'). In place of criterion (4) we will have the following:

- (4') If $\mathcal{M}$ is explicitly (α, n) -big above β and $(\alpha', n') <_{\text{lex}} (\alpha, n)$ then there is *an iterate* of $\mathcal{M}$ which is explicitly (α', n') -big above β .

Then, to recapture criterion (4), we will define as our primary notion: $\mathcal{M}$ is (α, n) -big above β iff for some (α', n') with $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$, $\mathcal{M}$ is explicitly (α', n') -big above β . Then this primary notion will satisfy criteria (1) through (5). (but not (5'))

There is one major drawback with the programme we have outlined above: we are only able to see how to carry it out for small ordinals α . In this paper, for the sake of concreteness, we will carry out the programme for $\alpha < \omega_1^{\omega_1}$. The ordinal $\omega_1^{\omega_1}$ has no special significance here. It is easy to see how the programme can be extended somewhat beyond $\omega_1^{\omega_1}$. We are not able to go very far beyond though. The difficulty seems to be with criterion (5') above. We are unable to see how to extend the definition of “explicitly (α, n) -big” to arbitrarily large ordinals α , so as to satisfy criterion (5'). We do not know if this is a limitation of the approach, or merely of the author. We suspect that it is a little of both.

In order to satisfy criterion (5') (described above), we will need a scheme for coding ordinals which yields a code for the uncountable ordinal α that may be found inside of a countable premouse. Recall that in the previous section we defined the coding scheme $(\mathfrak{C}, <)$ for ordinals $\alpha < \omega_1^{\omega_1}$.

We are now ready to begin our definition. For $\mathcal{M}$ a countable premouse and $\beta \in \text{OR}^{\mathcal{M}}$, and $2 \leq \alpha < \omega_1^{\omega_1}$, and $n \geq 0$, we wish to describe the notion: $\mathcal{M}$ is *explicitly* (α, n) -big above β . Our definition is by induction on (α, n) . As the basis of the induction, for $\alpha = 1$ and $n \geq 0$, we will say that $\mathcal{M}$ is *explicitly* $(1, n)$ -big above β iff $\mathcal{M}$ is n -big above β . (See Definition 3.1 above.) Now we turn to the inductive step. There are four different cases of the induction, and for ease of readability we split these up into four definitions below.

We begin with the case $n = 0$. In this context, recall that we have defined

$$A_{(\alpha, 0)} = \left\{ x \in \mathbb{R} \mid (\exists \alpha' < \alpha) x \in \text{OD}^{J_{\alpha'}(\mathbb{R})} \right\}.$$

Naively, in order to inductively arrange criterion (1) above, we would like to say that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β iff for all $\alpha' < \alpha$ and all n , there is some initial segment $\mathcal{N}_{(\alpha', n)} \trianglelefteq \mathcal{M}$ such that $\mathcal{N}_{(\alpha', n)}$ is explicitly (α', n) -big above β . (In fact this is what we will say for countable ordinals α . For example in the case $\alpha = 2$ we will have that $\mathcal{M}$ is explicitly $(2, 0)$ -big iff for all n there is an initial segment of $\mathcal{M}$ which is n -big.) But,

as was discussed above, we can not do this for uncountable ordinals α . So our actual definition will be an approximation of this. There are two cases, depending on whether or not $\text{cof}(\alpha) > \omega$. The case $\text{cof}(\alpha) \leq \omega$ is easy.

Definition 3.3. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$, and suppose that α is a successor ordinal, or a limit ordinal of cofinality ω . Then $\mathcal{M}$ is *explicitly* $(\alpha, 0)$ -big above β iff for cofinally many $\alpha' < \alpha$, and all n , there is a proper initial segment $\mathcal{N}_{(\alpha', n)} \triangleleft \mathcal{M}$ such that $\mathcal{N}_{(\alpha', n)}$ is explicitly (α', n) -big above β .

We next turn to the case $n = 0$ and $\text{cof}(\alpha) > \omega$. The following definition is motivated by criteria (4') and (5') above.

Definition 3.4. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Suppose that $\alpha < \omega_1^{\omega_1}$ is a limit ordinal of cofinality ω_1 . Then $\mathcal{M}$ is *explicitly* $(\alpha, 0)$ -big above β iff there is an *active* initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that, letting κ be the critical point of the last extender from the $\mathcal{N}$ -sequence, we have that $\beta < \kappa$ and $\text{code}(\alpha) \in J_{\kappa}^{\mathcal{N}}$, and for all $\alpha' < \alpha$, if $\text{code}(\alpha') \in J_{\kappa}^{\mathcal{N}}$, then for all n , there is a proper initial segment $\mathcal{N}_{(\alpha', n)} \triangleleft J_{\kappa}^{\mathcal{N}}$ such that $\mathcal{N}_{(\alpha', n)}$ is explicitly (α', n) -big above β .

Note. Let $\mathcal{N}$ be as in the above definition. Then we will say that $\mathcal{N}$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β .

We next turn our attention to the case $n > 0$. This is the successor case of our inductive definition. The procedure for passing from explicitly $(\alpha, n-1)$ -big to explicitly (α, n) -big is the same in our context as it is in the case $\alpha = 1$. That is, we add a Woodin cardinal.

Note. The case $\text{cof}(\alpha) \leq \omega$, and $n = 1$ is more complicated and is handled separately. We start with the easy case.

Definition 3.5. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$. If $\text{cof}(\alpha) > \omega$ then suppose that $n \geq 1$. If $\text{cof}(\alpha) \leq \omega$ then suppose that $n \geq 2$. Then $\mathcal{M}$ is *explicitly* (α, n) -big above β iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$, and an ordinal $\delta \in \mathcal{N}$ with $\beta < \delta$, such that δ is a Woodin cardinal of $\mathcal{N}$, and $\mathcal{N}$ is explicitly $(\alpha, n-1)$ -big above δ .

To emphasize the similarity with Definition 3.1 we state the following easy lemma.

Lemma 3.6. *Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$.*

First suppose that $\text{cof}(\alpha) > \omega$ and $n \geq 1$. Then $\mathcal{M}$ is explicitly (α, n) -big above β iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$, and n ordinals $\delta_1 < \dots < \delta_n \in \mathcal{N}$, with $\beta < \delta_1$, such that each δ_i is a Woodin cardinal of $\mathcal{N}$, and $\mathcal{N}$ is explicitly $(\alpha, 0)$ -big above δ_n .

Next suppose that $\text{cof}(\alpha) \leq \omega$ and $n \geq 2$. Then $\mathcal{M}$ is explicitly (α, n) -big above β iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$, and $n - 1$ ordinals $\delta_1 < \dots < \delta_{n-1} \in \mathcal{N}$, with $\beta < \delta_1$, such that each δ_i is a Woodin cardinal of $\mathcal{N}$, and $\mathcal{N}$ is explicitly $(\alpha, 1)$ -big above δ_{n-1} .

To complete our definition then, we must define the notion of explicitly $(\alpha, 1)$ -big in the case that α is a successor ordinal, or a limit ordinal of cofinality ω . It is instructive to consider the definition that we would have made if we had not excluded this case from Definition 3.5 above. Considering Definition 3.3 above, we see that we would have defined $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$, and an ordinal $\delta \in \mathcal{N}$ with $\beta < \delta$, such that δ is a Woodin cardinal of $\mathcal{N}$, and for cofinally many $\alpha' < \alpha$, and all n , there is a proper initial segment $\mathcal{N}_{(\alpha', n)} \triangleleft \mathcal{N}$ such that $\mathcal{N}_{(\alpha', n)}$ is explicitly (α', n) -big above δ . This is, at first sight, a reasonable definition. But it turns out that a premouse with this property is “bigger” than necessary. (We are trying to give the weakest possible definition which still meets our criteria.) So our actual definition will be a weakening of this property. The idea is that we do not need a single fixed δ for all of the pairs (α', n) . We can have a different $\delta_{(\alpha', n)}$ for each pair (α', n) , and these $\delta_{(\alpha', n)}$ do not have to be Woodin in all of $\mathcal{N}$, but only in the initial segment $\mathcal{N}_{(\alpha', n)} \triangleleft \mathcal{N}$. In exchange for this weakening we must demand that each $\delta_{(\alpha', n)}$ is a cardinal of $\mathcal{N}$, and that $\mathcal{N}$ is a model of KP. This brings us to the next two definitions.

Warning. Readers of [Ru1] should note that the indexing here is different than in that paper. For α a countable ordinal, what we are here calling “explicitly $(\alpha, 1)$ -big” was called “ $(\alpha, 0)$ -big” in [Ru1]. What we are here calling “explicitly $(\alpha, 0)$ -big” did not have a name in [Ru1]. In [Ru1] we did not deal with uncountable ordinals α , and when one does the indexing used here is better.

Definition 3.7. Let $\mathcal{M}$ be a premouse, $\alpha < \omega_1^{\omega_1}$, and $n \in \omega$. Let $\delta \in \text{OR}^{\mathcal{M}}$. Then we will say that δ is *locally* (α, n) -Woodin in $\mathcal{M}$ iff there is a proper initial segment $\mathcal{J}_\gamma^{\mathcal{M}} \triangleleft \mathcal{M}$ such that $\delta \in \mathcal{J}_\gamma^{\mathcal{M}}$ and

- (a) δ is a Woodin cardinal of $\mathcal{J}_\gamma^{\mathcal{M}}$, and
- (b) $\mathcal{J}_\gamma^{\mathcal{M}}$ is explicitly (α, n) -big above δ .

Note. In the above situation we will say that $\mathcal{J}_\gamma^{\mathcal{M}}$ witnesses that δ is locally (α, n) -Woodin in $\mathcal{M}$.

Definition 3.8. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Suppose that $2 \leq \alpha < \omega_1^{\omega_1}$ and that α is a successor ordinal, or a limit ordinal of cofinality ω . Then $\mathcal{M}$ is *explicitly* $(\alpha, 1)$ -big above β iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ with $\beta \in \mathcal{N}$, and an ordinal $\rho \in \mathcal{N}$ with $\beta < \rho$, such that ρ is a limit cardinal of $\mathcal{N}$ and:

- (i) $\mathcal{N}$ is an admissible set, and
- (ii) $\text{code}(\alpha) \in J_\rho^{\mathcal{N}}$, and
- (iii) for cofinally many $\alpha' < \alpha$, and all $n \in \omega$, and cofinally many $\delta < \rho$, we have that:
 - (a) δ is a cardinal of $\mathcal{N}$, and
 - (b) δ is locally (α', n) Woodin in $\mathcal{J}_\rho^{\mathcal{N}}$.

Note. Let $\mathcal{N}$ and ρ be as in the above definition. Then we will say that $(\mathcal{N}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β .

To understand the above definition, we invite the reader to contemplate what an explicitly $(2, 1)$ -big premouse looks like. In Remark 3.9 and Figure 1 below we explore the definition of explicitly $(\alpha, 1)$ -big in the case that α is a successor ordinal.

Remark 3.9. Suppose $\alpha = \alpha' + 1$ with $\alpha' \geq 1$. Suppose that $\mathcal{M}$ is a countable, iterable premouse and $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big (above 0) but no proper initial segment of $\mathcal{M}$ is. We will examine what $\mathcal{M}$ looks like. (See Figure 1 above.) For a particularly concrete example, consider the case $\alpha = 2$. Because no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big, there is an ordinal $\rho \in \mathcal{M}$ such that $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big. Again, because no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big, we have




FIGURE 1. $\alpha \geq 2$ is a successor and $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big

the following: ρ is the largest cardinal of $\mathcal{M}$, $\mathcal{M}$ is a passive premouse, $\mathcal{M} = J_{\gamma}^{\vec{E}}$, with $\vec{E} \subset J_{\rho}^{\mathcal{M}}$, and γ is the first admissible ordinal over $\mathcal{J}_{\rho}^{\mathcal{M}}$. There is a sequence of ordinals $\delta_0 < \gamma_0 < \delta_1 < \gamma_1 < \delta_2 < \dots$ such that $\max(\text{code}(\alpha)) < \delta_0$, and each δ_n is a cardinal of $\mathcal{M}$ and a Woodin cardinal of $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$, and $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$ is explicitly (α', n) -big above δ_n . We have that $\sup \{\delta_n\} = \sup \{\gamma_n\} = \rho$. (For otherwise let $\rho' = \sup \{\delta_n\}$. Then, letting $\mathcal{N}$ be the first admissible structure over $\mathcal{J}_{\rho'}^{\mathcal{M}}$, we have that $\mathcal{N} \triangleleft \mathcal{J}_{\rho'}^{\mathcal{M}}$ and $(\mathcal{N}, \rho')$ witnesses that $\mathcal{N}$ is explicitly $(\alpha, 0)$ -big above β .)

Later we will show that the sequence $\langle \delta_n, \gamma_n \mid n \in \omega \rangle$ is in $\mathcal{M}$. This turns out to be very important. We call this sequence an α -ladder. See Definition 3.28.

Because δ_n is a (strong limit) cardinal of $\mathcal{M}$ and $\mathcal{M}$ is strongly acceptable, $V_{\delta_n}^{\mathcal{M}} \subset \mathcal{J}_{\gamma_n}^{\mathcal{M}}$, and so δ_n is actually Woodin in $\mathcal{M}$ with respect to all functions $f : \delta_n \rightarrow \delta_n$ which are in $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$. But which functions are in $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$? The fact that $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$ is (α', n) -big above δ_n (and that $\mathcal{M}$ is iterable) implies that $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$ contains all functions $f : \delta_n \rightarrow \delta_n$ which are (in some appropriate sense) $\Delta_{(\alpha', n)}$ definable with $\mathcal{J}_{\delta_n}^{\mathcal{M}}$ as a parameter. This follows from the results in the next section.

This completes the inductive definition of the notion: $\mathcal{M}$ is explicitly (α, n) -big above β , for $\alpha < \omega_1^{\omega_1}$. As we remarked earlier, there is no special significance to the ordinal

$\omega_1^{\omega_1}$, and it is easy to see how to extend our definition somewhat beyond this ordinal. In fact it will actually be useful for us to extend the definition by one more step. We begin by extending our coding scheme by one more step. For uniformity, let us give a code to the ordinal $\omega_1^{\omega_1}$. We will say $\text{code}(\omega_1^{\omega_1}) = \emptyset$. Compare the following with Definition 3.4 above.

Definition 3.10. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Then $\mathcal{M}$ is *explicitly* $(\omega_1^{\omega_1}, 0)$ -big above β iff there is an *active* initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that, letting κ be the critical point of the last extender from the $\mathcal{N}$ -sequence, we have that $\beta < \kappa$, and for all $\alpha < \omega_1^{\omega_1}$, if $\text{code}(\alpha) \in J_{\kappa}^{\mathcal{N}}$, then for all n , there is a proper initial segment $\mathcal{N}_{(\alpha, n)} \triangleleft \mathcal{J}_{\kappa}^{\mathcal{N}}$ such that $\mathcal{N}_{(\alpha, n)}$ is explicitly (α, n) -big above β .

Definition 3.11. Suppose $(1, 0) \leq_{\text{lex}} (\alpha, n) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$. We will say that $\mathcal{M}$ is (α, n) -big above β iff $\mathcal{M}$ is explicitly (α', n') -big above β , for some (α', n') with $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$. We will say that $\mathcal{M}$ is (α, n) -small above β iff $\mathcal{M}$ is not (α, n) -big above β . We will say that $\mathcal{M}$ is (explicitly) (α, n) -big iff $\mathcal{M}$ is (explicitly) (α, n) -big above 0, equivalently above ω . Similarly with (α, n) -small.

In the rest of this section we will study the notions which we have just defined. In particular we will show that the notion of (α, n) -big satisfies criteria (2) through (5) mentioned above, and that the notion of explicitly (α, n) -big satisfies criteria (2), (3), (4'), and (5'). The proof that criteria (1) is satisfied will be given in the next section. We begin with the following lemma which follows immediately from the definitions.

Lemma 3.12. CRITERION 2. *If $\mathcal{N}$ is (α, n) -big above β and $\mathcal{N} \trianglelefteq \mathcal{M}$ then $\mathcal{M}$ is (α, n) -big above β . (Similarly with explicitly (α, n) -big.)*

CRITERION 3. *If $\mathcal{M}$ is (α, n) -big above β and $\beta' < \beta$ then $\mathcal{M}$ is (α, n) -big above β' . (Similarly with explicitly (α, n) -big.)*

CRITERION 4. *If $\mathcal{M}$ is (α, n) -big above β and $(\alpha', n') <_{\text{lex}} (\alpha, n)$ then $\mathcal{M}$ is (α', n') -big above β . (Not true with explicitly (α, n) -big.)*

Although the notion of explicitly (α, n) -big does not satisfy criterion (4), the following two lemmas show that it does “locally.”

Lemma 3.13. *If $\mathcal{M}$ is explicitly $(\alpha, n+1)$ -big above β then $\mathcal{M}$ is explicitly (α, n) -big above β .*

Proof. If $\text{cof}(\alpha) > \omega$ or $n \geq 1$ this follows immediately from Definition 3.5. If $\text{cof}(\alpha) \leq \omega$ and $n = 0$, then it follows immediately from Definitions 3.3 and 3.8. $\square$

Lemma 3.14. *Suppose $2 \leq \alpha < \omega_1^{\omega_1}$. Let ξ be a countable ordinal greater than 0. If $\mathcal{M}$ is explicitly $(\alpha + \xi, 0)$ -big above β then $\mathcal{M}$ is explicitly (α, n) -big above β for all n .*

Proof. By induction on ξ . First suppose $\xi = 1$, and $\mathcal{M}$ is explicitly $(\alpha + 1, 0)$ -big above β . By Definition 3.3 $\mathcal{M}$ is explicitly (α, n) -big above β for all n . The inductive step is similar. $\square$

In addition to the above two lemmas, it is also true that if $\mathcal{M}$ is explicitly (α, n) -big above β and $(\alpha', n') <_{\text{lex}} (\alpha, n)$ and $\max(\text{code}(\alpha'))$ is sufficiently small, then $\mathcal{M}$ is explicitly (α', n') -big above β . This idea is expressed in Lemmas 3.15, and 3.16 below.

Lemma 3.15. *Let $2 \leq \alpha \leq \omega_1^{\omega_1}$, and suppose that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . Let $\alpha' < \alpha$ and suppose that $\max(\text{code}(\alpha')) < \beta$. Then for all n , $\mathcal{M}$ is explicitly (α', n) -big above β .*

Proof. First suppose that $\text{cof}(\alpha) > \omega$. Then there is an active initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that, letting κ be the critical point of the last extender from the $\mathcal{N}$ -sequence, we have that $\beta < \kappa$, and for all $\hat{\alpha} < \alpha$, if $\text{code}(\hat{\alpha}) \in J_\kappa^{\mathcal{N}}$ then for all n , $\mathcal{M}$ is explicitly $(\hat{\alpha}, n)$ -big above β . Since $\max(\text{code}(\alpha')) < \beta < \kappa$ we have that $\text{code}(\alpha') \in J_\kappa^{\mathcal{N}}$, and so for all n , $\mathcal{M}$ is explicitly (α', n) -big above β .

Next suppose that $\text{cof}(\alpha) \leq \omega$. We argue by induction on α . By Lemma 3.14 we may assume that $\alpha' + 1 < \alpha$. Then, by definition, there is an $\hat{\alpha}$ with $\alpha' < \hat{\alpha} < \alpha$ such that $\mathcal{M}$ is explicitly $(\hat{\alpha}, 0)$ -big above β . So, by the lemma we are proving, for $\hat{\alpha}$, $\mathcal{M}$ is (α', n) -big above β for all n . $\square$

Lemma 3.16. *Let $2 \leq \alpha < \omega_1^{\omega_1}$, and suppose that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . Let $\alpha' < \alpha$ and suppose that $\max(\text{code}(\alpha')) < \max(\text{code}(\alpha))$. Then for all n , $\mathcal{M}$ is explicitly (α', n) -big above β .*

Proof. Suppose first that $\text{cof}(\alpha) = \omega_1$. Then there is an active initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ such that, letting κ be the critical point of the last extender from the $\mathcal{N}$ -sequence, we have that $\beta < \kappa$, and $\text{code}(\alpha) \in J_\kappa^\mathcal{N}$, and for all $\hat{\alpha} < \alpha$, if $\text{code}(\hat{\alpha}) \in J_\kappa^\mathcal{N}$ then for all n , $\mathcal{M}$ is explicitly $(\hat{\alpha}, n)$ -big above β . Since $\max(\text{code}(\alpha')) < \max(\text{code}(\alpha))$ we have that $\text{code}(\alpha') \in J_\kappa^\mathcal{N}$ and so for all n , $\mathcal{M}$ is explicitly (α', n) -big above β .

Next suppose that $\alpha = \alpha_0 + 1$ is a successor ordinal, where $\text{cof}(\alpha_0) = \omega_1$. By Lemma 3.14, $\mathcal{M}$ is explicitly $(\alpha_0, 0)$ -big above β . The argument is similar to the previous paragraph, using α_0 in place of α , and using the fact that by part (d) of Lemma 2.35, $\max(\text{code}(\alpha)) \leq \max(\text{code}(\alpha_0)) + 1$.

Now suppose that $\alpha = \alpha_0 + 1$ is a successor ordinal, where $\text{cof}(\alpha_0) \leq \omega$. By Lemma 3.14, $\mathcal{M}$ is explicitly $(\alpha_0, 1)$ -big above β , and we may assume that $\alpha' + 1 < \alpha_0$. Let $(\mathcal{N}, \rho)$ witness that $\mathcal{M}$ is explicitly $(\alpha_0, 1)$ -big above β . By definition, $\max(\text{code}(\alpha_0)) < \rho$. As ρ is a limit cardinal of $\mathcal{N}$, $\max(\text{code}(\alpha_0)) + 1 < \rho$. By part (d) of Lemma 2.35, $\max(\text{code}(\alpha)) \leq \max(\text{code}(\alpha_0)) + 1$. So $\max(\text{code}(\alpha')) < \max(\text{code}(\alpha)) < \rho$. Examining Definition 3.8 we see that there is an ordinal $\delta < \rho$ and an ordinal $\hat{\alpha} < \alpha_0$ such that $\max(\text{code}(\alpha')) < \delta$ and $\alpha' < \hat{\alpha}$, and there is an ordinal γ with $\delta < \gamma < \rho$ such that $\mathcal{J}_\gamma^\mathcal{N}$ is explicitly $(\hat{\alpha}, 0)$ -big above δ . By Lemma 3.15, for $\hat{\alpha}$, $J_\gamma^\mathcal{N}$ is explicitly (α', n) -big above δ for all n .

Finally suppose that α is a limit ordinal of cofinality ω . Let $\xi = \max(\text{code}(\alpha'))$. So $\xi < \max(\text{code}(\alpha))$. By part (c) of Lemma 2.35, there is an $\alpha_0 < \alpha$ such that for all $\hat{\alpha}$ with $\alpha_0 < \hat{\alpha} < \alpha$ we have that $\xi < \max(\text{code}(\hat{\alpha}))$. Let $\hat{\alpha}$ be least such that $\alpha' < \hat{\alpha}$ and $\alpha_0 < \hat{\alpha}$ and $\mathcal{M}$ is explicitly $(\hat{\alpha}, 0)$ -big above β . Then $\hat{\alpha}$ is not a limit ordinal of cofinality ω . So by the lemma we are proving, for $\hat{\alpha}$, $\mathcal{M}$ is explicitly (α', n) -big above β for all n . $\square$

Suppose that $\mathcal{M}$ is explicitly (α, n) -big above β . For which $\beta' > \beta$ is it also true that $\mathcal{M}$ is explicitly (α, n) -big above β' ? For example, if $n = k + 1$ and there is an ordinal $\delta > \beta$ such that δ is Woodin in $\mathcal{M}$ and $\mathcal{M}$ is explicitly (α, k) -big above δ , then $\mathcal{M}$ is explicitly (α, n) -big above β' for all $\beta' < \delta$. Also, if $\text{cof}(\alpha) \leq \omega$ and $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β , then in fact $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β' , for all $\beta' < \rho$. These considerations will crop up in the next section.

There is one case which is just convoluted enough to require a short proof. We describe this in the following proposition.

Proposition 3.17. *Let $\alpha \leq \omega_1^{\omega_1}$ and suppose $\text{cof}(\alpha) = \omega_1$. Suppose that $\mathcal{M}$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above some ordinal β . By definition $\mathcal{M}$ is active. Let κ be the critical point of the last extender from the $\mathcal{M}$ sequence. Then $\mathcal{M}$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β' , for all $\beta' < \kappa$.*

Proof. By definition, $\beta < \kappa$, and $\max(\text{code}(\alpha)) < \kappa$. Fix β' with $\beta < \beta' < \kappa$. Let $\alpha' < \alpha$ with $\max(\text{code}(\alpha')) < \kappa$. Let $n \in \omega$. We must show that there is a proper initial segment $\mathcal{N} \triangleleft J_\kappa^\mathcal{M}$ such that $\mathcal{N}$ is explicitly (α', n) -big above β' . By increasing β' we may assume that $\max(\text{code}(\alpha')) < \beta'$.

Let $\bar{\alpha} = \alpha' + \beta' + 1$. Notice that $\beta' < \max(\text{code}(\bar{\alpha})) \leq \max(\text{code}(\alpha')) + \beta' + 1$. As $\text{cof}(\alpha) = \omega_1$, $\bar{\alpha} < \alpha$. As κ is inaccessible in $\mathcal{M}$, $\max(\text{code}(\bar{\alpha})) < \kappa$. Thus there is a proper initial segment $\mathcal{N} \triangleleft J_\kappa^\mathcal{M}$ such that $\mathcal{N}$ is explicitly $(\bar{\alpha}, 1)$ -big above β . Fix the $\leq$ -least such $\mathcal{N}$. Since $\bar{\alpha}$ is a successor ordinal, there is some ρ in $\mathcal{N}$ such that $(\mathcal{N}, \rho)$ witnesses that $\mathcal{N}$ is explicitly $(\bar{\alpha}, 1)$ -big above β . By definition $\max(\text{code}(\bar{\alpha})) < \rho$. So $\beta' < \rho$. Thus $\mathcal{N}$ is explicitly $(\bar{\alpha}, 1)$ -big above β' .

Now $\alpha' < \bar{\alpha}$. Also, we are assuming that $\max(\text{code}(\alpha')) < \beta'$. So by Lemma 3.15, $\mathcal{N}$ is explicitly (α', n) -big above β' . $\square$

We now turn to criterion (5'). That is, we wish to show that the statement “I am explicitly (α, n) -big above β ” is definable. To begin with we need the following.

Lemma 3.18. *If $\mathcal{M}$ is explicitly (α, n) -big above β , then $\text{code}(\alpha) \in \mathcal{M}$. (Unless α is a limit ordinal of cofinality ω and $n = 0$. In this case we can only say that $\max(\text{code}(\alpha)) \leq \text{OR} \cap \mathcal{M}$.)*

Proof. First suppose $\text{cof}(\alpha) > \omega$. If $n = 0$ then the lemma is true by definition, and if $n > 0$ then we apply lemma 3.13 above.

Now suppose $\text{cof}(\alpha) \leq \omega$. If $n = 1$ then the lemma is true by definition, and if $n > 1$ then we apply lemma 3.13 above.

Next suppose $\alpha = \gamma + 1$ is a successor ordinal and $n = 0$, and $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . Then $\mathcal{M}$ is explicitly (γ, n) -big above β for all n , so $\text{code}(\gamma) \in \mathcal{M}$. By part (d) of Lemma 2.35, $\text{code}(\alpha) \leq \text{code}(\gamma) + 1$. So $\text{code}(\alpha) \in \mathcal{M}$.

Finally, suppose that α is a limit ordinal of cofinality ω and $n = 0$, and $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . We argue by induction on α . For cofinally many $\alpha' < \alpha$, there is a proper initial segment $\mathcal{N}_{\alpha'} \triangleleft \mathcal{M}$ such that $\mathcal{N}_{\alpha'}$ is explicitly $(\alpha', 0)$ -big above β . By induction, for cofinally many $\alpha' < \alpha$, $\text{code}(\alpha') \in \mathcal{M}$. By part (c) of Lemma 2.35, $\max(\text{code}(\alpha)) \leq \text{OR} \cap \mathcal{M}$. $\square$

Suppose that $\mathcal{M}$ is a countable premouse and $\beta \in \text{OR}^{\mathcal{M}}$, and $\alpha < \omega_1^{\omega_1}$ and $s = \text{code}(\alpha)$ and $s \in \mathcal{M}$. We want to show that the statement “I am explicitly $(|s|, n)$ -big above β ” is definable in $\mathcal{M}$. Notice that as $\mathcal{M}$ is countable, the term “ $|s|$ ” cannot mean the same thing in $\mathcal{M}$ as it does in V . So we must be careful about what we mean. To be precise, we want to show that there is a formula ψ in the language of premice such that for all premice $\mathcal{M}$, and all codes $s \in \mathfrak{C} \cap \mathcal{M}$, and all β in $\text{OR}^{\mathcal{M}}$, and all $n \in \omega$, $\mathcal{M}$ is explicitly $(|s|, n)$ -big above β iff $\mathcal{M} \models \psi[s, n, \beta]$.

It suffices to see that a premouse $\mathcal{M}$ can tell when its proper initial segments are big. This is true because the definition involves a “local Σ_1 induction.” We spell out all of the details below.

Definition 3.19. Let $\mathcal{M} = \mathcal{J}_{\gamma}^{\mathcal{M}}$ be a premouse. Then $H^{\mathcal{M}}$ is the function with domain equal to the set of all 4-tuples (ξ, β, s, n) such that $\xi \leq \gamma$, and $\beta \in \mathcal{J}_{\xi}^{\mathcal{M}}$, and $s \in \mathfrak{C}$ is a code, with $\max(s) \leq \text{OR} \cap \mathcal{J}_{\xi}^{\mathcal{M}}$, and $n \in \omega$, and

$$H^{\mathcal{M}} : \text{dom}(H^{\mathcal{M}}) \rightarrow \{0, 1\}$$

is defined by $H^{\mathcal{M}}(\xi, \beta, s, n) = 1$ iff $\mathcal{J}_{\xi}^{\mathcal{M}}$ is explicitly $(|s|, n)$ -big above β . For $\rho \leq \gamma$ we will write $H^{\mathcal{M}} \upharpoonright \rho$ to mean

$$\bigcup_{\xi < \rho} H^{\mathcal{M}} \upharpoonright \mathcal{J}_{\xi}^{\mathcal{M}}.$$

Lemma 3.20. Let $\mathcal{M} = \mathcal{J}_{\gamma}^{\mathcal{M}}$ be a premouse.

- (a) For all $\rho \leq \gamma$, $H^{\mathcal{M}} \upharpoonright \rho$ is Σ_1 definable over $\mathcal{J}_{\rho}^{\mathcal{M}}$, uniformly for all $\mathcal{M}$ and all ρ .

- (b) For all $\rho < \gamma$, $H^{\mathcal{M}} \restriction \rho \in \mathcal{J}_{\rho+1}^{\mathcal{M}}$.
- (c) For all $\rho < \gamma$, $H^{\mathcal{M}} \restriction (\rho + 1) = H^{\mathcal{J}_\rho^{\mathcal{M}}} \in \mathcal{J}_{\rho+1}^{\mathcal{M}}$

Proof. First observe that $H^{\mathcal{N}}$ is Σ_0 definable from the parameter $\mathcal{N}$, uniformly for all premice $\mathcal{N}$. That is, there is a Σ_0 formula θ such that for all $\mathcal{N}$, $h = H^{\mathcal{N}}$ iff $\theta[h, \mathcal{N}]$ holds. $\theta(v_1, v_2)$ just says that v_1 is a function which obeys the appropriate recursive definition, with v_2 as a parameter. The fact that θ can be taken to be Σ_0 uses certain facts about our coding scheme: (See Lemma 2.35.)

1. The relation $|s| < |t|$ is Σ_0 definable.
2. The relation $\text{cof}(|s|) \leq \omega$ is Σ_0 definable.
3. If $s \in \mathfrak{C}$ and $|s|$ is a limit ordinal of cofinality ω , and if we set $\mathfrak{C} \restriction s = \{s' \in \mathfrak{C} : |s'| < |s| \wedge \max(s') \leq \max(s)\}$, then we have that $\{|s'| : s' \in \mathfrak{C} \restriction s\}$ is cofinal in $|s|$.

Now let $\varphi(\xi, \beta, s, n, b)$ be the Σ_1 formula in the language of premice which, when interpreted in the premouse $\mathcal{M}$, asserts:

$$b \in \{0, 1\} \text{ and there exists a function } h \text{ such that } \theta(h, \mathcal{J}_\xi^{\mathcal{M}}) \text{ and } h(\xi, \beta, s, n) = b.$$

We now prove (a), (b), and (c) by simultaneous induction on ρ .

Given (c) for $\rho' < \rho$, it is easy to see that $H^{\mathcal{M}} \restriction \rho(\xi, \beta, s, n) = b$ iff $\mathcal{J}_\rho^{\mathcal{M}} \models \varphi[\xi, \beta, s, n, b]$.

This proves (a).

(b) follows immediately from (a).

We turn to (c). Let $h = H^{\mathcal{M}} \restriction (\rho + 1) - H^{\mathcal{M}} \restriction \rho$. By (b) it suffices to show that $h \in \mathcal{J}_{\rho+1}^{\mathcal{M}}$.

Let

$$X = \{(\beta, s, n) \in \mathcal{J}_\rho^{\mathcal{M}} \mid \mathcal{J}_\rho^{\mathcal{M}} \text{ is explicitly } (|s|, n)\text{-big above } \beta\}.$$

It suffices to show that X is definable over $\mathcal{J}_\rho^{\mathcal{M}}$. Let

$$Y = X \cap \{(\beta, s, n) \mid (\forall \xi < \rho) \mathcal{J}_\xi^{\mathcal{M}} \text{ is not explicitly } (|s|, n)\text{-big above } \beta\}.$$

By (a) it suffices to show that Y is definable over $\mathcal{J}_\rho^{\mathcal{M}}$. But to determine whether or not $(\beta, s, n) \in Y$ only requires looking at some simple first order properties of $\mathcal{J}_\rho^{\mathcal{M}}$ (like whether or not it is active, whether or not it is admissible, and whether or not it has n Woodin cardinals) and then looking at whether or not certain proper initial segments of

$\mathcal{J}_\rho^\mathcal{M}$ are big. By (a) this last task can be done definably over $\mathcal{J}_\rho^\mathcal{M}$. So Y is definable over $\mathcal{J}_\rho^\mathcal{M}$. $\square$

Corollary 3.21. CRITERION (5'). *There is a formula ψ in the language of premice such that for all premice $\mathcal{M}$, and all codes $s \in \mathfrak{C} \cap \mathcal{M}$, and all β in $\text{OR}^\mathcal{M}$, and all $n \in \omega$, $\mathcal{M}$ is explicitly $(|s|, n)$ -big above β iff $\mathcal{M} \models \psi[s, n, \beta]$.*

Proof. This follows from the proof of (c) in the previous lemma. $\square$

Note. If ψ is as above, then we will express the fact that $\mathcal{M} \models \psi[s, n, \beta]$ by saying that $\mathcal{M} \models$ “I am explicitly (α, n) -big above β .”

We will use the above corollary heavily. Our first application of the corollary is to prove a series of lemmas which say that the property of being explicitly (α, n) -big can be reflected downward. We will then go on to use these reflection lemmas to further study the structure of (α, n) -big mice. Among other things, we will learn that the mice we are interested in studying occur below the first mouse with ω Woodin cardinals. (See Proposition 3.25)

Lemma 3.22. *Let $\mathcal{M}$ be a countable, realizable, meek premouse and suppose that λ is a limit cardinal of $\mathcal{M}$. Let $\alpha < \omega_1^{\omega_1}$ with $\text{code}(\alpha) \in J_\lambda^\mathcal{M}$. Let $\beta < \lambda$, and let $n \in \omega$. Suppose that there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β . Then, there is a proper initial segment $\mathcal{N} \triangleleft J_\lambda^\mathcal{M}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β .*

Proof. Let $\kappa < \lambda$ be a cardinal of $\mathcal{M}$ such that $\text{code}(\alpha) \in J_\kappa^\mathcal{M}$ and $\beta < \kappa$. Suppose that $\mathcal{N} \triangleleft \mathcal{M}$ and $\kappa \in \mathcal{N}$ and $\mathcal{N}$ is explicitly (α, n) -big above β , and $\mathcal{N}$ is $\trianglelefteq$ -least with these properties. By Corollary 3.21, there is a formula θ such that $\mathcal{N}$ is the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ such that $\kappa \in \mathcal{N}$ and $\mathcal{N} \models \theta[\text{code}(\alpha), \beta]$. Then, by our first corollary to the Condensation Theorem, Corollary 1.11 on page 12, $\rho_\omega(\mathcal{N}) = \kappa$. So $\mathcal{N} \triangleleft J_\lambda^\mathcal{M}$. $\square$

The next lemma says that the least cardinal which is locally (α, n) -Woodin is not regular.

Lemma 3.23. *Let $\mathcal{M}$ be a countable, realizable, meek premouse and suppose that δ is an inaccessible cardinal of $\mathcal{M}$. Let $\alpha < \omega_1^{\omega_1}$ with $\text{code}(\alpha) \in J_\delta^{\mathcal{M}}$, and let $n \in \omega$. Suppose that δ is locally (α, n) -Woodin in $\mathcal{M}$. Then, there are cofinally many $\bar{\delta} < \delta$ such that $\bar{\delta}$ is a cardinal of $\mathcal{M}$ and $\bar{\delta}$ is locally (α, n) -Woodin in $\mathcal{J}_\delta^{\mathcal{M}}$.*

Proof. Let $\mathcal{N} \triangleleft \mathcal{M}$ be $\leq$ -least such that $\mathcal{N}$ witnesses that δ is locally (α, n) -Woodin in $\mathcal{M}$. So δ is Woodin in $\mathcal{N}$ and $\mathcal{N}$ is the $\leq$ -least initial segment of $\mathcal{M}$ which is explicitly (α, n) -big above δ .

Claim. $\rho_\omega(\mathcal{N}) = \delta$.

Proof of Claim. We want to apply our first corollary to the Condensation Theorem, Corollary 1.11, and this corollary has a rather carefully-worded hypothesis. We need to see that there is some formula θ and some parameter $q \in J_\delta^{\mathcal{M}}$, so that $\mathcal{N}$ is the $\leq$ -least initial segment of $\mathcal{M}$ such that $\delta \in \mathcal{N}$ and $\mathcal{N} \models \theta[q]$. We can take $q = \text{code}(\alpha)$ and we can take $\theta[\text{code}(\alpha)]$ to say: “There is some ordinal δ' such that I am explicitly (α, n) -big above δ' and no proper initial segment of me is.” Clearly $\mathcal{N}$ satisfies this with $\delta' = \delta$. Suppose towards a contradiction that there is some $\mathcal{N}'$ so that $\delta \in \mathcal{N}' \triangleleft \mathcal{N}$, and $\mathcal{N}' \models \theta[\text{code}(\alpha)]$. Fix δ' witnessing this. Then $\delta' < \delta$. Let κ be a cardinal of $\mathcal{M}$ such that $\kappa < \delta$ and the maximum of $\max(\text{code}(\alpha))$ and δ' is below κ . Then $\mathcal{N}'$ is the $\leq$ -least initial segment of $\mathcal{M}$ such that $\kappa \in \mathcal{N}'$ and $\mathcal{N}'$ satisfies “I am explicitly (α, n) -big above δ' .” By Corollary 1.11, $\rho_\omega(\mathcal{N}') = \kappa$. But this is impossible since δ is a cardinal of $\mathcal{M}$. This shows that $\mathcal{N}$ is the $\leq$ -least initial segment of $\mathcal{M}$ such that $\delta \in \mathcal{N}$ and $\mathcal{N} \models \theta[\text{code}(\alpha)]$. By Corollary 1.11, $\rho_\omega(\mathcal{N}) = \delta$. $\diamond$

Now fix $\beta < \delta$. We will find $\bar{\delta}$ with $\beta < \bar{\delta} < \delta$ such that $\bar{\delta}$ is locally (α, n) -Woodin in $\mathcal{J}_\delta^{\mathcal{M}}$. Working in $\mathcal{M}$ where δ is inaccessible, let $X \prec \mathcal{N}$ be a fully elementary submodel of $\mathcal{N}$ such that $|X| < \delta$ and $X \cap \delta$ is a cardinal greater than the maximum of $\max(\text{code}(\alpha))$ and β . Let $\pi : \bar{\mathcal{N}} \rightarrow X$ be the inverse of the collapse map. Let $\bar{\delta} = \text{crit}(\pi)$. Then $\pi(\bar{\delta}) = \delta$ and $\bar{\delta}$ is a cardinal of $\mathcal{M}$. So $\rho_\omega(\bar{\mathcal{N}}) = \bar{\delta}$. So by the Condensation Theorem, $\bar{\mathcal{N}} \triangleleft J_\delta^{\mathcal{M}}$. (Case (b) of the Condensation Theorem cannot occur because $\bar{\delta}$ is a cardinal of $\mathcal{M}$, and so $\mathcal{J}_\delta^{\mathcal{M}}$ is passive.) We have: $\beta < \bar{\delta}$ and $\bar{\delta}$ is a cardinal of $\mathcal{M}$ and $\bar{\delta}$ is a Woodin cardinal of

$\bar{N}$ and $\bar{N}$ is explicitly (α, n) -big above $\bar{\delta}$. So $\bar{N}$ witnesses that $\bar{\delta}$ is locally (α, n) -Woodin in $\mathcal{J}_\delta^\mathcal{M}$. $\square$

Suppose $\text{cof}(\alpha) \leq \omega$. The next lemma says that if $\mathcal{M}$ has a single fixed cardinal δ , such that for cofinally many $\alpha' < \alpha$ and all n , δ is locally (α', n) -Woodin in $\mathcal{M}$, then there is a proper initial segment of $\mathcal{M}$ which is explicitly $(\alpha, 1)$ -big. Compare this with the discussion following Lemma 3.6 on page 37.

Lemma 3.24. *Let $\mathcal{M}$ be a countable, realizable, meek premouse and suppose that δ is an inaccessible cardinal of $\mathcal{M}$. Let $\alpha < \omega_1^{\omega_1}$ with $\text{code}(\alpha) \in J_\delta^\mathcal{M}$. Suppose that $\text{cof}(\alpha) \leq \omega$ and that for cofinally many $\alpha' < \alpha$ and all n , δ is locally (α, n) -Woodin in $\mathcal{M}$. Then, for all $\beta < \delta$, $J_\delta^\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β .*

Proof. Fix $\beta < \delta$. Let $s = \text{code}(\alpha)$. So $s \in J_\delta^\mathcal{M}$ by assumption. Recall that $\mathfrak{C}$ is our set of codes for ordinals less than $\omega_1^{\omega_1}$. Set $\mathfrak{C} \restriction s = \{s' \in \mathfrak{C} : |s'| < |s| \wedge \max(\text{code}(s')) \leq \max(\text{code}(s))\}$.

Claim. *For all $s' \in \mathfrak{C} \restriction s$, and all $n \in \omega$, and all $\beta' < \delta$, there is an ordinal δ' such that $\beta' < \delta' < \delta$ and δ' is a cardinal of $\mathcal{M}$ and δ' is locally (α', n) -Woodin in $J_\delta^\mathcal{M}$.*

Proof. Fix s' , n , and β' . By hypothesis there is an ordinal $\hat{\alpha}$ such that $|s'| < \hat{\alpha} < \alpha$ and there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\delta \in \mathcal{N}$, and δ is a Woodin cardinal of $\mathcal{N}$, and $\mathcal{N}$ is explicitly $(\hat{\alpha}, n)$ -big above δ . By Lemma 3.15, $\mathcal{N}$ is explicitly $(|s'|, n)$ -big above δ . Then the conclusion of the claim follows from the previous lemma. $\diamond$

Since $\mathfrak{C} \restriction s \in J_\delta^\mathcal{M}$ and $J_\delta^\mathcal{M} \models \text{ZFC}$, there is a $\rho < \delta$ such that $\beta < \rho$ and $\max(s) < \rho$ and the above claim is true with ρ in place of δ . By part (e) of Lemma 2.35, $\{|s'| : s' \in \mathfrak{C} \restriction s\}$ is cofinal in α . Thus $(J_\delta^\mathcal{M}, \rho)$ witnesses that $J_\delta^\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . $\square$

Because the definition of (α, n) -big involves a fairly complicated induction, it may not be clear that there are *any* premice which are (α, n) -big. We will now show that if $\mathcal{M}$ is a countable, iterable premouse and $\mathcal{M}$ has ω Woodin cardinals, then $\mathcal{M}$ is $(\omega_1^{\omega_1}, 0)$ -big. The contrapositive of this implies, in particular, that if $\mathcal{M}$ is $(\omega_1^{\omega_1}, 0)$ -small, then $\mathcal{M}$ does not have ω Woodin cardinals, and in particular $\mathcal{M}$ is meek.

Proposition 3.25. *Let $\mathcal{M}$ be a countable, realizable, premouse. For $i \in \omega$, suppose that δ_i is a Woodin cardinal of $\mathcal{M}$, and $\delta_i < \delta_{i+1}$. Then $\mathcal{M}$ is $(\omega_1^{\omega_1}, 0)$ -big.*

Proof. We may assume that no proper initial segment of $\mathcal{M}$ satisfies our hypotheses on $\mathcal{M}$. Thus the δ_i are cofinal in the ordinals of $\mathcal{M}$, and $\mathcal{M}$ is, in particular, meek.

Claim. *For all $\alpha < \omega_1^{\omega_1}$ such that $\text{code}(\alpha) \in \mathcal{M}$, if $\text{code}(\alpha) \in J_{\delta_i}^{\mathcal{M}}$, then for all $\beta < \delta_i$, $J_{\delta_i}^{\mathcal{M}}$ is explicitly (α, n) -big above β for all n .*

Proof of Claim. By Lemmas 3.13 and 3.14, it suffices to prove the claim with “for $n = 1$ ” in place of “for all n ”. We do this by induction on α . So let α be the least counterexample. Fix i such that $\text{code}(\alpha) \in J_{\delta_i}^{\mathcal{M}}$, and fix $\beta < \delta_i$.

First suppose that $\text{cof}(\alpha) \leq \omega$. To conclude that $J_{\delta_i}^{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above β , we will apply the previous lemma with $\delta = \delta_i$. So we must see that the hypotheses of this lemma are satisfied. By Lemma 2.35, there are cofinally many $\alpha' < \alpha$ such that $\text{code}(\alpha') \in J_{\delta_i}^{\mathcal{M}}$. Fix such an α' and fix $n \geq 1$. By induction, $J_{\delta_{i+n}}^{\mathcal{M}}$ is explicitly $(\alpha', 1)$ -big above δ_{i+n-1} . So $J_{\delta_{i+n}}^{\mathcal{M}}$ is explicitly (α', n) -big above δ_i . Thus the hypotheses of the previous lemma are satisfied, and our claim is proved.

Now suppose that $\text{cof}(\alpha) = \omega_1$. Let κ be a measurable cardinal of $\mathcal{M}$ such that $\delta_i < \kappa < \delta_{i+1}$. Then for all $\alpha' < \alpha$, if $\text{code}(\alpha') \in J_{\kappa}^{\mathcal{M}}$, then by induction, $J_{\delta_{i+1}}^{\mathcal{M}}$ is explicitly $(\alpha' + 1, 1)$ -big above δ_i . By Lemma 3.13, for all n , $J_{\delta_{i+1}}^{\mathcal{M}}$ is explicitly (α', n) -big above δ_i . By Lemma 3.22, for all $\alpha' < \alpha$, if $\text{code}(\alpha') \in J_{\kappa}^{\mathcal{M}}$, then for all n there is a proper initial segment $\mathcal{N} \triangleleft J_{\kappa}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α', n) -big above δ_i . Thus $J_{\delta_{i+1}}^{\mathcal{M}}$ is explicitly $(\alpha, 0)$ -big above δ_i . Thus $J_{\delta_{i+1}}^{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above β . By Lemma 3.22, there is a proper initial segment $\mathcal{N} \triangleleft J_{\delta_i}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly $(\alpha, 1)$ -big above β . $\diamond$

Now, let κ be the least measurable cardinal of $\mathcal{M}$. Let $\alpha < \omega_1^{\omega_1}$ be such that $\text{code}(\alpha) \in J_{\kappa}^{\mathcal{M}}$. Then for all n , $J_{\delta_0}^{\mathcal{M}}$ is explicitly (α, n) -big (above 0). By Lemma 3.22, for all n there is a proper initial segment $\mathcal{N} \triangleleft J_{\kappa}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α, n) -big. Thus $\mathcal{M}$ is explicitly $(\omega_1^{\omega_1}, 0)$ -big. $\square$

The next lemma says that the least mouse which is (α, n) -big is not $(\alpha, n + 1)$ -big, at least if $\max(\text{code}(\alpha))$ is sufficiently small. This is a somewhat technical lemma which will be used several times in the sequel.

Lemma 3.26. *Let $\mathcal{M}$ be a countable, realizable, meek premouse, and $\beta \in \text{OR}^{\mathcal{M}}$. Let $\alpha < \omega_1^{\omega_1}$, let $n \in \omega$, and suppose that $\mathcal{M}$ is explicitly (α, n) -big above β . Suppose that $(\alpha', n') <_{\text{lex}} (\alpha, n)$ and $\max(\text{code}(\alpha')) < \beta$. Then there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is explicitly (α', n') -big above β .*

Proof. If $\text{cof}(\alpha) > \omega$ and $n = 0$ this follows from the proof of Lemma 3.15. If $\text{cof}(\alpha) \leq \omega$ and $n = 0$ the statement follows from the definition of explicitly $(\alpha, 0)$ -big, and Lemma 3.15.

So now let $n \geq 0$ and suppose that $\mathcal{M}$ is explicitly $(\alpha, n + 1)$ -big above β , and $\max(\text{code}(\alpha)) < \beta$, and we will show that there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β . Clearly we may assume that no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, n + 1)$ -big above β .

If $\text{cof}(\alpha) \leq \omega$ and $n = 0$ this follows from the definitions. So suppose otherwise. Then there is an ordinal $\delta \in \mathcal{M}$ with $\beta < \delta$ such that δ is a Woodin cardinal of $\mathcal{M}$, and $\mathcal{M}$ is explicitly (α, n) -big above δ . We will show that there is a proper initial segment $\mathcal{N} \triangleleft J_{\delta}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β .

First suppose that $\text{cof}(\alpha) \leq \omega$. Then by case hypothesis we have that $n \geq 1$. There are n ordinals, $\delta_1, \dots, \delta_n$, with $\delta = \delta_1 < \delta_2 < \dots < \delta_n$ such that each δ_i is a Woodin cardinal of $\mathcal{M}$, and $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ_n . In particular, $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above δ_n . Thus, for cofinally many $\alpha' < \alpha$ and all n' , δ_n is locally (α', n') -Woodin in $\mathcal{M}$. By Lemma 3.24, $J_{\delta_n}^{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above β' , for all $\beta' < \delta_n$. Thus $J_{\delta_n}^{\mathcal{M}}$ is explicitly (α, n) -big above β . By Lemma 3.22, there is a proper initial segment $\mathcal{N} \triangleleft J_{\delta_1}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β .

Next suppose that $\text{cof}(\alpha) > \omega$. Then there are $n + 1$ ordinals, $\delta_1, \dots, \delta_{n+1}$, with $\delta = \delta_1 < \delta_2 < \dots < \delta_{n+1}$ such that each δ_i is a Woodin cardinal of $\mathcal{M}$, and $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above δ_{n+1} . By definition, $\mathcal{M}$ is an active premouse, and letting κ be the critical point of the last extender from the $\mathcal{M}$ sequence we have that $\delta_{n+1} < \kappa$.

To simplify the notation, let us assume that $n > 0$. Let μ be a measurable cardinal of $\mathcal{M}$ such that $\delta_n < \mu < \delta_{n+1}$. Let $\mathcal{P} \triangleleft J_{\delta_{n+1}}^{\mathcal{M}}$ be such that $\mathcal{P}$ is active, and μ is the critical point of the last extender on the $\mathcal{P}$ -sequence. We will show that $\mathcal{P}$ is explicitly $(\alpha, 0)$ -big above δ_n . Let $\alpha' < \alpha$ with $\max(\text{code}(\alpha')) < \mu$, and let $n' \in \omega$. We must show that

there is an initial segment $\mathcal{Q} \triangleleft J_\mu^{\mathcal{M}}$ such that $\mathcal{Q}$ is explicitly (α', n') -big above δ_n . But $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above δ_{n+1} , so there is a proper initial segment of $J_\kappa^{\mathcal{M}}$ which is explicitly (α', n') -big above δ_{n+1} , and so in particular above δ_n . So by Lemma 3.22, there is an initial segment $\mathcal{Q} \triangleleft J_\mu^{\mathcal{M}}$ such that $\mathcal{Q}$ is explicitly (α', n') -big above δ_n .

Thus $\mathcal{P}$ is explicitly $(\alpha, 0)$ -big above δ_n , and so $\mathcal{P}$ is explicitly (α, n) -big above β . By Lemma 3.22, there is a proper initial segment $\mathcal{N} \triangleleft J_{\delta_1}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α, n) -big above β . $\square$

The next lemma says that if α is greatest so that $\mathcal{M}$ is $(\alpha, 0)$ -big, then $\max(\text{code}(\alpha))$ is below the least measurable cardinal of $\mathcal{M}$. This is another technical lemma which will be used several times in the sequel.

Lemma 3.27. *Let $\mathcal{M}$ be a countable, realizable premouse, and $\beta \in \text{OR}^{\mathcal{M}}$. Let $\alpha < \omega_1^{\omega_1}$ and suppose that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . Let $\kappa \in \text{OR}^{\mathcal{M}}$ and suppose that $\beta < \kappa$, and κ is a cardinal of $\mathcal{M}$, and κ is the critical point of an extender from the $\mathcal{M}$ -sequence. Then either $\max(\text{code}(\alpha)) < \kappa$, or $\mathcal{M}$ is explicitly $(\alpha^*, 0)$ -big above β , for some $\alpha^* > \alpha$ such that $\text{cof}(\alpha^*) = \omega_1$.*

Proof. Suppose $\max(\text{code}(\alpha)) \geq \kappa$. Let $\alpha^* \leq \omega_1^{\omega_1}$ be least such that $\alpha^* > \alpha$ and $\max(\text{code}(\alpha^*)) < \kappa$. (There is such an α^* since we have set $\text{code}(\omega_1^{\omega_1}) = \emptyset$, and $\max(\emptyset) = 0$.) Then, by Lemma 2.35, α^* is a limit ordinal of uncountable cofinality. We will show that $\mathcal{M}$ is explicitly $(\alpha^*, 0)$ -big above β . For all $\alpha' < \alpha^*$, if $\max(\text{code}(\alpha')) < \kappa$, then $\alpha' < \alpha$, and $\max(\text{code}(\alpha')) < \max(\text{code}(\alpha))$. By Lemma 3.16, for all $\alpha' < \alpha^*$, if $\max(\text{code}(\alpha')) < \kappa$, then $\mathcal{M}$ is explicitly (α', n) -big above β for all n . Since we may assume that $\mathcal{M}$ does not have ω Woodin cardinals, we must have that for all $\alpha' < \alpha^*$, if $\max(\text{code}(\alpha')) < \kappa$, then for all n , there is a proper initial segment of $\mathcal{M}$ that is explicitly (α', n) -big above β . By Lemma 3.22, for all $\alpha' < \alpha^*$, if $\max(\text{code}(\alpha')) < \kappa$, then for all n , there is a proper initial segment of $\mathcal{J}_\kappa^{\mathcal{M}}$ that is explicitly (α', n) -big above β . Thus letting $\mathcal{N} \trianglelefteq \mathcal{M}$ be such that $\mathcal{N}$ is active and κ is the critical point of the last extender from the $\mathcal{N}$ sequence, we have that $\mathcal{N}$ witnesses that $\mathcal{M}$ is explicitly $(\alpha^*, 0)$ -big above β . $\square$

In Corollary 3.21 above, we showed that the statement “I am explicitly (α, n) -big above β ” is definable. We then examined some consequences of this. We now turn to another

consequence: If $\text{cof}(\alpha) \leq \omega$ and $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big, then there is a sequence of ordinals witnessing this. We call this sequence an α -ladder. (See Figure 1 on page 39.) It turns out to be important that this α -ladder is an element of $\mathcal{M}$. We turn to this now. The technical details concerning the organization of the α -ladder in the following definition are motivated by our use of the α -ladder in the next section.

Definition 3.28. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Suppose that $2 \leq \alpha < \omega_1^{\omega_1}$ and that $\text{cof}(\alpha) \leq \omega$. Let f be a function. Then f is an α -ladder above β relative to $\mathcal{M}$ iff the following conditions hold. There are two cases. First suppose that $\alpha = \alpha' + 1$ is a successor ordinal. (See Figure 1 on page 39.) Then:

- (i) $\text{dom}(f) = \omega$, and
- (ii) for all n , $f(n) = (\delta_n, \gamma_n)$, with $\beta < \delta_n < \gamma_n < \delta_{n+1} < \text{OR}^{\mathcal{M}}$, and
- (iii) δ_n is a cardinal of $\mathcal{M}$, and $J_{\gamma_n}^{\mathcal{M}} \models \text{“}\delta_n \text{ is a Woodin cardinal,”}$ and γ_n is least such that $J_{\gamma_n}^{\mathcal{M}}$ is explicitly $(\alpha', 2n)$ -big above δ_n , and
- (iv) $\max(\text{code}(\alpha)) < \delta_0$.

Next suppose that α is a limit ordinal of cofinality ω . Then:

- (i) $\text{dom}(f)$ is a countable limit ordinal $\mu \leq \max(\text{code}(\alpha))$, and
- (ii) for all $\xi \in \text{dom}(f)$ we have that: $f(\xi) = (\delta_\xi, \gamma_\xi, s_\xi)$ with $\beta < \delta_\xi < \gamma_\xi < \text{OR}^{\mathcal{M}}$, and $\delta_\xi > \sup \{ \gamma_\eta \mid \eta < \xi \}$, and $s_\xi \in \mathfrak{C}$ is a code for an ordinal less than ω_1^ω , and, letting $\alpha_\xi = |s_\xi|$, we have that $\alpha_\xi < \alpha$ and
- (iii) δ_ξ is a cardinal of $\mathcal{M}$, and $J_{\gamma_\xi}^{\mathcal{M}} \models \text{“}\delta_\xi \text{ is a Woodin cardinal,”}$ and γ_ξ is least such that $J_{\gamma_\xi}^{\mathcal{M}}$ is explicitly $(\alpha_\xi, 2)$ -big above δ_ξ , and
- (iv) for all ξ , $\max(s_\xi) \leq \max(\text{code}(\alpha)) < \delta_0$, and
- (v) $\{ \alpha_\xi \mid \xi \in \text{dom}(f) \}$ is cofinal in α .

Definition 3.29. Suppose f is an α -ladder above β relative to $\mathcal{M}$. Then, using the notation from the previous definition, we define

$$\sup(f) = \sup \{ \gamma_\xi \mid \xi \in \text{dom}(f) \} = \sup \{ \delta_\xi \mid \xi \in \text{dom}(f) \}.$$

Lemma 3.30. Let $\mathcal{M}$ be a countable, realizable, meek premouse, and $\beta \in \text{OR}^{\mathcal{M}}$. Suppose that $2 \leq \alpha < \omega_1^{\omega_1}$ and that α is a successor ordinal, or a limit ordinal of cofinality ω .

Suppose that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β , and that $\mathcal{M}$ itself witnesses this. Let ρ be the least ordinal such that $(\mathcal{M}, \rho)$ witnesses this. Then there is a function f such that:

- (i) f is an α -ladder above β relative to $\mathcal{M}$, and
- (ii) $f \in \mathcal{M}$, and
- (iii) $\rho = \sup(f)$.

Proof. We will assume that α is a limit ordinal of cofinality ω . If α is a successor ordinal then the proof is similar but simpler.

Recall that ρ is a limit cardinal of $\mathcal{M}$, and notice that there are cofinally many $\gamma < \rho$ such that $J_\gamma^\mathcal{M} \models \text{ZFC}$. Let $s = \text{code}(\alpha)$. Then $s \in J_\rho^\mathcal{M}$ by definition. Recall that $\mathfrak{C}$ is our set of codes for ordinals below $\omega_1^{\omega_1}$. Set $\mathfrak{C} \restriction s = \{s' \in \mathfrak{C} : |s'| < |s| \wedge \max(s') \leq \max(s)\}$. Fix some $\gamma < \rho$ such that $\mathfrak{C} \restriction s \in J_\gamma^\mathcal{M}$ and $J_\gamma^\mathcal{M} \models \text{ZFC}$. We work in $J_\gamma^\mathcal{M}$ for a moment. Let μ be the cofinality, in $J_\gamma^\mathcal{M}$, of the wellorder $(\mathfrak{C} \restriction s, <)$. It is easy to see that $\mu \leq \max(s)$. Fix a sequence $\langle s_\xi \mid \xi < \mu \rangle \in J_\gamma^\mathcal{M}$ which is increasing and cofinal in $\mathfrak{C} \restriction s$.

Now we work in $J_\rho^\mathcal{M}$. For $\xi < \mu$ let δ_ξ be the least cardinal of $\mathcal{M}$ greater than $\max(s)$ which is locally $(|s_\xi|, 2)$ -Woodin in $J_\rho^\mathcal{M}$. By assumption on $\mathcal{M}$, there is such a $\delta_\xi < \rho$. Let γ_ξ be the least ordinal greater than δ_ξ such that $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$ witnesses that δ_ξ is locally $(|s_\xi|, 2)$ -Woodin. So δ_ξ is Woodin in $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$ and $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$ is explicitly $(|s_\xi|, 2)$ -big above δ_ξ . By assumption on $\mathcal{M}$, $\delta_\xi < \gamma_\xi < \rho$. By Corollary 3.21, the function $f = \langle \delta_\xi, \gamma_\xi, s_\xi \mid \xi < \mu \rangle$ is definable over $J_\rho^\mathcal{M}$. So $f \in \mathcal{M}$. We will show that f is an α -ladder above β relative to $\mathcal{M}$, and that $\sup(f) = \rho$. We start by showing that the sequence of $\langle \delta_\xi \rangle$ is strictly increasing.

Claim 1. *Let $\eta < \xi < \mu$. Then $\delta_\eta < \gamma_\eta < \delta_\xi$.*

Proof of Claim 1. Since $\max(s) < \delta_\xi$, it follows from Lemma 3.26 that for all $s' \in \mathfrak{C} \restriction s$, if $|s'| < |s_\xi|$ then there is a *proper* initial segment of $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$ which is explicitly $(|s'|, 2)$ -big above δ_ξ . Since δ_ξ is inaccessible in $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$, Lemma 3.23 for $\mathcal{J}_{\gamma_\xi}^\mathcal{M}$ and δ_ξ implies that $\delta_\eta < \gamma_\eta < \delta_\xi$. $\diamond$

It then follows from the leastness of ρ that $\sup(f) = \rho$. To complete our proof we must show that the sequence of δ_ξ does not contain any of its limit point.

Claim 2. *Let $\xi < \mu$ and suppose that ξ is a limit ordinal. Then $\delta_\xi > \sup \{ \delta_\eta \mid \eta < \xi \}$.*

Proof of Claim 2. We have that $\xi < \mu \leq \max(s) < \delta_\xi$, and δ_ξ is regular in $\mathcal{J}_{\gamma_\xi}^{\mathcal{M}}$, and $f \restriction \xi \in \mathcal{J}_{\gamma_\xi}^{\mathcal{M}}$. $\diamond$

This completes the proof of the lemma. $\square$

We now turn to yet another consequence of the fact that the statement “I am explicitly (α, n) -big” is definable. We have mentioned several times that the property of being explicitly (α, n) -big does not satisfy our criterion (4). That is to say, if $\mathcal{M}$ is explicitly (α, n) -big it does not follow that $\mathcal{M}$ is explicitly (α', n') -big for all $(\alpha', n') <_{\text{lex}} (\alpha, n)$. We proved four lemmas above which gave approximations to this condition though. (See Lemmas 3.13 through 3.16.) Another approximation is that our criterion (4') is satisfied. That is the content of the next lemma.

Lemma 3.31. CRITERION (4'). *Suppose that $\mathcal{M}$ is a countable, realizable premouse, and $(1, 0) <_{\text{lex}} (\alpha_0, n_0) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$, and $\beta \in \mathcal{M}$, and $\mathcal{M}$ is explicitly (α_0, n_0) -big above β . Then for all $(\alpha', n') <_{\text{lex}} (\alpha_0, n_0)$, there is a countable premouse $\mathcal{N}'$ such that*

- (a) $\mathcal{N}'$ is a linear iterate of some initial segment $\mathcal{N} \leq \mathcal{M}$,
- (b) all critical points used in the iteration to $\mathcal{N}'$ are above β ,
- (c) $\mathcal{N}'$ is realizable, and
- (d) $\mathcal{N}'$ is explicitly (α', n') -big above β .

Proof. Let (α, n) be the lexicographically least pair such that $(\alpha', n') \leq_{\text{lex}} (\alpha, n)$ and $\mathcal{M}$ is explicitly (α, n) -big above β . If $(\alpha', n') = (\alpha, n)$ then we may set $\mathcal{N}' = \mathcal{M}$ and we are done. So suppose that $(\alpha', n') <_{\text{lex}} (\alpha, n)$. Then by Lemma 3.13 $n = 0$, and so by definition α is not a limit ordinal of cofinality ω , and by Lemma 3.14 α is not a successor ordinal. So α is a limit ordinal of cofinality ω_1 . Let $\mathcal{N} \leq \mathcal{M}$ witness that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . By definition, $\mathcal{N}$ is active. Let κ be the critical point of the last extender from the $\mathcal{N}$ -sequence. By definition, $\beta < \kappa$. Let $\mathcal{N}'$ be the result of iterating $\mathcal{N}$ linearly, via its last extender, $\alpha' + 1$ -times. We will show that $\mathcal{N}'$ is explicitly (α', n') -big above β . Let $i : \mathcal{N} \rightarrow \mathcal{N}'$ be the iteration map. So $\kappa = \text{crit}(i)$. If $\alpha < \omega_1^{\omega_1}$ then let $s = \text{code}(\alpha)$. By definition, $s \in J_\kappa^{\mathcal{N}}$. Let φ be a Σ_1 formula such that for any premouse $\mathcal{P}$, and any $\xi, \bar{\beta}, \bar{s}, \bar{n}$, $\mathcal{P} \models \varphi[\xi, \bar{\beta}, \bar{s}, \bar{n}]$ iff $\mathcal{J}_\xi^{\mathcal{P}}$ is explicitly $(|\bar{s}|, \bar{n})$ -big above $\bar{\beta}$.

First suppose that $\alpha < \omega_1^{\omega_1}$. Then

$$\mathcal{N} \models (\forall \bar{s} \in J_\kappa^{\mathcal{N}}, \forall \bar{n})[|\bar{s}| < |s| \implies (\exists \xi < \kappa) \varphi(\xi, \beta, \bar{s}, \bar{n})].$$

So

$$\mathcal{N}' \models (\forall \bar{s} \in J_{i(\kappa)}^{\mathcal{N}'}, \forall \bar{n})[|\bar{s}| < |s| \implies (\exists \xi < i(\kappa)) \varphi(\xi, \beta, \bar{s}, \bar{n})].$$

Let $s' = \text{code}(\alpha')$. Then $s' \in J_{i(\kappa)}^{\mathcal{N}'}$. So $\mathcal{N}'$ is explicitly (α', n') -big above β .

If $\alpha = \omega_1^{\omega_1}$ then

$$\mathcal{N} \models (\forall \bar{s} \in J_\kappa^{\mathcal{N}}, \forall \bar{n})[(\exists \xi < \kappa) \varphi(\xi, \beta, \bar{s}, \bar{n})]$$

and the argument is similar. $\square$

Finally, we turn to criterion 5. That is, we will show that the property of being (α, n) -big is preserved by non-dropping iterations. As in the proof of the previous lemma, it is easy to see that the property of having a proper initial segment which is explicitly (α, n) -big is preserved by non-dropping iterations. So we want to see that if $\mathcal{M}$ itself witnesses that $\mathcal{M}$ is explicitly (α, n) -big, then the same is true of a non-dropping iterate of $\mathcal{M}$. We will focus on the one difficult case, which is the case $\text{cof}(\alpha) \leq \omega$ and $n = 1$. This turns out to be somewhat tricky because it involves the preservation of admissibility under Σ_0 ultrapowers. We do not know whether admissibility is preserved under Σ_0 ultrapowers in general, but the next lemma says that it is preserved given a few extra assumptions.

The following lemma is due jointly to the author and his PhD thesis advisor, John Steel.

Lemma 3.32. *Let $M = J_\xi^A$, where A is some set in M . Suppose that $M \models \text{KP}$. Suppose that there is an ordinal $\rho \in M$ such that*

- (a) ρ is the largest cardinal of M .
- (b) For all $x \in J_\rho^A$, $\mathcal{P}(x) \cap \mathcal{M} \subset J_\rho^A$.

Let G be a (κ, λ) -extender over M , with $\kappa < \rho$. Suppose that for all $a \in [\lambda]^{<\omega}$, G_a is $\Sigma_1(M)$. Suppose that $\text{Ult}_0(M, G)$ is wellfounded, and let $i = i_G$ so that $\text{Ult}_0(M, G) = J_\eta^{i(A)}$ for some η . Finally suppose that i is continuous at ρ . Then $\text{Ult}_0(M, G) \models \text{KP}$.

Proof. Write $\text{Ult} = \text{Ult}_0(M, G)$. Recall that i is a cofinal Σ_1 embedding. Let φ be a Σ_0 formula of the Language of Set Theory, and let $p \in \text{Ult}$. Suppose that $(\forall x \in J_{i(\rho)}^{i(A)})(\exists y \in$

Ult) such that $\text{Ult} \models \varphi[x, y, p]$. It suffices to show that there is an ordinal $\gamma \in \text{Ult}$ such that $(\forall x \in J_{i(\rho)}^{i(A)})(\exists y \in J_\gamma^{i(A)})$ such that $\text{Ult} \models \varphi[x, y, p]$. (To see that this is sufficient, use the fact that, since $i(\rho)$ is the largest cardinal of Ult , for all domains $D \in \text{Ult}$, there is a function $h \in \text{Ult}$ such that $h : J_{i(\rho)}^{i(A)} \xrightarrow{\text{onto}} D$.)

Let $a \in [\lambda]^{<\omega}$ and let $h \in M$ be a function with $\text{dom } h = [\kappa]^{|a|}$ so that $p = [a, h]_G^M$. First we will show that there is an ordinal $\gamma \in \text{Ult}$ which works for all x which are generated by a . Let

$$B = \{ f \in J_\rho^A \mid f \text{ is a function with } \text{dom } f = [\kappa]^{|a|} \}.$$

Then $B \in M$, and, using Σ_0 Los's Theorem and the fact that the range of i is cofinal in OR^{Ult} we have

$$(\forall f \in B)(\exists \gamma \in M) [\{ z \in [\kappa]^{|a|} \mid (\exists y \in J_\gamma^A) M \models \varphi[f(z), y, h(z)] \} \in G_a].$$

Now, using the fact the G_a is $\Sigma_1(M)$ and the fact that $M \models \text{KP}$, we get that there is a fixed ordinal $\gamma \in M$ such that

$$(\forall f \in B) [\{ z \in [\kappa]^{|a|} \mid (\exists y \in J_\gamma^A) M \models \varphi[f(z), y, h(z)] \} \in G_a].$$

Fix such an ordinal γ . Now we will show that this same ordinal works for elements of Ult which are generated by coordinates other than a .

Claim. *Let $b \in [\kappa]^{<\omega}$. Let f be any function in J_ρ^A with $\text{dom } f = [\kappa]^{|a \cup b|}$. Then,*

$$\{ z \in [\kappa]^{|a \cup b|} \mid (\exists y \in J_\gamma^A) M \models \varphi[f(z), y, h(z_{a \cup b}^a)] \} \in G_{a \cup b}.$$

Proof of Claim. Suppose otherwise. Let

$$X = \{ z \in [\kappa]^{|a \cup b|} \mid (\forall y \in J_\gamma^A) M \models \neg \varphi[f(z), y, h(z_{a \cup b}^a)] \}.$$

So $X \in G_{a \cup b}$. Let $Y = \{ z_{a \cup b}^a \mid z \in X \}$. Then $Y \in G_a$. Let $\pi : Y \rightarrow X$ be defined by $\pi(w) =$ the lexicographically least $z \in X$ such that $z_{a \cup b}^a = w$. Now let $\bar{f} : [\kappa]^{|a|} \rightarrow J_\rho^A$ be defined by

$$\bar{f}(w) = \begin{cases} f(\pi(w)) & \text{if } w \in Y; \\ \emptyset & \text{otherwise.} \end{cases}$$

Then $\bar{f} \in J_\rho^A$ so $\bar{f} \in B$. Thus

$$\{ w \in [\kappa]^{|a|} \mid (\exists y \in J_\gamma^A) M \models \varphi[\bar{f}(w), y, h(w)] \} \in G_a.$$

Fix $w \in Y$ such that $(\exists y \in J_\gamma^A)M \models \varphi[\bar{f}(w), y, h(w)]$. Let $z = \pi(w)$. Then $z \in X$. But this contradicts the fact that

$$(\exists y \in J_\gamma^A)M \models \varphi[f(z), y, h(z_{a \cup b}^a)].$$

◇

To finish the proof, let $x \in J_{i(\rho)}^{i(A)}$. We will show that $(\exists y \in J_{i(\gamma)}^{i(A)}) \text{Ult} \models \varphi[x, y, p]$.

Using our assumption that i is continuous at ρ , we have that $x \in J_{i(\beta)}^{i(A)}$ for some $\beta < \rho$. Thus, there is some function $f \in M$ such that $\text{ran}(f) \subset J_\beta^A$, and some $b \in [\lambda]^{<\omega}$ so that $x = [a \cup b, f]_G^M$. Using our assumption that for all $w \in J_\rho^A$, $\mathcal{P}(w) \cap \mathcal{M} \subset J_\rho^A$, we have that $f \in J_\rho^A$. Then, using the Claim and Σ_0 Los's Theorem, we are done. □

The next lemma states that admissibility is preserved under direct limits.

Lemma 3.33. *Let λ be a limit ordinal. Let $\langle M_\xi \mid \xi < \lambda \rangle$ be a sequence of admissible sets. For $\xi < \eta < \lambda$, suppose $i_{\xi, \eta}$ is a Σ_1 embedding of M_ξ into M_η . Suppose that for $\xi < \eta < \nu < \lambda$, $i_{\xi, \nu} = i_{\eta, \nu} \circ i_{\xi, \eta}$. Suppose that the direct limit of this system is wellfounded, and let M_λ be the (transitive collapse of) the direct limit. Then M_λ is admissible.*

Proof. For $\xi < \lambda$, let $i_{\xi, \lambda}$ be the direct limit map. Let $a, p \in M_\lambda$ and let φ be a Σ_1 formula. Suppose that $M \models (\forall x \in a)(\exists y)\varphi[x, y, p]$. Let $\xi < \lambda$ be large enough that both a and p are in the range of $i_{\xi, \lambda}$. Let $\bar{a}$ be such that $i_{\xi, \lambda}(\bar{a}) = a$ and let $\bar{p}$ be such that $i_{\xi, \lambda}(\bar{p}) = p$. Then $M_\xi \models (\forall x \in \bar{a})(\exists y)\varphi[x, y, \bar{p}]$. Since M_ξ is admissible, there is a $\bar{b} \in M_\xi$ such that $M_\xi \models (\forall x \in \bar{a})(\exists y \in \bar{b})\varphi[x, y, \bar{p}]$. Let $b = i_{\xi, \lambda}(\bar{b})$. Then $M_\lambda \models (\forall x \in a)(\exists y \in b)\varphi[x, y, p]$. Thus M_λ is admissible. □

Recall that our current goal is to prove that if $\mathcal{M}$ is explicitly (α, n) -big above δ , and $i : \mathcal{M} \rightarrow \mathcal{M}'$ is a non-dropping iteration map, then $\mathcal{M}'$ is explicitly (α, n) -big above $i(\delta)$. This is pretty easy to see in all but one case. The difficult case is the case in which $\text{cof}(\alpha) \leq \omega$, and $n = 1$, and no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ . In this case we have that $\mathcal{M}$ is admissible, and we must see that $\mathcal{M}'$ is admissible. The previous two lemmas will give this to us, if we add one more hypothesis. The following lemma gives the details.

Lemma 3.34. *Let $\mathcal{M}$ be a countable, realizable, pre-mouse, and $\delta \in \text{OR}^{\mathcal{M}}$. Let $\alpha < \omega_1^{\omega_1}$ and suppose that $\text{cof}(\alpha) \leq \omega$. Suppose that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ and no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ . Furthermore, suppose that there is some ordinal $\beta < \delta$ such that $\max(\text{code}(\alpha)) < \beta$. Let $\mathcal{T}$ be a 0-maximal iteration tree on $\mathcal{M}$, and suppose that $\mathcal{T}$ is above β . Then $\forall \xi < \text{lh}(\mathcal{T})$, if $i_{0,\xi}$ exists, then $\mathcal{M}_\xi^{\mathcal{T}}$ is explicitly $(\alpha, 1)$ -big above $i_{0,\xi}(\delta)$.*

Proof. By induction on ξ . For $\xi < \text{lh } \mathcal{T}$, we will write $\mathcal{M}_\xi$ instead of $\mathcal{M}_\xi^{\mathcal{T}}$. Suppose that $i_{0,\xi}$ exists.

By hypothesis, there is an ordinal $\rho \in \mathcal{M}$ such that $\delta < \rho$ and $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ . Recall that this means that ρ is a limit cardinal of $\mathcal{M}$ and that:

- (i) $\mathcal{M}$ is an admissible set, and
- (ii) $\text{code}(\alpha) \in J_\rho^{\mathcal{M}}$, and
- (iii) for cofinally many $\alpha' < \alpha$, and all $n' \in \omega$, and cofinally many $\delta' < \rho$ we have that:
 - (a) δ' is a cardinal of $\mathcal{M}$, and
 - (b) δ' is locally (α', n') -Woodin in $\mathcal{J}_\rho^{\mathcal{M}}$.

Let $\kappa_0 = \text{crit}(i_{0,\xi})$. Let $s = \text{code}(\alpha)$. So we are assuming that $\max(s) < \beta < \kappa_0$.

We will show that $(\mathcal{M}_\xi, i_{0,\xi}(\rho))$ witnesses that $\mathcal{M}_\xi$ is explicitly $(\alpha, 1)$ -big above $i_{0,\xi}(\delta)$.

It is easy to see that:

- (ii) $s \in J_{i_{0,\xi}(\rho)}^{\mathcal{M}_\xi}$, and
- (iii) for cofinally many $\alpha' < \alpha$, and all $n' \in \omega$, and cofinally many $\delta' < i_{0,\xi}(\rho)$ we have that:
 - (a) δ' is a cardinal of $\mathcal{M}_\xi$, and
 - (b) δ' is locally (α', n') Woodin in $J_{i_{0,\xi}(\rho)}^{\mathcal{M}_\xi}$.

So we have only to see that $\mathcal{M}_\xi$ is an admissible set.

By induction, for all $\nu \in [0, \xi)_T$, $\mathcal{M}_\nu$ is an admissible set.

If ξ is a limit ordinal then $\mathcal{M}_\xi$ is an admissible set by Lemma 3.33. So suppose that $\xi = \eta + 1$. Let $\nu = T\text{-pred}(\xi)$. By induction $\mathcal{M}_\nu$ is admissible. Let $G = E_\eta^{\mathcal{T}}$, and let $i = i_{\nu,\xi}$, and let $\kappa = \text{crit}(i)$. We can conclude from Lemma 3.32 that $\mathcal{M}_\xi$ is admissible,

if we can see that $\mathcal{M}_\nu$ and G and i and κ satisfy the hypotheses of that lemma. Let $\rho' = i_{(0,\nu)}(\rho)$. Then ρ' is a limit cardinal of $\mathcal{M}_\nu$, and since no proper initial segment of $\mathcal{M}_\nu$ is explicitly $(\alpha, 1)$ -big above $i_{(0,\nu)}(\delta)$, we have that $\mathcal{M}_\nu$ is the first admissible structure over $\mathcal{J}_{\rho'}^{\mathcal{M}_\nu}$. Thus ρ' is the largest cardinal of $\mathcal{M}_\nu$ and there are no extenders on the $\mathcal{M}_\nu$ sequence with index above ρ' . This implies that $\kappa < \rho'$. Since premice are strongly acceptable we have that for all $x \in J_{\rho'}^{\mathcal{M}_\nu}$, $\mathcal{P}^{\mathcal{M}_\nu}(x) \in J_{\rho'}^{\mathcal{M}_\nu}$. Since $\mathcal{T}$ is a 0-maximal tree, G is close to $\mathcal{M}_\nu$, and so for all $a \in [\text{lh}(G)]^{<\omega}$, G_a is $\Sigma_1(\mathcal{M}_\nu)$. So to complete our proof it suffices to see that i is continuous at ρ' . This follows from the next claim.

Claim 3. $\text{cof}^{\mathcal{M}_\nu}(\rho') < \kappa$.

Proof of Claim 3. As there is no proper initial segment of $\mathcal{M}$ which is explicitly $(\alpha, 1)$ -big above δ , ρ is the least ordinal such that $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ . By Lemma 3.30, there is a function f such that:

- (i) f is an α -ladder above δ relative to $\mathcal{M}$, and
- (ii) $f \in \mathcal{M}$, and
- (iii) $\rho = \sup(f)$.

By definition, $\text{dom}(f) \leq \max(\text{code}(\alpha))$. Thus $\text{cof}^{\mathcal{M}}(\rho) \leq \max(s) < \kappa_0$. So $\text{cof}^{\mathcal{M}_\nu}(\rho') < \kappa_0 < \kappa$. ◇

This completes the proof of the lemma. □

4. A CORRECTNESS THEOREM

In this section we prove that if a countable, iterable premouse $\mathcal{M}$ is (α, n) -big, then $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$. This gives us one half of the proof that $A_{(\alpha,n)}$ is a mouse set. We work in the theory ZFC.

We shall use heavily a result due to H. Woodin concerning genericity over $L[\vec{E}]$ models. This result is often referred to as “iterating to make a real generic.” We state the result here in the form in which we shall need it. The following is also Theorem 4.3 from [St2].

Lemma 4.1 (Woodin). *Let $\mathcal{M}$ be a countable, realizable, premouse. Suppose $\kappa < \delta < \text{OR}^{\mathcal{M}}$. Suppose $\mathcal{M} \models \text{“}\delta \text{ is a Woodin cardinal.”}$ Let $\mathbb{P} \subseteq J_{\kappa}^{\mathcal{M}}$ be a partial order in $\mathcal{M}$ and let G be $\mathcal{M}$ -generic over $\mathbb{P}$. Then there is a partial order $\mathbb{Q} \subseteq J_{\delta}^{\mathcal{M}}$, with $\mathbb{Q} \in \mathcal{M}$, such that for any real w , there is a 0-maximal iteration tree $\mathcal{T}$ on $\mathcal{M}$ of countable length $\theta + 1$ such that*

- (a) $\mathcal{M}_{\theta}^{\mathcal{T}}$ is realizable, and
- (b) $D^{\mathcal{T}} = \emptyset$ so that $i_{0,\theta}^{\mathcal{T}}$ is defined, and
- (c) $\text{lh}(E_{\xi}^{\mathcal{T}}) < i_{0,\xi}^{\mathcal{T}}(\delta)$ for all $\xi < \theta$, and
- (d) $\text{crit}(E_{\xi}^{\mathcal{T}}) > \kappa$ for all $\xi < \theta$ (so G is $\mathcal{M}_{\theta}^{\mathcal{T}}$ -generic over $\mathbb{P}$), and
- (e) w is $\mathcal{M}_{\theta}^{\mathcal{T}}[G]$ -generic over $i_{0,\theta}^{\mathcal{T}}(\mathbb{Q})$.

For an idea of how to prove this lemma, see the exercises at the end of chapters 6 and 7 in [Ma2]. We shall also need a slightly different form of the above lemma.

Lemma 4.2. *Let $\mathcal{M}$ be a countable, realizable premouse. Suppose $\kappa < \delta < \gamma < \text{OR}^{\mathcal{M}}$, and suppose that δ is a cardinal of $\mathcal{M}$, and $J_{\gamma}^{\mathcal{M}} \models \text{“}\delta \text{ is a Woodin cardinal.”}$ Let $\mathbb{P} \subseteq J_{\kappa}^{\mathcal{M}}$ be a partial order in $\mathcal{M}$ and let G be $\mathcal{M}$ -generic over $\mathbb{P}$. Then there is a partial order $\mathbb{Q} \subseteq J_{\delta}^{\mathcal{M}}$, with $\mathbb{Q} \in J_{\gamma}^{\mathcal{M}}$, such that for any real w , there is a 0-maximal iteration tree $\mathcal{T}$ on $\mathcal{M}$ of countable length $\theta + 1$ such that*

- (a) $\mathcal{M}_{\theta}^{\mathcal{T}}$ is realizable, and
- (b) $D^{\mathcal{T}} = \emptyset$ so that $i_{0,\theta}^{\mathcal{T}}$ is defined, and
- (c) $\text{lh}(E_{\xi}^{\mathcal{T}}) < i_{0,\xi}^{\mathcal{T}}(\delta)$ for all $\xi < \theta$, and
- (d) $\text{crit}(E_{\xi}^{\mathcal{T}}) > \kappa$ for all $\xi < \theta$ (so G is $\mathcal{M}_{\theta}^{\mathcal{T}}$ -generic over $\mathbb{P}$), and
- (e) w is $i_{0,\theta}^{\mathcal{T}}(J_{\gamma}^{\mathcal{M}})[G]$ -generic over $i_{0,\theta}^{\mathcal{T}}(\mathbb{Q})$.

Remark 4.3. Lemma 4.2 does not seem to follow from Lemma 4.1 applied to the premouse $J_{\gamma}^{\mathcal{M}}$. Since δ is not necessarily a regular cardinal of $\mathcal{M}$, if w is a real, and if $\mathcal{T}'$ is the iteration tree on $J_{\gamma}^{\mathcal{M}}$ given by Lemma 4.1, and if $\mathcal{T}$ is the tree on $\mathcal{M}$ induced by $\mathcal{T}'$, and if $\text{lh}(\mathcal{T}) = \theta + 1$, then it is possible that $i_{0,\theta}^{\mathcal{T}'}(\delta) < i_{0,\theta}^{\mathcal{T}}(\delta)$. This means that w may not be $i_{0,\theta}^{\mathcal{T}}(J_{\gamma}^{\mathcal{M}})[G]$ -generic.

However Lemma 4.2 does follow from the proof of Lemma 4.1. The idea is that in the situation described in the above paragraph, we would simply continue to iterate. We sketch this idea below.

Proof of Lemma 4.2. We assume that the reader is familiar with the proof of Lemma 4.1. Let $\mathbb{Q}$ be the partial order given by Lemma 4.1 applied to the premouse $J_\gamma^{\mathcal{M}}$. Fix any real w . The proof of Lemma 4.1 proceeds by building an iteration tree $\mathcal{T}'$ on $J_\gamma^{\mathcal{M}}$ such that for all $\xi < \text{lh } \mathcal{T}'$, $E_\xi^{\mathcal{T}'}$ is the “least witness of the non-genericity of w .” (That is $E_\xi^{\mathcal{T}'}$ is the shortest extender E from the $\mathcal{M}_\xi^{\mathcal{T}'}$ sequence such that there is a set $X \subseteq \text{crit } E$ such that w is not consistent with X but w is consistent with $i_E(X)$.) To prove Lemma 4.2, we can build a tree $\mathcal{T}$ on $\mathcal{M}$ such that for all $\xi < \text{lh } \mathcal{T}$, $E_\xi^{\mathcal{T}}$ is the least witness that w is not generic over $i_{0,\xi}^{\mathcal{T}}(J_\gamma^{\mathcal{M}})[G]$. (Note that since δ is a cardinal of $\mathcal{M}$, any extender in $J_\delta^{\mathcal{M}}$ which is total on $J_\gamma^{\mathcal{M}}$ is also total on $\mathcal{M}$.) The proof that this process must terminate goes through in our setting even though δ is not Woodin in $\mathcal{M}$. $\square$

We are now ready to state the main theorem of this section. This theorem says that our Criterion 1 from the previous section is satisfied. First we give a definition which will help us to state this criterion.

Definition 4.4. Let $\alpha \geq 2$ be projective-like. Let $n \in \omega$. Let $X \subset \mathbb{R}$. Then we say that X is *closed under* $A_{(\alpha,n)}$ iff for all $y \in X$, $A_{(\alpha,n)}(y) \subseteq X$.

Using this definition we can state Criterion 1 from the previous section as follows: If a countable, iterable premouse $\mathcal{M}$ is (α, n) -big above β , and if $\mathbb{P} \in \mathcal{M}$ is a partial order with $(\text{card}(\mathbb{P})^+)^{\mathcal{M}} \leq \beta$, and if G is $\mathcal{M}$ -generic over $\mathbb{P}$, then $\mathbb{R} \cap \mathcal{M}[G]$ should be closed under $A_{(\alpha,n)}$. The motivation for this is that the criterion is met in the case $\alpha = 1$. This was proved by Steel and Woodin in [St5]. Our proof below borrows many ideas from the proof in [St5].

In theorem 4.6 below, we will actually prove something which is ostensibly stronger than the statement of Criterion 1 in the previous paragraph. Recall that on page 21 we discussed Countable Sets of Ordinal Definable Reals, in $L(\mathbb{R})$. In that discussion we defined the sets $\text{OD}_n^{J_\alpha(\mathbb{R})}$ for $n \geq 1$, and we pointed out that for $n \geq 0$, $A_{(\alpha,n)} \subseteq \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. We also asked the question: Is it true that $A_{(\alpha,n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$? As we mentioned before,

we are not able to answer this question using techniques from descriptive set theory, but we are able to shed some light on the question using inner model theory. The way we will do this is by replacing $A_{(\alpha,n)}$ with $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$ in Criterion 1, from the previous paragraph. We start with the obvious analogue of Definition 4.4 above.

Definition 4.5. Let $\alpha \geq 2$ be projective-like. Let $n \in \omega$. Let $X \subset \mathbb{R}$. Then we say that X is *closed under* $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$ iff for all $y \in X$, $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}(y) \subseteq X$.

Now we state the main theorem of this section.

Theorem 4.6. Assume $\text{Det}(\sum_1(J_{\omega_1^{\omega_1}}(\mathbb{R})))$. Let (α, n) be such that $(1, 0) \leq_{\text{lex}} (\alpha, n) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$. Let $\mathcal{M}$ be a countable, realizable premouse. Let $\mathbb{P}$ be a poset in $\mathcal{M}$ and let $\kappa \geq \omega$ be such that $\mathbb{P} \subseteq J_\kappa^{\mathcal{M}}$. Suppose $\mathcal{M} \models \kappa^+$ exists, and let $\beta = (\kappa^+)^{\mathcal{M}}$. Suppose that $\mathcal{M}$ is (α, n) -big above β . Let G be $\mathcal{M}$ -generic over $\mathbb{P}$. Then $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$.

Note. The statement of the theorem includes a determinacy hypothesis. We have not assumed the optimal determinacy hypothesis, but rather an easily statable one. It is possible that the optimal determinacy hypothesis actually follows from the other assumptions of the theorem, and therefore could be omitted. We do not know if this is true.

Before proving Theorem 4.6, we will give a corollary to the theorem. The following is the only result from this section which we will actually use in the rest of the paper.

Corollary 4.7. Assume $\text{Det}(\sum_1(J_{\omega_1^{\omega_1}}(\mathbb{R})))$. Let (α, n) be such that $(1, 0) \leq_{\text{lex}} (\alpha, n) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$. Let $\mathcal{M}$ be a countable, realizable premouse. Suppose that $\mathcal{M}$ is (α, n) -big (above 0.) Then $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$.

Proof. Let (α', n') be the lexicographically least pair such that $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$, and $\mathcal{M}$ is *explicitly* (α', n') -big. Let $\mathcal{N} \trianglelefteq \mathcal{M}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ which is explicitly (α', n') -big. The case $n' = 0$ and $\text{cof}(\alpha') \leq \omega$ is a bit awkward, so let us consider the other cases first. In all other cases, $\mathcal{N} \models \omega_1$ exists, and $\mathcal{N}$ is explicitly (α', n') -big above $\omega_1^{\mathcal{N}}$. Then we may apply Theorem 4.6 to $\mathcal{N}$, with $\mathbb{P} = \emptyset$ and $\beta = \omega_1^{\mathcal{N}}$, to conclude that $\mathbb{R} \cap \mathcal{N}$ is closed under $\text{OD}_{n'+1}^{J_{\alpha'}(\mathbb{R})}$. In particular, $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$.

Now suppose that $\text{cof}(\alpha') \leq \omega$ and $n' = 0$. In this case it may not be true the $\mathcal{N} \models \text{"}\omega_1 \text{ exists."}$ Nevertheless, we will show that $\text{OD}_1^{J_{\alpha'}(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{N}$. By proposition 2.22 we have that $\text{OD}_1^{J_{\alpha'}(\mathbb{R})} = A_{(\alpha', 0)} =$

$$\left\{ x \in \mathbb{R} \mid (\exists \gamma < \alpha') x \in \text{OD}^{J_\gamma(\mathbb{R})} \right\}.$$

Fix $\gamma < \alpha'$. We will show that $\text{OD}^{J_\gamma(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{N}$. Fix $n_0 \geq 1$. We will show that $\text{OD}_{n_0+1}^{J_\gamma(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{N}$. By definition of explicitly $(\alpha', 0)$ -big, there is an α_0 such that $\gamma \leq \alpha_0 < \alpha'$, and $\mathcal{N}$ is explicitly (α_0, n_0) -big. Let $\mathcal{P} \trianglelefteq \mathcal{N}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{N}$ such that $\mathcal{P}$ is explicitly (α_0, n_0) -big. Since $n_0 \geq 1$, we have as above that $\mathcal{P} \models \text{"}\omega_1 \text{ exists"}$, and $\mathcal{P}$ is explicitly (α_0, n_0) -big above $\omega_1^{\mathcal{P}}$. Then we may apply Theorem 4.6 to $\mathcal{P}$, with $\mathbb{P} = \emptyset$ and $\beta = \omega_1^{\mathcal{P}}$, to conclude that $\mathbb{R} \cap \mathcal{P}$ is closed under $\text{OD}_{n_0+1}^{J_{\alpha_0}(\mathbb{R})}$. In particular $\text{OD}_{n_0+1}^{J_\gamma(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{N}$. $\square$

We now turn to the proof of Theorem 4.6. We want to prove the theorem by induction on (α, n) . It turns out that the induction is easier to do if we add a few extra assumptions. What we will do is the following. First we will state a lemma which is almost identical to the theorem, but with a few extra assumptions. Then we will derive the theorem from the lemma. Then we will prove the lemma by induction on (α, n) .

The following lemma is almost identical to Theorem 4.6, with a few exceptions. Notice, in particular, that below we say “explicitly (α, n) -big” and in the theorem above we say “ (α, n) -big.” Also, in the lemma below we put a requirement on $\max(\text{code}(\alpha))$.

Lemma 4.8. *Let (α, n) be such that $(1, 0) \leq_{\text{lex}} (\alpha, n) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $\mathcal{M}$ be a countable, realizable premouse. Let $\mathbb{P}$ be a poset in $\mathcal{M}$ and let κ be such that $\mathbb{P} \subseteq J_\kappa^{\mathcal{M}}$. Suppose $\mathcal{M} \models \kappa^+$ exists, and let $\beta \geq (\kappa^+)^{\mathcal{M}}$. Suppose that $\mathcal{M}$ is explicitly (α, n) -big above β , and that, if $\alpha < \omega_1^{\omega_1}$, $\max(\text{code}(\alpha)) < \beta$. Let G be $\mathcal{M}$ -generic over $\mathbb{P}$. Then $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$.*

First we will derive the theorem from the lemma.

Proof of Theorem 4.6. Fix (α, n) , $\mathcal{M}$, $\mathbb{P}$, κ , β , and G as in the statement of the theorem. Let (α', n') be the lexicographically greatest pair such that $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$ and $\mathcal{M}$ is explicitly (α', n') -big above β . It is easy to see that there is such a pair. (Let X be the

set of pairs (α', n') such that $\mathcal{M}$ is explicitly (α', n') -big above β . Notice that X is countable. Suppose towards a contradiction that $(X, \leq_{\text{lex}})$ has limit order type. Let $(\alpha_0, 0)$ be the supremum of X . Then by lemma 3.13, for cofinally many $\alpha' < \alpha_0$, and all n' , $\mathcal{M}$ is explicitly (α', n') -big above β . If $\mathcal{M}$ had ω Woodin cardinals above β , then $\mathcal{M}$ would be explicitly $(\omega_1^{\omega_1}, 0)$ -big above β , and so $(\omega_1^{\omega_1}, 0)$ would be the greatest pair in X . So suppose that $\mathcal{M}$ does not have ω Woodin cardinals. Then for cofinally many $\alpha' < \alpha_0$, and all n' , some *proper* initial segment of $\mathcal{M}$ is explicitly (α', n') -big above β . But then, by definition, $\mathcal{M}$ is explicitly $(\alpha_0, 0)$ -big above β . Now let $\mathcal{N} \trianglelefteq \mathcal{M}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ such that $\mathcal{N}$ is explicitly (α', n') -big above β . Notice that $\mathbb{R} \cap \mathcal{N}[G] = \mathbb{R} \cap \mathcal{M}[G]$. So it suffices to show that $\mathbb{R} \cap \mathcal{N}[G]$ is closed under $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. We will in fact show that $\mathbb{R} \cap \mathcal{N}[G]$ is closed under $\text{OD}_{n'+1}^{J_{\alpha'}(\mathbb{R})}$. If $(\alpha', n') = (\omega_1^{\omega_1}, 0)$, then we may now apply Lemma 4.8 to $\mathcal{N}$, and we are done. So suppose that $\alpha' < \omega_1^{\omega_1}$. Then we have to worry about the hypothesis in Lemma 4.8 about $\max(\text{code}(\alpha'))$.

The case $\text{cof}(\alpha') \leq \omega$ and $n' = 0$ is slightly awkward, so let us handle the other cases first. In all other cases, there is at least one total-on- $\mathcal{N}$ extender from the $\mathcal{N}$ -sequence, with critical point above β . Let μ be the least such critical point. By Lemma 3.27, $\max(\text{code}(\alpha')) < \mu$. Let β' be any ordinal with $\max(\text{code}(\alpha')) < \beta' < \mu$. By Lemma 3.17 on page 43, and the discussion preceding that lemma, we have that $\mathcal{N}$ is explicitly (α', n') -big above β' . So we may apply Lemma 4.8 to $\mathcal{N}$ and β' to conclude that $\mathbb{R} \cap \mathcal{N}[G]$ is closed under $\text{OD}_{n'+1}^{J_{\alpha'}(\mathbb{R})}$.

Finally, let us consider the case $\text{cof}(\alpha') \leq \omega$ and $n' = 0$. By proposition 2.22 we have that $\text{OD}_1^{J_{\alpha'}(\mathbb{R})} = A_{(\alpha', 0)} =$

$$\left\{ x \in \mathbb{R} \mid (\exists \gamma < \alpha') x \in \text{OD}^{J_\gamma(\mathbb{R})} \right\}.$$

We have that for all $\gamma < \alpha'$, and all m , $\mathcal{N}$ is (γ, m) -big above β . By the previous cases of the theorem we are proving, for all $\gamma < \alpha'$, and all $m \neq 0$, $\mathbb{R} \cap \mathcal{N}[G]$ is closed under $\text{OD}_{m+1}^{J_\gamma(\mathbb{R})}$. Thus $\mathbb{R} \cap \mathcal{N}[G]$ is closed under $\text{OD}_1^{J_{\alpha'}(\mathbb{R})}$. $\square$

Now we turn to the proof of Lemma 4.8. We will prove the lemma by induction on (α, n) . There will be several different cases for the inductive step. Some of these cases are direct analogues of cases in the proof by Steel and Woodin in [St5]. Other cases are

unique to our setting. Two of the unique cases are: $\text{cof}(\alpha) \leq \omega$ with $n = 1$, and $n = 2$. In these two cases, our primary tool will be the following lemma. This lemma is the motivation for the definition of explicitly $(\alpha, 1)$ -big, in the case $\text{cof}(\alpha) \leq \omega$. The lemma says the if $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big, then $\mathcal{M}$ can tell whether or not a given $\Sigma_2(J_\alpha(\mathbb{R}))$ fact is true.

Lemma 4.9. *Suppose $2 \leq \alpha < \omega_1^{\omega_1}$, and that $\text{cof}(\alpha) \leq \omega$. Assume $\text{Det}(J_\alpha(\mathbb{R}))$. Let $\mathcal{M}$ be a countable, realizable premouse. Let $\mathbb{P}$ be a poset in $\mathcal{M}$ and let κ be such that $\mathbb{P} \subseteq J_\kappa^\mathcal{M}$. Suppose $\mathcal{M} \models \kappa^+$ exists, and let $\beta = (\kappa^+)^{\mathcal{M}}$. Suppose that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β and no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . Let φ be a Σ_2 formula in the language of $J_\alpha(\mathbb{R})$, with one free variable. Then there is a Π_1 formula ψ in the language of premice, and a parameter $a \in \mathcal{M}$, such that whenever G is $\mathcal{M}$ -generic over $\mathbb{P}$ and $x \in \mathbb{R} \cap \mathcal{M}[G]$ we have that:*

$$J_\alpha(\mathbb{R}) \models \varphi[x]$$

iff

$$\mathcal{M}[G] \models \psi[x, a, G].$$

Lemma 4.8 and Lemma 4.9 will be proved by simultaneous induction. For ease of readability we will separate these into two proofs. We begin with the proof of Lemma 4.9.

Proof of Lemma 4.9. We assume that Lemma 4.8 holds for all (α', n) with $1 \leq \alpha' < \alpha$.

Since no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β , there is an ordinal $\rho \in \mathcal{M}$ such $\beta < \rho$ and $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . Recall that this implies that ρ is a limit cardinal of $\mathcal{M}$, and that $\mathcal{M}$ is the first admissible structure over $\mathcal{J}_\rho^\mathcal{M}$. Also, by Lemma 3.30, there is a function f such that:

- (a) f is an α -ladder above β relative to $\mathcal{M}$, and
- (b) $f \in \mathcal{M}$, and
- (c) $\rho = \sup(f)$.

Recall what it means that f is an α -ladder above β relative to $\mathcal{M}$. There are two cases. First suppose that $\alpha = \alpha' + 1$ is a successor ordinal. (See Figure 1 on page 39.) Then:

- (i) $\text{dom}(f) = \omega$, and
- (ii) for all n , $f(n) = (\delta_n, \gamma_n)$, with $\beta < \delta_n < \gamma_n < \delta_{n+1} < \text{OR}^{\mathcal{M}}$, and
- (iii) δ_n is a cardinal of $\mathcal{M}$, and $J_{\gamma_n}^{\mathcal{M}} \models$ “ δ_n is a Woodin cardinal,” and γ_n is least such that $J_{\gamma_n}^{\mathcal{M}}$ is explicitly $(\alpha', 2n)$ -big above δ_n , and
- (iv) $\max(\text{code}(\alpha)) < \delta_0$.

Next suppose that α is a limit ordinal of cofinality ω . Then:

- (i) $\text{dom}(f)$ is a countable limit ordinal $\mu \leq \max(\text{code}(\alpha))$, and
- (ii) for all $\xi \in \text{dom}(f)$ we have that: $f(\xi) = (\delta_\xi, \gamma_\xi, s_\xi)$ with $\beta < \delta_\xi < \gamma_\xi < \text{OR}^{\mathcal{M}}$, and $\delta_\xi > \sup \{ \gamma_\eta \mid \eta < \xi \}$, and $s_\xi \in \mathfrak{C}$ is a code for an ordinal less than ω_1^ω , and, letting $\alpha_\xi = |s_\xi|$, we have that $\alpha_\xi < \alpha$ and
- (iii) δ_ξ is a cardinal of $\mathcal{M}$, and $J_{\gamma_\xi}^{\mathcal{M}} \models$ “ δ_ξ is a Woodin cardinal,” and γ_ξ is least such that $J_{\gamma_\xi}^{\mathcal{M}}$ is explicitly $(\alpha_\xi, 2)$ -big above δ_ξ , and
- (iv) for all ξ , $\max(s_\xi) \leq \max(\text{code}(\alpha)) < \delta_0$, and
- (v) $\{ \alpha_\xi \mid \xi \in \text{dom}(f) \}$ is cofinal in α .

Also recall that $\sup(f) = \sup \{ \gamma_\xi \mid \xi \in \text{dom}(f) \} = \sup \{ \delta_\xi \mid \xi \in \text{dom}(f) \}$.

For notational simplicity we make the following definitions. Let $\mu = \text{dom}(f)$. So $\mu = \omega$ if α is a successor ordinal, and if α is a limit ordinal of cofinality ω , then $\mu \leq \max(\text{code}(\alpha))$ is some countable limit ordinal. If α is a limit ordinal, then for $\xi < \mu$ the codes s_ξ were defined above, and we set $\alpha_\xi = |s_\xi|$. If α is a successor ordinal, then for $\xi < \mu$ let $\alpha_\xi = \alpha - 1$ and let $s_\xi = \text{code}(\alpha - 1)$. Also, for $\xi < \mu$, let

$$n_\xi = \begin{cases} 2\xi & \text{if } \alpha \text{ is a successor ordinal,} \\ 2 & \text{if } \alpha \text{ is a limit ordinal.} \end{cases}$$

Notice then that for all $\xi < \mu$, $J_{\gamma_\xi}^{\mathcal{M}} \models \delta_\xi^+$ exists, and $J_{\gamma_\xi}^{\mathcal{M}}$ is explicitly (α_ξ, n_ξ) -big above $(\delta_\xi^+)^{J_{\gamma_\xi}^{\mathcal{M}}}$. This is important because we want to apply Lemma 4.8 for (α_ξ, n_ξ) to the premouse $J_{\gamma_\xi}^{\mathcal{M}}$.

Now, fix a Σ_2 formula φ , as in the statement of the Lemma. Recall that there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ partial function from $\mathbb{R}$ onto $J_\alpha(\mathbb{R})$. As a consequence of this,

$$\Sigma_2(J_\alpha(\mathbb{R})) = \exists^{\mathbb{R}}[\Sigma_1(J_\alpha(\mathbb{R})) \wedge \Pi_1(J_\alpha(\mathbb{R}))].$$

So there are two Σ_1 formulas φ_1 and φ_2 so that for all $x \in \mathbb{R}$

$$J_\alpha(\mathbb{R}) \models \varphi[x]$$

iff

$$(\exists y \in \mathbb{R}) J_\alpha(\mathbb{R}) \models (\varphi_1 \wedge \neg \varphi_2)[x, y].$$

Fix such formulas φ_1 and φ_2 .

We now fix two sequences of formulas $\langle \theta_\xi^1 \mid \xi < \mu \rangle$ and $\langle \theta_\xi^2 \mid \xi < \mu \rangle$ such that $(\forall x, y \in \mathbb{R})$

$$J_\alpha(\mathbb{R}) \models \varphi_1[x, y]$$

iff

$$(\exists \xi < \mu) \quad J_{\alpha_\xi}(\mathbb{R}) \models \theta_\xi^1[x, y]$$

and

$$J_\alpha(\mathbb{R}) \models \varphi_2[x, y]$$

iff

$$(\exists \xi < \mu) \quad J_{\alpha_\xi}(\mathbb{R}) \models \theta_\xi^2[x, y].$$

If α is a limit ordinal, then we simply set $\theta_\xi^1 = \varphi_1$ and $\theta_\xi^2 = \varphi_2$ for all $\xi < \alpha$. If α is a successor ordinal, then by Lemma 1.1, we may take $\langle \theta_\xi^1 \mid \xi \in \omega \rangle$ and $\langle \theta_\xi^2 \mid \xi \in \omega \rangle$ to be recursive. Notice that in either case we have that $\langle \theta_\xi^1 \mid \xi < \mu \rangle$ and $\langle \theta_\xi^2 \mid \xi < \mu \rangle$ are in $J_\rho^{\mathcal{M}}$. Also, we may assume that for $\xi < \mu$, θ_ξ^1 and θ_ξ^2 are both (logically equivalent to) Σ_{n_ξ} formulas.

Suppose for a moment that G is $\mathcal{M}$ -generic over $\mathbb{P}$, and that for each $\xi \in \text{dom}(f)$, H_ξ is $\mathcal{J}_{\gamma_\xi}^{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \delta_\xi)$, where $\text{Coll}(\nu_1, \nu_2)$ is the partial order for collapsing ν_2 to ν_1 . In this situation we will want to show later that $\mathcal{J}_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi]$ has enough “correctness” that it can tell when θ_ξ^1 and θ_ξ^2 are true in the model $J_{\alpha_\xi}(\mathbb{R})$. To do this, we must consider the following technical point. For $\xi < \mu$, let $\bar{\mathbb{R}}_\xi = \mathbb{R} \cap J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi]$. Notice that $\omega_1^{L(\bar{\mathbb{R}}_\xi)} = (\delta_\xi^+)^{J_{\gamma_\xi}^{\mathcal{M}}}$. Recall that $s_\xi = \text{code}(\alpha_\xi)$ and that $\max(s_\xi) < \delta_0 \leq \delta_\xi$. Thus

$\max(s_\xi) < \omega_1^{L(\bar{\mathbb{R}}_\xi)}$. Thus it makes sense to consider the term $|s_\xi|^{L(\bar{\mathbb{R}}_\xi)}$. This will be important below. (cf. the discussion preceding Lemma 2.36 on page 30.)

For any $\mathcal{M}$, $\mathbb{P}$ as in the statement of the lemma, and any ladder f as above, and any G which is $\mathcal{M}$ generic over $\mathbb{P}$, let $S(\mathcal{M}, f, \mathbb{P}, G)$ be the collection of all $(x, y) \in \mathbb{R}^2$ such that:

There exists (in V) some sequence $\langle H_\xi \mid \xi \in \text{dom}(f) \rangle$ s.t. $(\forall \xi \in \text{dom}(f)) \left[H_\xi \right.$
 is $J_{\gamma_\xi}^{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \delta_\xi)$ and $(x, y) \in J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi] \left. \right]$ and, letting $\bar{\mathbb{R}}_\xi =$
 $\mathbb{R} \cap J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi]$, and $\bar{\alpha}_\xi = |s_\xi|^{L(\bar{\mathbb{R}}_\xi)}$ we have that $(\exists \xi \in \text{dom}(f)) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models$
 $\theta_\xi^1[x, y]]$ and $(\forall \xi \in \text{dom}(f)) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \neg \theta_\xi^2[x, y]]$.

Then $S(\mathcal{M}, f, \mathbb{P}, G)$ is Σ_1^1 uniformly in any real coding $(\mathcal{J}_\rho^{\mathcal{M}}, f, \mathbb{P}, G)$, where $\rho = \sup(f)$. That is, there is a recursive tree T on $\omega \times \omega \times \omega$ such that for any $\mathcal{M}, \mathbb{P}$ as in the statement of the lemma, and any ladder f as above, and any G which is $\mathcal{M}$ generic over $\mathbb{P}$, and any real z coding $(\mathcal{J}_\rho^{\mathcal{M}}, f, \mathbb{P}, G)$, where $\rho = \sup(f)$, we have that

$$S(\mathcal{M}, f, \mathbb{P}, G) = [T_z] = \{ (x, y) \mid (\forall i \in \omega) (x \restriction i, y \restriction i, z \restriction i) \in T \}.$$

Fix such a recursive tree T .

Now we return to our fixed $\mathcal{M}$, $\mathbb{P}$, f , and $\rho = \sup(f)$. (We have not yet fixed a G .) We will use the following notation: If τ is a term in a forcing language, and K is a generic filter, then τ_K denotes the K -interpretation of τ . Now, let $\dot{z}$ be a $\mathbb{P}$ -name in $\mathcal{M}$ such that whenever G is $\mathcal{M}$ -generic over $\mathbb{P}$, $\dot{z}_G$ is a $\text{Coll}(\omega, \rho)$ -name in $\mathcal{M}[G]$, and whenever H is $\mathcal{M}[G]$ -generic over $\text{Coll}(\omega, \rho)$, $(\dot{z}_G)_H$ is a real coding $(\mathcal{J}_\rho^{\mathcal{M}}, f, \mathbb{P}, G)$. Let $a = \langle \rho, \dot{z} \rangle$. We claim that this a , in conjunction with an appropriate ψ , satisfies the conclusion of the lemma. Now we define the appropriate formula ψ .

For any reals x, z let $T_{x,z}$ be the tree on ω given by:

$$s \in T_{x,z} \Leftrightarrow (x \restriction \text{lh } s, s, z \restriction \text{lh } s) \in T.$$

Claim. *There is a Π_1 formula ψ such that whenever G is $\mathcal{M}$ generic over $\mathbb{P}$, and $x \in \mathbb{R} \cap \mathcal{M}[G]$, we have that:*

$$\mathcal{M}[G] \models \psi[x, a, G]$$

iff

whenever H is $\mathcal{M}[G]$ -generic over $\text{Coll}(\omega, \rho)$, $T_{x, (\dot{z}_G)_H}$ is ill-founded.

Proof of Claim. We can take ψ to say:

$$\emptyset \Vdash \frac{\mathcal{M}[G]}{\text{Coll}(\omega, \rho)} \neg[\exists \text{ a rank function for } T_{\dot{x}, \dot{z}_G}].$$

More precisely, we can take ψ to say: “For all forcing conditions q , and all names σ_1 and σ_2 , if $\sigma_2 = \dot{z}_G$, then $\neg[q \Vdash \sigma_1 \text{ is a rank function for } T_{\dot{x}, \sigma_2}]$.” We leave it to the reader to see that this can be expressed over $\mathcal{M}[G]$ as a Π_1 statement about x , a , and G .

Now, if $T_{x, (\dot{z}_G)_H}$ is ill-founded whenever H is $\mathcal{M}[G]$ -generic over $\text{Coll}(\omega, \rho)$, then clearly $\mathcal{M}[G] \models \psi[x, a, G]$. The converse is true because for all such H , $\mathcal{M}[G][H]$ is a model of KP. $\diamond$

Fix ψ as in the proof of the claim. We claim that this ψ satisfies the conclusion of the lemma. Let G be $\mathcal{M}$ -generic over $\mathbb{P}$, and let $x \in \mathbb{R} \cap \mathcal{M}[G]$. We will show that

$$J_\alpha(\mathbb{R}) \models \varphi[x]$$

iff

$$\mathcal{M}[G] \models \psi[x, a, G].$$

Suppose first that $\mathcal{M}[G] \models \psi[x, a, G]$. Let H be $\mathcal{M}[G]$ -generic over $\text{Coll}(\omega, \rho)$. Let $z = (\dot{z}_G)_H$. Then $\exists y \in \mathbb{R}$ such that $(x, y, z) \in [T]$. Fix such a y . Then $(x, y) \in S(\mathcal{M}, f, \mathbb{P}, G)$. So let $\langle H_\xi \mid \xi \in \text{dom } f \rangle$ be such that:

$$\begin{aligned} & (\forall \xi \in \text{dom}(f)) \left[H_\xi \text{ is } J_{\gamma_\xi}^{\mathcal{M}}[G]\text{-generic over } \text{Coll}(\omega, \delta_\xi) \text{ and } (x, y) \in J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi] \right] \\ & \text{and, letting } \bar{\mathbb{R}}_\xi = \mathbb{R} \cap J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi], \text{ and } \bar{\alpha}_\xi = |s_\xi|^{L(\bar{\mathbb{R}}_\xi)} \text{ we have that } (\exists \xi \in \\ & \text{dom}(f)) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \theta_\xi^1[x, y]] \text{ and } (\forall \xi \in \text{dom}(f)) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \neg \theta_\xi^2[x, y]]. \end{aligned}$$

For a moment, fix $\xi \in \text{dom } f$. Let $\bar{\mathbb{R}}_\xi = \mathbb{R} \cap J_{\gamma_\xi}^{\mathcal{M}}[G][H_\xi]$. Let $\bar{\alpha}_\xi = |s_\xi|^{L(\bar{\mathbb{R}}_\xi)}$. We will show that $J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi)$ is Σ_{n_ξ} elementarily embeddable into $J_{\alpha_\xi}(\mathbb{R})$. There is a K which is $J_{\gamma_\xi}^{\mathcal{M}}$ -generic over $\text{Coll}(\omega, \delta_\xi)$ such that $\bar{\mathbb{R}}_\xi = \mathbb{R} \cap J_{\gamma_\xi}^{\mathcal{M}}[K]$. As $J_{\gamma_\xi}^{\mathcal{M}}$ is (α_ξ, n_ξ) -big above $(\delta_\xi^+)^{J_{\gamma_\xi}^{\mathcal{M}}}$, Lemma 4.8 for (α_ξ, n_ξ) implies that $\bar{\mathbb{R}}_\xi$ is closed under $\text{OD}_{n_\xi+1}^{J_{\alpha_\xi}(\mathbb{R})}$. In particular $\bar{\mathbb{R}}_\xi$ is closed downward under $\Delta_{(\alpha_\xi, n_\xi)}$. As $J_{\gamma_\xi}^{\mathcal{M}}$ has a measurable cardinal above δ_ξ , $\bar{\mathbb{R}}_\xi = \mathbb{R} \cap L(\bar{\mathbb{R}}_\xi)$. Since n_ξ is even, we can apply Lemma 2.36 on page 30 to conclude that $J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi)$ is in fact $\Sigma_{n_\xi+1}$ elementarily embeddable into $J_{\alpha_\xi}(\mathbb{R})$. (This is why we organized the definition of an α -ladder so that n_ξ is even.)

Now notice that we have:

$$(\exists \xi \in \text{dom } f) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \theta_\xi^1[x, y]],$$

and

$$(\forall \xi \in \text{dom } f) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \neg \theta_\xi^2[x, y]].$$

Thus we have:

$$(\exists \xi \in \text{dom } f) [J_{\alpha_\xi}(\mathbb{R}) \models \theta_\xi^1[x, y]],$$

and

$$(\forall \xi \in \text{dom } f) [J_{\alpha_\xi}(\mathbb{R}) \models \neg \theta_\xi^2[x, y]].$$

And so we have:

$$J_\alpha(\mathbb{R}) \models (\varphi_1 \wedge \neg \varphi_2)[x, y].$$

And so we have:

$$J_\alpha(\mathbb{R}) \models \varphi[x].$$

Conversely, assume that $J_\alpha(\mathbb{R}) \models \varphi[x]$. We will show that $\mathcal{M}[G] \models \psi[x, a, G]$.

Since $J_\alpha(\mathbb{R}) \models \varphi[x]$, $\exists y \in \mathbb{R}$ such that $J_\alpha(\mathbb{R}) \models (\varphi_1 \wedge \neg \varphi_2)[x, y]$. Fix such a y . We are going to iterate $\mathcal{M}$, yielding an iteration map $i : \mathcal{M} \rightarrow \bar{\mathcal{M}}$, with $\text{crit}(i) > \beta$, so that for all $\xi \in \text{dom}(f)$, $\langle x, y \rangle$ is $i(J_{\gamma_\xi}^{\mathcal{M}})[G]$ -generic over a partial order contained in $i(J_{\delta_\xi}^{\mathcal{M}})$.

We apply Lemma 4.2 iteratively, for each $\xi \in \text{dom } f$. This gives us, for each $\xi \in \text{dom } f$: a premouse $\mathcal{M}_\xi$, and a partial order $\mathbb{Q}_\xi \in \mathcal{M}_\xi$, and an iteration tree $\mathcal{T}_\xi$ on $\mathcal{M}_\xi$ of countable

length $\theta_\xi + 1$; and a system of commuting maps $i_{\xi,\eta} : \mathcal{M}_\xi \rightarrow \mathcal{M}_\eta$ for $\xi < \eta \in \text{dom } f$; such that:

- (a) $\mathcal{M}_0 = \mathcal{M}$, and
- (b) for each $\xi \in \text{dom}(f)$, $\mathcal{M}_{\xi+1}$ is the last model of $\mathcal{T}_\xi$, and
- (c) for each $\xi \in \text{dom}(f)$, $D^{\mathcal{T}_\xi} = \emptyset$ and $i_{\xi,\xi+1} = i_{0,\theta_\xi}^{\mathcal{T}_\xi}$, and for $\xi' < \xi$, $i_{\xi',\xi+1} = i_{\xi',\xi} \circ i_{\xi,\xi+1}$, and
- (d) for limit $\eta \in \text{dom}(f)$, $\mathcal{M}_\eta$ is the direct limit of the $\mathcal{M}_\xi$ under the $i_{\xi,\xi'}$ for $\xi < \xi' < \eta$, and $i_{\xi,\eta}$ is the direct limit map, and
- (e) for each $\xi \in \text{dom}(f)$, $\mathcal{M}_\xi$ is realizable, and
- (f) for each $\xi \in \text{dom}(f)$, $\text{lh}(E_\nu^{\mathcal{T}_\xi}) < i_{0,\nu}^{\mathcal{T}_\xi} \circ i_{0,\xi}(\delta_\xi)$ for all $\nu < \theta_\xi$, and
- (g) for each $\xi \in \text{dom}(f)$, $\text{crit}(E_\nu^{\mathcal{T}_\xi}) > \sup \{ i_{0,\xi}(\gamma_{\xi'}) \mid \xi' < \xi \}$ for all $\nu < \theta_\xi$ (in particular $\text{crit}(i_{0,\xi+1}) > \beta$ so that G is $\mathcal{M}_{\xi+1}$ -generic over $\mathbb{P}$), and
- (h) for each $\xi \in \text{dom } f$, $\mathbb{Q}_\xi \subseteq i_{0,\xi}(J_{\delta_\xi}^{\mathcal{M}})$ and $\mathbb{Q}_\xi \in i_{0,\xi}(J_{\gamma_\xi}^{\mathcal{M}})$ and $\langle x, y \rangle$ is $i_{0,\xi+1}(J_{\gamma_\xi}^{\mathcal{M}})[G]$ -generic over $i_{\xi,\xi+1}(\mathbb{Q}_\xi)$.

The partial order $\mathbb{Q}_\xi$ and the tree $\mathcal{T}_\xi$ comes from applying Lemma 4.2 with

- (i) $\mathcal{M} = \mathcal{M}_\xi$
- (ii) $\kappa = \sup \{ i_{0,\xi}(\gamma_{\xi'}) \mid \xi' < \xi \}$, (In the case $\xi = 0$ take $\kappa = \max(\text{code}(\alpha))$.)
- (iii) $\delta = i_{0,\xi}(\delta_\xi)$,
- (iv) $\gamma = i_{0,\xi}(\gamma_\xi)$,
- (v) $w = \langle x, y \rangle$,
- (vi) $\mathbb{P} = \mathbb{P}$, and $G = G$.

The reason that we can have (ii) above is that by the definition of an α -ladder $\delta_\xi > \sup \{ \gamma_{\xi'} \mid \xi' < \xi \}$, and so $i_{0,\xi}(\delta_\xi) > \sup \{ i_{0,\xi}(\gamma_{\xi'}) \mid \xi' < \xi \}$.

The above remarks clearly define by recursion a sequence

$$\langle \mathcal{M}_\xi, \mathcal{T}_\xi, \mathbb{Q}_\xi \mid \xi \in \text{dom } f \rangle$$

and a system of maps

$$\{ i_{\xi,\eta} \mid \xi < \eta < \text{dom}(f) \}$$

such that properties (a) through (h) above hold. The reason that we are able to maintain that (e) holds at limit ξ is that for all but finitely many $\eta_1 < \eta_2 < \xi$, the realization

maps for $\mathcal{M}_{\eta_1}$ and $\mathcal{M}_{\eta_2}$ commute with each other. (cf. the proof in [St2] that if $\mathcal{M}$ is a countable realizable mouse, then player II wins the weak iteration game on $\mathcal{M}$ of length ω_1 .)

Let $\bar{\mathcal{M}}$ be the direct limit of the $\mathcal{M}_\xi$ under the $i_{\xi,\eta}$, and let $i_\xi : \mathcal{M}_\xi \rightarrow \bar{\mathcal{M}}$ be the direct limit map. Let $i = i_0$. As in the previous paragraph, $\bar{\mathcal{M}}$ is realizable. By (g) we have that $\text{crit}(i) > \beta$ so that G is $\bar{\mathcal{M}}$ -generic over $\mathbb{P}$. And by (g) and (h) we have that for all $\xi \in \text{dom}(f)$, $\langle x, y \rangle$ is $i(J_{\gamma_\xi}^{\mathcal{M}})[G]$ -generic over $i_\xi(\mathbb{Q}_\xi)$.

To complete the proof of Lemma 4.9, we must show that $\mathcal{M}[G] \models \psi[x, a, G]$. Let $\dot{x}$ and $\dot{G}$ be $\mathbb{P}$ -names in $J_\beta^{\mathcal{M}}$ such that $\dot{x}_G = x$ and $\dot{G}_G = G$. Since ψ is a Π_1 formula, the relation $R(p, \sigma_1, \sigma_2, \sigma_3) \Leftrightarrow p \Vdash \psi(\sigma_1, \sigma_2, \sigma_3)$ is Π_1 definable over $\mathcal{M}$. Then since $\text{crit}(i) > \beta$, we have

$$\begin{aligned}
& \mathcal{M}[G] \models \psi[x, a, G] \\
& \text{iff} \\
& (\exists p \in G) \mathcal{M} \models "p \Vdash \psi(\dot{x}, \check{a}, \dot{G})" \\
& \text{iff} \\
& (\exists p \in G) \bar{\mathcal{M}} \models "p \Vdash \psi(\dot{x}, i(\check{a}), \dot{G})" \\
& \text{iff} \\
& \bar{\mathcal{M}}[G] \models \psi[x, i(a), G]
\end{aligned}$$

So it suffices to show that $\bar{\mathcal{M}}[G] \models \psi[x, i(a), G]$. Now, $i(a) = \langle i(\rho), i(\dot{z}) \rangle$. Let H be $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, i(\rho))$. Let $z = (i(\dot{z})_G)_H$. It suffices to show that $\mathcal{M}[G][H] \models$ "There is no rank function for $T_{x,z}$." For this it suffices to show that $(x, y, z) \in [T]$.

Since $\max(\text{code}(\alpha)) < \text{crit}(i)$, $i(f)$ is an α -ladder above β relative to $\bar{\mathcal{M}}$. Thus $S(\bar{\mathcal{M}}, i(f), \mathbb{P}, G)$ is defined. It is not difficult to see that z codes $(J_{i(\rho)}^{\bar{\mathcal{M}}}, i(f), \mathbb{P}, G)$. Thus $[T_z] = S(\bar{\mathcal{M}}, i(f), \mathbb{P}, G)$. So to complete the proof it suffices to show that $(x, y) \in S(\bar{\mathcal{M}}, i(f), \mathbb{P}, G)$.

Since $\text{dom}(f) \leq \max(\text{code}(\alpha)) < \text{crit}(i)$, $\text{dom}(i(f)) = \text{dom}(f)$. For $\xi \in \text{dom}(f)$, let $(\bar{\delta}_\xi, \bar{\gamma}_\xi) = i(\delta_\xi, \gamma_\xi)$. For each $\xi \in \text{dom } f$, $\langle x, y \rangle$ is $J_{\bar{\gamma}_\xi}^{\bar{\mathcal{M}}}[G]$ -generic over $i_\xi(\mathbb{Q}_\xi)$, and $i_\xi(\mathbb{Q}_\xi) \subseteq J_{\bar{\delta}_\xi}^{\bar{\mathcal{M}}}$. Let H_ξ be $J_{\bar{\gamma}_\xi}^{\bar{\mathcal{M}}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta}_\xi)$ with $\langle x, y \rangle \in J_{\bar{\gamma}_\xi}^{\bar{\mathcal{M}}}[G][H_\xi]$.

For a moment, fix $\xi \in \text{dom } f$. Let $\bar{\mathbb{R}}_\xi = \mathbb{R} \cap J_{\bar{\gamma}_\xi}^{\bar{\mathcal{M}}}[G][H_\xi]$, and let $\bar{\alpha}_\xi = |s_\xi|^{L(\bar{\mathbb{R}}_\xi)}$. Exactly as we did before, we may show that $J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi)$ is Σ_{n_ξ} elementarily embeddable into $J_{\alpha_\xi}(\mathbb{R})$.

Now we have:

$$J_\alpha(\mathbb{R}) \models (\varphi_1 \wedge \neg \varphi_2)[x, y].$$

So we have:

$$(\exists \xi \in \text{dom } f) [J_{\alpha_\xi}(\mathbb{R}) \models \theta_\xi^1[x, y]],$$

and

$$(\forall \xi \in \text{dom } f) [J_{\alpha_\xi}(\mathbb{R}) \models \neg \theta_\xi^2[x, y]].$$

And so we have:

$$(\exists \xi \in \text{dom } f) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \theta_\xi^1[x, y]],$$

and

$$(\forall \xi \in \text{dom } f) [J_{\bar{\alpha}_\xi}(\bar{\mathbb{R}}_\xi) \models \neg \theta_\xi^2[x, y]].$$

Thus the sequence $\langle H_\xi \mid \xi \in \text{dom } f \rangle$ witnesses that $(x, y) \in S(\bar{\mathcal{M}}, i(f), \mathbb{P}, G)$. $\square$

We now turn to the proof of Lemma 4.8. The statement of the lemma is on page 63.

Proof of Lemma 4.8. The lemma is proved by induction on (α, n) . For $\alpha = 1$ the lemma is essentially proved in Section 4 of [St2]. So we assume that $\alpha \geq 2$. We also assume that Lemma 4.9 on page 65 holds for α .

Without loss of generality we may assume that there is no proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\beta \in \mathcal{N}$ and $\mathcal{N}$ is explicitly (α, n) -big above β . (If there were such an $\mathcal{N} \triangleleft \mathcal{M}$ then, since $\beta \geq (\kappa^+)^{\mathcal{M}}$, $\mathbb{R} \cap \mathcal{N}[G] = \mathbb{R} \cap \mathcal{M}[G]$, and so we may replace $\mathcal{M}$ with $\mathcal{N}$.)

We split into 6 cases depending on (α, n) .

CASE 1 $\text{cof}(\alpha) \leq \omega$ and $n = 0$. By Proposition 2.22 we have that $\text{OD}_1^{J_\alpha(\mathbb{R})} = A_{(\alpha,0)} =$

$$\left\{ x \in \mathbb{R} \mid (\exists \alpha' < \alpha) x \in \text{OD}^{J_{\alpha'}(\mathbb{R})} \right\}.$$

We are assuming that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . This implies that for cofinally many $\alpha' < \alpha$, and all n , $\mathcal{M}$ is explicitly (α', n) -big above β . By Lemma 2.35 on page 29, for all sufficiently large $\alpha' < \alpha$, $\max(\text{code}(\alpha')) \leq \max(\text{code}(\alpha))$, and so $\max(\text{code}(\alpha')) < \beta$. Thus we may apply our induction hypothesis and conclude that for cofinally many $\alpha' < \alpha$, and all n , $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_{n+1}^{J_{\alpha'}(\mathbb{R})}$. Thus $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_1^{J_\alpha(\mathbb{R})}$.

CASE 2 $\text{cof}(\alpha) > \omega$ and $n = 0$. We are assuming that $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β , and no proper initial segment of $\mathcal{M}$ is. This implies that $\mathcal{M}$ is active. Let κ be the critical point of the last extender on the $\mathcal{M}$ sequence. Then $\kappa > \beta$. Fix any $\alpha' < \alpha$ and any $n \in \omega$. As in the proof of Lemma 3.31, there is a countable premouse $\mathcal{M}'$ such that

- (a) $\mathcal{M}'$ is a linear iterate of $\mathcal{M}$ via its last extender, and
- (b) letting $i : \mathcal{M} \rightarrow \mathcal{M}'$ be the iteration map we have that $i(\kappa) > \max(\text{code}(\alpha'))$, and
- (c) $\mathcal{M}'$ is realizable, and
- (c) $\mathcal{M}'$ is explicitly (α', n) -big above β .

Let β' be any ordinal with $\max(\text{code}(\alpha')) < \beta' < i(\kappa)$. By Lemma 3.17, $\mathcal{M}'$ is explicitly (α', n) -big above β' . Thus we may apply the Lemma we are proving, for (α', n) , to $\mathcal{M}'$ and β' , to conclude that $\mathbb{R} \cap \mathcal{M}'[G]$ is closed under $\text{OD}_{n+1}^{J_{\alpha'}(\mathbb{R})}$.

Notice that for any such $\mathcal{M}'$ as above, $\mathbb{R} \cap \mathcal{M}'[G] = \mathbb{R} \cap \mathcal{M}[G]$. So we have that for all $\alpha' < \alpha$ and all n , $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_{n+1}^{J_{\alpha'}(\mathbb{R})}$. As in the previous case, this implies that $\mathbb{R} \cap \mathcal{M}[G]$ is closed under $\text{OD}_1^{J_\alpha(\mathbb{R})}$.

CASE 3 $\text{cof}(\alpha) \leq \omega$ and $n = 1$. We are assuming that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β and that no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 1)$ big above β . This means that there is an ordinal ρ in $\mathcal{M}$ such that $(\mathcal{M}, \rho)$ witnesses that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . Recall that this means that ρ is a limit cardinal of $\mathcal{M}$ and:

- (i) $\mathcal{M}$ is the first admissible structure over $\mathcal{J}_\rho^\mathcal{M}$, and
- (ii) $\text{code}(\alpha) \in J_\rho^\mathcal{M}$, and
- (iii) for cofinally many $\alpha' < \alpha$, and all $n \in \omega$, and cofinally many $\delta < \rho$ we have that

- (a) δ is a cardinal of $\mathcal{M}$, and
- (b) δ is locally (α', n) Woodin in $\mathcal{J}_\rho^\mathcal{M}$.

Let $x_0 \in \mathbb{R} \cap \mathcal{M}[G]$, and let $y_0 \in \text{OD}_2^{J_\alpha(\mathbb{R})}(x_0)$. We will show that $y_0 \in \mathcal{M}[G]$. Fix $\xi < \omega_1$ such that for all $w \in \text{WO}$ with $|w| = \xi$, $\{y_0\} \in \Sigma_2(J_\alpha(\mathbb{R}), \{x_0, w\})$.

We begin with a claim that says that we may ignore the ordinal parameter ξ .

Claim 1. *Without loss of generality we may assume that $\{y_0\} \in \Sigma_2(J_\alpha(\mathbb{R}), \{x_0\})$.*

Proof of Claim 1. Recall that we are assuming that $\max(\text{code}(\alpha)) < \beta$. Notice that $\mathcal{M}$ has a measurable cardinal μ such $\beta < \mu$, and such that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above μ . Let $j : \mathcal{M} \rightarrow \bar{\mathcal{M}}$ be the iteration map resulting from iterating $\mathcal{M}$, via μ , $\xi + 1$ times. Then $\xi < j(\mu)$. By Lemma 3.34, $\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above $j(u)$. Notice that $\mathbb{R} \cap \mathcal{M}[G] = \mathbb{R} \cap \bar{\mathcal{M}}[G]$. So to see that $y_0 \in \mathcal{M}[G]$, it suffices to see that $y_0 \in \bar{\mathcal{M}}[G]$.

In order to see that $y_0 \in \bar{\mathcal{M}}[G]$ it suffices to see that $y_0 \in \bar{\mathcal{M}}[G][H]$, whenever H is $\bar{\mathcal{M}}[G]$ generic over $\text{Coll}(\omega, j(\mu))$. Fix an arbitrary such H . Let $J = G \times H$. So it suffices to see that $y_0 \in \bar{\mathcal{M}}[J]$. But $\bar{\mathcal{M}}[J] = \bar{\mathcal{M}}[K]$, for some K which is $\bar{\mathcal{M}}$ -generic over $\text{Coll}(\omega, j(\mu))$. So in order to show that $y_0 \in \mathcal{M}[G]$ it suffices to show that $y_0 \in \bar{\mathcal{M}}[K]$ whenever K is $\bar{\mathcal{M}}$ -generic over $\text{Coll}(\omega, j(\mu))$ and $x_0 \in \bar{\mathcal{M}}[K]$. Since $\bar{\mathcal{M}} \models (j(\mu))^+$ exists, and $\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above $(j(\mu)^+)^{\bar{\mathcal{M}}}$, it suffices to replace $\mathcal{M}$ with $\bar{\mathcal{M}}$, and $\mathbb{P}$ with $\text{Coll}(\omega, j(\mu))$, and κ with $j(\mu)$, and β with $(j(\mu)^+)^{\bar{\mathcal{M}}}$. In other words, without loss of generality we may assume that $\mathbb{P} = \text{Coll}(\omega, \kappa)$ and that $\xi < \kappa$. So we do assume this now.

Then there is a real w in WO such that w codes ξ , and such that w in $\mathcal{M}[G]$. We are assuming that $\{y_0\} \in \Sigma_2(J_\alpha(\mathbb{R}), \{x_0, w\})$. So we may replace x_0 by $\langle x_0, w \rangle$ to yield Claim 1. $\diamond$

So we do now assume that $\{y_0\} \in \Sigma_2(J_\alpha(\mathbb{R}), \{x_0\})$. This means that as a set of integers $y_0 \in \Delta_2(J_\alpha(\mathbb{R}), \{x_0\})$. By lemma 4.9, y_0 is a $\Delta_1(\mathcal{M}[G])$ set of integers. As $\mathcal{M}[G]$ is admissible, $y_0 \in \mathcal{M}[G]$.

CASE 4 $\text{cof}(\alpha) \leq \omega$ and $n = 2$. We are assuming that $\mathcal{M}$ is explicitly $(\alpha, 2)$ -big above β and that no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 2)$ big above β . This

means that there is an ordinal $\delta \in \mathcal{M}$ with $\beta < \delta$ such that $\mathcal{M} \models \text{“}\delta \text{ is a Woodin cardinal”}$, and $\mathcal{M}$ is explicitly $(\alpha, 1)$ big above δ .

Let $x_0 \in \mathbb{R} \cap \mathcal{M}[G]$, and let $y_0 \in \text{OD}_3^{J_\alpha(\mathbb{R})}(x_0)$. We will show that $y_0 \in \mathcal{M}[G]$. Fix $\xi < \omega_1$ such that for all $w \in \text{WO}$ with $|w| = \xi$, $\{y_0\} \in \Sigma_3(J_\alpha(\mathbb{R}), \{x_0, w\})$.

Exactly as in Case 3, we may now ignore the ordinal parameter ξ and assume that $\{y_0\} \in \Sigma_3(J_\alpha(\mathbb{R}), \{x_0\})$. Recall that there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ partial function from $\mathbb{R}$ onto $J_\alpha(\mathbb{R})$. This implies that $\Sigma_3(J_\alpha(\mathbb{R})) = \exists^\mathbb{R} \Pi_2(J_\alpha(\mathbb{R}))$. Fix a Π_2 formula φ such that

$$\{y_0\} = \{y \mid (\exists z \in \mathbb{R}) J_\alpha(\mathbb{R}) \models \varphi[x_0, y, z]\}.$$

Fix $z_0 \in \mathbb{R}$ such that $J_\alpha(\mathbb{R}) \models \varphi[x_0, y_0, z_0]$. We now apply Lemma 4.1. This gives us a partial order $\mathbb{Q} \subseteq J_\delta^\mathcal{M}$, with $\mathbb{Q} \in \mathcal{M}$, and a 0-maximal iteration tree $\mathcal{T}$ on $\mathcal{M}$ of countable length $\theta + 1$ such that

- (a) $\mathcal{M}_\theta^\mathcal{T}$ is realizable, and
- (b) $D^\mathcal{T} = \emptyset$ so that $i_{0,\theta}^\mathcal{T}$ is defined, and
- (c) $\text{lh } E_\xi^\mathcal{T} < i_{0,\xi}^\mathcal{T}(\delta)$ for all $\xi < \theta$, and
- (d) $\text{crit } E_\xi^\mathcal{T} > \beta$ for all $\xi < \theta$ (so G is $\mathcal{M}_\theta^\mathcal{T}$ -generic over $\mathbb{P}$), and
- (e) $\langle y_0, z_0 \rangle$ is $\mathcal{M}_\theta^\mathcal{T}[G]$ -generic over $i_{0,\theta}^\mathcal{T}(\mathbb{Q})$.

As in the proof of Claim 1, we may apply Lemma 3.34 to conclude that $\mathcal{M}_\theta^\mathcal{T}$ is explicitly $(\alpha, 1)$ -big above $i_{0,\theta}^\mathcal{T}(\delta)$. Let $\bar{\mathcal{M}} = \mathcal{M}_\theta^\mathcal{T}$, and $\bar{\delta} = i_{0,\theta}^\mathcal{T}(\delta)$. Since $\mathbb{R} \cap \bar{\mathcal{M}}[G] = \mathbb{R} \cap \mathcal{M}[G]$, it suffices to show that $y_0 \in \bar{\mathcal{M}}[G]$.

$\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ big above $\bar{\delta}$, and no proper initial segment of $\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above $\bar{\delta}$. This implies that $\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ big above $(\bar{\delta}^+)^{\bar{\mathcal{M}}}$. By Lemma 4.9 applied to the formula $\neg\varphi$, there is a Σ_1 formula ψ , and there is a parameter $a \in \bar{\mathcal{M}}$ such that whenever K is $\bar{\mathcal{M}}$ -generic over $\mathbb{P} \times \text{Coll}(\omega, \bar{\delta})$, and $x, y, z \in \mathbb{R} \cap \bar{\mathcal{M}}[K]$, we have

$$J_\alpha(\mathbb{R}) \models \varphi[x, y, z]$$

iff

$$\bar{\mathcal{M}}[K] \models \psi[x, y, z, a, K].$$

Fix such a formula ψ and such a parameter a .

Since $\bar{\mathcal{M}}[G]$ is admissible, to show that $y_0 \in \bar{\mathcal{M}}[G]$ it is sufficient to show that y_0 is a $\Delta_1(\bar{\mathcal{M}}[G])$ set of integers. That is the content of the following claim. Let $\dot{H}$ be the canonical $\text{Coll}(\omega, \bar{\delta})$ -name in $\bar{\mathcal{M}}[G]$ such that whenever H is $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta})$, $\dot{H}_H = H$.

Claim. *Let $n, m \in \omega$. Then the following are equivalent:*

- (a) $y_0(n) = m$
- (b) $\bar{\mathcal{M}}[G] \models \exists q \in \text{Coll}(\omega, \bar{\delta}) \text{ s.t. } q \Vdash “(\exists y, z \in \mathbb{R}) [\psi(\check{x}_0, y, z, \check{a}, \check{G} \times \dot{H}) \wedge y(n) = m].”$
- (c) $\bar{\mathcal{M}}[G] \models \forall q \in \text{Coll}(\omega, \bar{\delta}), q \Vdash “(\forall y, z \in \mathbb{R}) [\psi(\check{x}_0, y, z, \check{a}, \check{G} \times \dot{H}) \implies y(n) = m].”$

Proof of Claim. (a) $\Rightarrow$ (b). Suppose $y_0(n) = m$. Let H be $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta})$ and such that $\langle y_0, z_0 \rangle \in \bar{\mathcal{M}}[G][H]$. Then $G \times H$ is $\bar{\mathcal{M}}$ -generic over $\mathbb{P} \times \text{Coll}(\omega, \bar{\delta})$ and $J_\alpha(\mathbb{R}) \models \varphi[x_0, y_0, z_0]$, so $\bar{\mathcal{M}}[G][H] \models \psi[x_0, y_0, z_0, a, G \times H]$. From this (b) easily follows. (b) $\Rightarrow$ (a). Fix q witnessing that (b) holds. Let H be $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta})$ and such that $q \in H$. Then fix $y, z \in \mathbb{R}$ such that $y(n) = m$ and $\bar{\mathcal{M}}[G][H] \models \psi[x_0, y, z, a, G \times H]$. Then $J_\alpha(\mathbb{R}) \models \varphi[x_0, y, z]$, so $y = y_0$.

(a) $\Rightarrow$ (c). Suppose $y_0(n) = m$. Let H be $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta})$. Let $y, z \in \mathbb{R} \cap \bar{\mathcal{M}}[G][H]$, and suppose that $\bar{\mathcal{M}}[G][H] \models \psi[x_0, y, z, a, G \times H]$. Then $J_\alpha(\mathbb{R}) \models \varphi[x_0, y, z]$, so $y = y_0$. So $y(n) = m$. Since H, y, z were arbitrary, (c) follows.

(c) $\Rightarrow$ (a). Let H be $\bar{\mathcal{M}}[G]$ -generic over $\text{Coll}(\omega, \bar{\delta})$ and such that $\langle y_0, z_0 \rangle \in \bar{\mathcal{M}}[G][H]$. Since $J_\alpha(\mathbb{R}) \models \varphi[x_0, y_0, z_0]$, we have that $\bar{\mathcal{M}}[G][H] \models \psi[x_0, y_0, z_0, a, G \times H]$. By (c), $y_0(n) = m$. $\diamond$

This completes CASE 4.

CASE 5 $n \geq 1$ is odd, and if $\text{cof}(\alpha) \leq \omega$, then $n \geq 3$. We are assuming that $\mathcal{M}$ is explicitly (α, n) -big above β and that no proper initial segment of $\mathcal{M}$ is explicitly (α, n) -big above β . If $\text{cof}(\alpha) > \omega$ this means that $n \geq 1$ and there are n ordinals $\delta_1, \dots, \delta_n \in \mathcal{M}$ with $\beta < \delta_1 < \dots < \delta_n$ such that for $1 \leq i \leq n$, $\mathcal{M} \models “\delta_i \text{ is a Woodin cardinal}”$, and $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above δ_n . If $\text{cof}(\alpha) \leq \omega$ this means that $n \geq 3$ and there are $n - 1$

ordinals $\delta_1, \dots, \delta_{n-1} \in \mathcal{M}$ with $\beta < \delta_1 < \dots < \delta_{n-1}$ such that for $1 \leq i \leq n-1$, $\mathcal{M} \models \text{"}\delta_i \text{ is a Woodin cardinal"}$, and $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above δ_{n-1} .

Let $x_0 \in \mathbb{R} \cap \mathcal{M}[G]$, and let $y_0 \in \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}(x_0)$. We will show that $y_0 \in \mathcal{M}[G]$. Fix $\xi < \omega_1$ such that for all $w \in \text{WO}$ with $|w| = \xi$, $\{y_0\} \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x_0, w\})$.

Exactly as in Case 3, we may now ignore the ordinal parameter ξ and assume that $\{y_0\} \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x_0\})$. Recall that there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ partial function from $\mathbb{R}$ onto $J_\alpha(\mathbb{R})$. This implies that $\Sigma_{n+1}(J_\alpha(\mathbb{R})) = \exists^\mathbb{R} \Pi_n(J_\alpha(\mathbb{R}))$. Fix a Π_n formula φ such that

$$\{y_0\} = \{y \mid (\exists z \in \mathbb{R}) J_\alpha(\mathbb{R}) \models \varphi[x_0, y, z]\}.$$

Fix $z_0 \in \mathbb{R}$ such that $J_\alpha(\mathbb{R}) \models \varphi[x_0, y_0, z_0]$. We now apply Lemma 4.1. This gives us a partial order $\mathbb{Q} \subseteq J_{\delta_1}^\mathcal{M}$, with $\mathbb{Q} \in \mathcal{M}$, and a 0-maximal iteration tree $\mathcal{T}$ on $\mathcal{M}$ of countable length $\theta + 1$ such that

- (a) $\mathcal{M}_\theta^\mathcal{T}$ is realizable, and
- (b) $D^\mathcal{T} = \emptyset$ so that $i_{0,\theta}^\mathcal{T}$ is defined, and
- (c) $\text{lh } E_\xi^\mathcal{T} < i_{0,\xi}^\mathcal{T}(\delta_1)$ for all $\xi < \theta$, and
- (d) $\text{crit } E_\xi^\mathcal{T} > \beta$ for all $\xi < \theta$ (so G is $\mathcal{M}_\theta^\mathcal{T}$ -generic over $\mathbb{P}$), and
- (e) $\langle y_0, z_0 \rangle$ is $\mathcal{M}_\theta^\mathcal{T}[G]$ -generic over $i_{0,\theta}^\mathcal{T}(\mathbb{Q})$.

Let $\bar{\mathcal{M}} = \mathcal{M}_\theta^\mathcal{T}$, and let $\bar{\delta}_i = i_{0,\theta}^\mathcal{T}(\delta_i)$. Since $\mathbb{R} \cap \bar{\mathcal{M}}[G] = \mathbb{R} \cap \mathcal{M}[G]$, it suffices to show that $y_0 \in \bar{\mathcal{M}}[G]$.

Notice that $\bar{\mathcal{M}}$ is explicitly $(\alpha, n-1)$ -big above $(\bar{\delta}_1^+)^{\bar{\mathcal{M}}}$. (If $\text{cof}(\alpha) \leq \omega$ then, as in the proof of Claim 1, we may apply Lemma 3.34 to conclude that $\bar{\mathcal{M}}$ is explicitly $(\alpha, 1)$ -big above $\bar{\delta}_{n-1}$. If $\text{cof}(\alpha) > \omega$ then it is easy to see that $\bar{\mathcal{M}}$ is explicitly $(\alpha, 0)$ -big above $\bar{\delta}_n$.)

Let H be $\bar{\mathcal{M}}[G]$ generic over $\text{Coll}(\omega, \delta_1)$ and such that $\langle y_0, z_0 \rangle \in \bar{\mathcal{M}}[G][H]$. Let $\bar{\mathbb{R}} = \mathbb{R} \cap \bar{\mathcal{M}}[G][H]$. By the lemma we are proving, for $(\alpha, n-1)$, $\bar{\mathbb{R}}$ is closed under $\text{OD}_n^{J_\alpha(\mathbb{R})}$. In particular $\bar{\mathbb{R}}$ is closed downward under $\Delta_{(\alpha, n-1)}$. Also, clearly $\bar{\mathbb{R}} = \mathbb{R} \cap L(\bar{\mathbb{R}})$. By assumption, $\max(\text{code}(\alpha)) < \beta < \bar{\delta}_1 < (\omega_1)^{L(\bar{\mathbb{R}})}$. Since $n-1$ is even, we can apply Lemma 2.36 to conclude that there is some ordinal $\bar{\alpha}$ in $\bar{\mathcal{M}}$ such that $J_{\bar{\alpha}}(\bar{\mathbb{R}})$ is Σ_n elementarily embeddable into $J_\alpha(\mathbb{R})$.

Since φ is a Π_n formula and $y_0, z_0 \in \bar{\mathbb{R}}$, it follows that in $\bar{\mathcal{M}}[G][H]$ the following is true:

$$\{y_0\} = \{y \mid (\exists z \in \mathbb{R}) J_\alpha(\mathbb{R}) \models \varphi[x_0, y, z]\}.$$

As the collapse forcing is homogeneous, this implies that $y_0 \in \bar{\mathcal{M}}[G]$.

This completes CASE 5.

CASE 6 $n \geq 2$ is even, and if $\text{cof}(\alpha) \leq \omega$ then $n \geq 4$. We are assuming that $\mathcal{M}$ is explicitly (α, n) -big above β and that no proper initial segment of $\mathcal{M}$ is explicitly (α, n) -big above β . This means that there are two ordinals, δ_1, δ_2 , such that $\beta < \delta_1 < \delta_2$ and both δ_1 and δ_2 are Woodin in $\mathcal{M}$ and $\mathcal{M}$ is explicitly $(\alpha, n-2)$ -big above δ_2 .

Let $x_0 \in \mathbb{R} \cap \mathcal{M}[G]$, and let $y_0 \in \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}(x_0)$. We will show that $y_0 \in \mathcal{M}[G]$. Fix $\xi < \omega_1$ such that for all $w \in \text{WO}$ with $|w| = \xi$, $\{y_0\} \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x_0, w\})$.

Exactly as in Case 3, we may now ignore the ordinal parameter ξ and assume that $\{y_0\} \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x_0\})$. The proof that we may ignore the ordinal parameter ξ also yields the following:

Claim. *Without loss of generality we may assume that $\max(\text{code}(\alpha))$ is countable in $\mathcal{M}[G]$.*

Proof of Claim. Recall that $\beta \geq (\kappa^+)^{\mathcal{M}}$, where $\mathbb{P} \subseteq J_\kappa^{\mathcal{M}}$. By assumption $\max(\text{code}(\alpha)) < \beta$. So, exactly as in the proof of Claim 1 in CASE 3, without loss of generality we may assume that $\max(\text{code}(\alpha)) < \kappa$ and $\mathbb{P} = \text{Coll}(\omega, \kappa)$. $\diamond$

So we do now assume that $\max(\text{code}(\alpha))$ is countable in $\mathcal{M}[G]$. We will now show that $\mathcal{M}[G]$ is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ correct. More precisely, let $\bar{\mathbb{R}} = \mathbb{R} \cap \mathcal{M}[G]$. Let $s = \text{code}(\alpha)$ and let $\bar{\alpha} = |s|^{\mathcal{M}[G]}$. We will show that there is a Σ_{n+1} elementary embedding

$$j : J_{\bar{\alpha}}(\bar{\mathbb{R}}) \rightarrow J_\alpha(\mathbb{R}).$$

Since, $\{y_0\} \in \Sigma_{n+1}(J_\alpha(\mathbb{R}), \{x_0\})$, this will give us that $y_0 \in \bar{\mathbb{R}}$.

Remark 4.10. The original $\mathcal{M}[G]$ with which we started the proof may *not* be $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -correct. Because of our application of the above Claim, we are now working with a (possibly) modified $\mathcal{M}[G]$ in which $\max(\text{code}(\alpha))$ is countable. This is what will give us that $\bar{\mathbb{R}}$ is correct.

As $\mathcal{M}$ is explicitly (α, n) -big above β , in particular $\mathcal{M}$ is explicitly $(\alpha, n-2)$ -big above β . By the lemma we are proving, for $(\alpha, n-2)$, $\bar{\mathbb{R}}$ is closed under $\text{OD}_{n-1}^{J_\alpha(\mathbb{R})}$. In particular, $\bar{\mathbb{R}}$ is closed down under $\Delta_{(\alpha, n-2)}$. Also, $\bar{\mathbb{R}} = \mathbb{R} \cap L(\bar{\mathbb{R}})$. Also, by assumption, $\max(\text{code}(\alpha)) < \omega_1^{L(\bar{\mathbb{R}})}$. By Lemma 2.36, since $n-2$ is even, there is a Σ_{n-1} elementary map $j : (J_{\bar{\alpha}}(\bar{\mathbb{R}}); \in, \bar{\mathbb{R}}) \rightarrow (J_\alpha(\mathbb{R}); \in, \mathbb{R})$. We will complete our proof by showing that j is Σ_{n+1} elementary.

Recall that there is a $\Sigma_1(J_\alpha(\mathbb{R}))$ partial function from $\mathbb{R}$ onto $J_\alpha(\mathbb{R})$. So to show that j is $\Sigma_{n+1}(J_\alpha(\mathbb{R}))$ -elementary, it suffices to show that for every Π_n formula φ and every $x \in \bar{\mathbb{R}}$, if $J_\alpha(\mathbb{R}) \models (\exists y \in \mathbb{R})\varphi[x, y]$, then $J_{\bar{\alpha}}(\bar{\mathbb{R}}) \models (\exists y \in \bar{\mathbb{R}})\varphi[x, y]$. So fix a Π_n formula φ , and fix an $x \in \bar{\mathbb{R}}$. Now fix a $y \in \mathbb{R}$ and suppose that $J_\alpha(\mathbb{R}) \models \varphi[x, y]$. We will show that $J_{\bar{\alpha}}(\bar{\mathbb{R}}) \models (\exists y \in \bar{\mathbb{R}})\varphi[x, y]$.

We now apply Lemma 4.1. This gives us a partial order $\mathbb{Q} \subseteq J_{\delta_1}^{\mathcal{M}}$, with $\mathbb{Q} \in \mathcal{M}$, and a 0-maximal iteration tree $\mathcal{T}$ on $\mathcal{M}$ of countable length $\theta + 1$ such that

- (a) $\mathcal{M}_\theta^{\mathcal{T}}$ is realizable, and
- (b) $D^{\mathcal{T}} = \emptyset$ so that $i_{0, \theta}^{\mathcal{T}}$ is defined, and
- (c) $\text{lh } E_\xi^{\mathcal{T}} < i_{0, \xi}^{\mathcal{T}}(\delta_1)$ for all $\xi < \theta$, and
- (d) $\text{crit } E_\xi^{\mathcal{T}} > \beta$ for all $\xi < \theta$ (so G is $\mathcal{M}_\theta^{\mathcal{T}}$ -generic over $\mathbb{P}$), and
- (e) y is $\mathcal{M}_\theta^{\mathcal{T}}[G]$ -generic over $i_{0, \theta}^{\mathcal{T}}(\mathbb{Q})$.

Let $\bar{\mathcal{M}} = \mathcal{M}_\theta^{\mathcal{T}}$, and for $1 \leq i \leq 2$, let $\bar{\delta}_i = i_{0, \theta}^{\mathcal{T}}(\delta_i)$. Notice that $\mathbb{R} \cap \bar{\mathcal{M}}[G] = \bar{\mathbb{R}}$. As in the previous case, we have that $\bar{\mathcal{M}}$ is explicitly $(\alpha, n-2)$ -big above $\bar{\delta}_2$. As no proper initial segment of $\bar{\mathcal{M}}$ is explicitly $(\alpha, n-2)$ -big above $\bar{\delta}_2$, we have that $\bar{\mathcal{M}}$ is explicitly $(\alpha, n-2)$ -big above $(\bar{\delta}_2^+)^{\bar{\mathcal{M}}}$.

Since y is $\bar{\mathcal{M}}[G]$ generic over $i_{0, \theta}^{\mathcal{T}}(\mathbb{Q})$, and $i_{0, \theta}^{\mathcal{T}}(\mathbb{Q}) \subset J_{\delta_1}^{\bar{\mathcal{M}}}$, we can absorb y into a generic object for the stationary tower forcing over $\bar{\mathcal{M}}[G]$ up to $\bar{\delta}_2$. (For a description of Woodin's Stationary Tower Forcing, see [Ma2].) Let H be this generic object. Let $\mathcal{N}$ be the generic ultrapower of $\bar{\mathcal{M}}[G]$ by H , and let $\pi : \bar{\mathcal{M}}[G] \rightarrow \mathcal{N}$ be the generic elementary embedding. It is one of the fundamental properties of the Stationary Tower forcing that $\mathbb{R} \cap \mathcal{N} = \mathbb{R} \cap \bar{\mathcal{M}}[G][H]$. Let $\mathbb{R}^* = \mathbb{R} \cap \bar{\mathcal{M}}[G][H]$.

Recall that $s = \text{code}(\alpha)$, that $\max(s) < \omega_1^{L(\bar{\mathbb{R}})}$, and that $\bar{\alpha} = |s|^{L(\bar{\mathbb{R}})}$. Let $\alpha^* = |s|^{L(\mathbb{R}^*)}$. Then $\pi(J_{\bar{\alpha}}(\bar{\mathbb{R}})) = J_{\alpha^*}(\mathbb{R}^*)$.

Now $\bar{\mathcal{M}}[G][H]$ is a generic extension of $\bar{\mathcal{M}}$ via a partial order $\mathbb{Z} \subset J_{\delta_2}^{\bar{\mathcal{M}}}$. Since $\bar{\mathcal{M}}$ is explicitly $(\alpha, n-2)$ big above $(\bar{\delta}_2^+)^{\bar{\mathcal{M}}}$, we have by the lemma we are proving, for $(\alpha, n-2)$, that $\mathbb{R}^*$ is closed under $\text{OD}_{n-1}^{J_\alpha(\mathbb{R})}$. In particular, $\mathbb{R}^*$ is closed downward under $\Delta_{(\alpha, n-2)}$. Also, $\mathbb{R}^* = \mathbb{R} \cap L(\mathbb{R}^*)$. By Lemma 2.36, there is a Σ_{n-1} elementary map $i : (J_{\alpha^*}(\mathbb{R}^*); \in, \mathbb{R}^*) \rightarrow (J_\alpha(\mathbb{R}); \in, \mathbb{R})$.

As φ is a Π_n formula, and $x, y \in \mathbb{R}^*$, $J_{\alpha^*}(\mathbb{R}^*) \models \varphi[x, y]$. As π is fully elementary and $x \in \bar{\mathbb{R}}$, we have that $J_{\bar{\alpha}}(\bar{\mathbb{R}}) \models (\exists y \in \bar{\mathbb{R}})\varphi[x, y]$. This completes our proof. $\square$

5. THE RANKED ITERATION GAME

Suppose that $\mathcal{M}$ is a countable, iterable premouse, and $\mathcal{M}$ is (α, n) -big, but every proper initial segment of $\mathcal{M}$ is (α, n) -small. In the previous section we proved that $A_{(\alpha, n)} \subseteq \mathbb{R} \cap \mathcal{M}$. Our next major goal is to prove that $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha, n)}$. This will give us that $A_{(\alpha, n)} = \mathbb{R} \cap \mathcal{M}$, and so $A_{(\alpha, n)}$ is a mouse set.

Showing that $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha, n)}$ essentially amounts to showing that every proper initial segment of $\mathcal{M}$ is sufficiently simply definable. Suppose $\mathcal{N} \triangleleft \mathcal{M}$ is a proper initial segment of $\mathcal{M}$. Suppose further that $\mathcal{N}$ projects to ω . Why should $\mathcal{N}$ be simply definable? The reason is that $\mathcal{N}$ is the unique premouse $\mathcal{N}'$ such that $\mathcal{N}'$ projects to ω , $\mathcal{N}'$ is sound, $\mathcal{N}'$ has the same ordinals as $\mathcal{N}$, and $\mathcal{N}'$ is fully iterable. (This fact is demonstrated by performing a *comparison* of $\mathcal{N}'$ with $\mathcal{N}$.) This would give us a simple definition of $\mathcal{N}$, except for the fact that being fully iterable is not simply definable.

The strategy for showing that $\mathcal{N}$ is simply definable is the following: Find a condition other than “full iterability” which is simply definable, and yet is sufficient to allow us to perform a comparison. Such a condition is called a definable “iterability condition.” That is the topic of this section.

We follow [St5] in using “iteration games” as the basis for our iterability conditions. We work in the theory $\text{ZF} + \text{DC}$.

We will need a coding function $x \mapsto x^*$ from $\mathbb{R}$ onto HC = the collection of hereditarily countable sets. For the sake of concreteness we will describe a particular such coding

function. Let $\mathcal{T}$ be the collection of wellfounded trees on ω . First we define a function $h : \mathcal{T} \xrightarrow{\text{onto}} \text{HC}$ by induction on the rank of a tree $T \in \mathcal{T}$. Let

$$T_{\langle n \rangle} = \{ s \in \omega^{<\omega} \mid \langle n \rangle \smallfrown s \in T \},$$

and set

$$h(T) = \{ h(T_{\langle n \rangle}) \mid \langle n \rangle \in T \}.$$

Now let $\pi : \mathbb{R} \xrightarrow{\text{onto}} \mathcal{P}(\omega^{<\omega})$ be recursive, and set $\text{WF} = \{ x \mid \pi(x) \in \mathcal{T} \}$. Finally, for $x \notin \text{WF}$, set $x^* = \emptyset$, and for $x \in \text{WF}$, set $x^* = h(\pi(x))$. Then if $n \geq 1$ and θ is any Σ_n formula then the set

$$\{ x \in \mathbb{R} \mid \text{HC} \models \theta[x^*] \}$$

is Σ_{n+1}^1 .

We will need several concepts from [St5]. The following definition is taken from that paper.

Definition 5.1. Let $\mathcal{M}$ be a premouse, and $\delta < \text{OR}^{\mathcal{M}}$.

- (a) $k(\mathcal{M}, \delta)$ is the unique $k < \omega$ such that $\mathcal{M}$ is k -sound, $k+1$ solid, and $\rho_{k+1}(\mathcal{M}) \leq \delta < \rho_k(\mathcal{M})$, if such a k exists. $k(\mathcal{M}, \delta)$ is undefined otherwise.
- (b) $\mathcal{M}$ is a δ -mouse iff $k(\mathcal{M}, \delta)$ is defined, and letting $k = k(\mathcal{M}, \delta)$, we have that $\mathcal{M} = \mathcal{H}_{k+1}^{\mathcal{M}}(\delta \cup p_{k+1}(\mathcal{M}))$.
- (c) An iteration tree $\mathcal{T}$ on $\mathcal{M}$ is *above* δ iff $\text{crit}(E) \geq \delta$ for all extenders E used on $\mathcal{T}$.

See [MiSt] for the definitions of the various fine structural terms used above.

Definition 5.2. If $x \in \mathbb{R}$ then we say that x is a *mouse code* iff $x^* = (\mathcal{M}, \delta)$ where $\mathcal{M}$ is a countable premouse and $\delta \in \text{OR}$ and either

- (1) $\delta = \text{OR}^{\mathcal{M}}$, or
- (2) $\delta \in \text{OR}^{\mathcal{M}}$ and $\mathcal{M}$ is a δ -mouse.

We will need a few more concepts from [St5]. The following definition is also taken from that paper.

Definition 5.3. Let $\mathcal{T}$ be an iteration tree of limit length.

- (a) $\delta(\mathcal{T})$ is the supremum of the lengths of the extenders used in $\mathcal{T}$.

- (b) Let b be a cofinal wellfounded branch of $\mathcal{T}$. Then $Q(b, \mathcal{T}) = \mathcal{J}_\alpha^{\mathcal{M}_b^{\mathcal{T}}}$, where α is the largest β such that $\beta = \delta(\mathcal{T})$, or $\delta(\mathcal{T}) < \beta \leq \text{OR}^{\mathcal{M}_b^{\mathcal{T}}}$ and $\mathcal{J}_\beta^{\mathcal{M}_b^{\mathcal{T}}} \models \delta(\mathcal{T})$ is a Woodin cardinal.

In [St5] it is shown that if $\mathcal{M}$ is a δ -mouse and $\mathcal{T}$ is on ω -maximal tree on $\mathcal{M}$ above δ , and $\delta(\mathcal{T}) < \text{OR}^{\mathcal{Q}(b, \mathcal{T})}$, then $Q(b, \mathcal{T})$ is a $\delta(\mathcal{T})$ -mouse.

We will also need from [St5] the notion of a *putative iteration tree*. This is a tree $\mathcal{T}$ of premice which has all of the properties of an iteration tree, except that if $\mathcal{T}$ has successor length, then the last model of $\mathcal{T}$ is allowed to be illfounded.

Let $\alpha \geq 1$. Let x be a mouse code. We now describe $G^\alpha(x)$, the *rank- α iteration game* on x . $G^\alpha(x)$ is a clopen game on $\mathbb{R} \times \alpha$, so a run of $G^\alpha(x)$ is completed after some finite number of rounds.

At the start of round n with $1 \leq n < \omega$, we have an ordinal α_{n-1} and a real x_{n-1} such that x_{n-1} is a mouse code. (We get started by setting $\alpha_0 = \alpha$ and $x_0 = x$.) Round n begins with player I playing an ordinal $\alpha_n < \alpha_{n-1}$. If $\alpha_{n-1} = 0$ so that player I can not play α_n , then player I loses. Otherwise let $(\mathcal{M}, \delta) = x_{n-1}^*$. Then player I must play a real y_n such that $y_n^* = \mathcal{T}$ where $\mathcal{T}$ is an ω -maximal, putative iteration tree on $\mathcal{M}$ above δ . (We allow player I to play the trivial tree with only one model, so that player I can always meet this obligation.) If $\mathcal{T}$ has a last model which is wellfounded then player I loses. Otherwise, player II must play a mouse code x_n such that there is some maximal wellfounded branch b of $\mathcal{T}$ so that $x_n^* = \langle Q(b, \mathcal{T} \upharpoonright \sup(b)), \delta(\mathcal{T} \upharpoonright \sup(b)) \rangle$. If there is no maximal wellfounded branch b of $\mathcal{T}$ then player II loses. This completes round n .

Since there is no infinite decreasing sequence of ordinals, one of the two players will win after a finite number of rounds. So this completes the description of $G^\alpha(x)$.

Definition 5.4. Let $\mathcal{M}$ be a countable δ -mouse. Let $\alpha \geq 1$. Then $\mathcal{M}$ is *rank- α iterable above δ* iff for some (equivalently, for all) $x \in \mathbb{R}$ such that $x^* = (\mathcal{M}, \delta)$, there is a winning quasi-strategy for player II in $G^\alpha(x)$. (From now on we will abbreviate this to “ II wins $G^\alpha(x)$ ”.)

Our immediate goal is to explore the definability of the set

$$\{ x \in \mathbb{R} \mid II \text{ wins } G^\alpha(x) \}.$$

Definition 5.5. For $\alpha \geq 2$ set

$$\lambda(\alpha) = \begin{cases} \omega\alpha & \text{if } \alpha \geq \omega \\ \omega(\alpha - 1) & \text{if } \alpha < \omega. \end{cases}$$

In Corollary 5.8 below we show that if $\alpha \geq 2$ and $\lambda = \lambda(\alpha)$, then the set

$$\{ x \in \mathbb{R} \mid x \text{ is a mouse code} \wedge II \text{ wins } G^\lambda(x) \}$$

is $\Pi_1(J_\alpha(\mathbb{R}))$ uniformly in α .

If x is a mouse code and $p = \langle y_1, x_1, \dots, y_n, x_n \rangle$ is a sequence of reals of even length then we will say that p is a *legal play on x with I to move* iff there are ordinals $\alpha, \alpha_1 \dots \alpha_n$ such that letting $p' = \langle \alpha_1, y_1, x_1, \dots, \alpha_n, y_n, x_n \rangle$ we have that p' is a legal play in $G^\alpha(x)$. Notice that this is equivalent to saying that if we let $p' = \langle n-1, y_1, x_1, \dots, 0, y_n, x_n \rangle$ then p' is a legal play in $G^n(x)$. Notice that the relation “ p is a legal play on x with I to move” is a simply definable relation of p and x . (The relation is Σ_2^1 .) The definition of a *legal play on x with II to move* is similar.

The following lemma gives an alternate characterization of the property “ II wins $G^\alpha(x)$ ”. The lemma says that we could also have defined this relation on x and α by induction on α .

Lemma 5.6. *Let x be a mouse code and $\alpha \geq 1$.*

- (1) *If $\alpha = \beta + 1$ then II wins $G^\alpha(x)$ iff $(\forall y_1 \in \mathbb{R})(\exists x_1 \in \mathbb{R})[\langle y_1 \rangle \text{ is a legal play on } x \text{ with } II \text{ to move} \Rightarrow (\langle y_1, x_1 \rangle \text{ is a legal play on } x \text{ with } I \text{ to move, and } II \text{ wins } G^\beta(x_1))]$.*
- (2) *If α is a limit ordinal then II wins $G^\alpha(x)$ iff $(\forall \beta < \alpha) II$ wins $G^\beta(x)$.*

Proof. One direction of both (1) and (2) is trivial. For the other direction, the cautious reader might ask whether we need to use the Axiom of Choice in order to pick winning quasi-strategies to piece together. But since $G^\beta(x)$ is a clopen game, if player II wins

$G^\beta(x)$, then he has a canonical winning quasi-strategy. So the Axiom of Choice is in fact not needed. $\square$

Recall the definition of $\lambda(\alpha)$ in Definition 5.5 above. Notice that $\lambda(\alpha) \leq \text{OR}^{J_\alpha(\mathbb{R})}$. For $\gamma \in \text{OR}$, let $F(\gamma) = \{x \in \mathbb{R} : x \text{ is a mouse code and player } II \text{ wins } G^\gamma(x)\}$.

Lemma 5.7. *Let $\alpha \geq 2$. Let $\lambda = \lambda(\alpha)$.*

(1) *For all $\gamma < \lambda$, $F \upharpoonright \gamma \in J_\alpha(\mathbb{R})$.*

(2) *$F \upharpoonright \lambda$ is $\Delta_1(J_\alpha(\mathbb{R}))$.*

Proof. Let P be the binary relation defined by $P(f, \gamma)$ iff $f = F \upharpoonright \gamma$. Notice that P is a Σ_0 relation of f and γ , with $\mathbb{R}$ as a parameter. This follows from the previous lemma.

First we will show that (1) $\implies$ (2). Assuming (1), we have that for $\gamma < \lambda$, $S = F(\gamma)$ iff there is a function $f \in J_\alpha(\mathbb{R})$ such that $P(f, \gamma + 1)$ and $S = f(\gamma)$. This shows that $F \upharpoonright \lambda$ is $\Sigma_1(J_\alpha(\mathbb{R}))$. Since $S = F(\gamma)$ iff $(\forall T) [T = F(\gamma) \rightarrow S = T]$, we have that $F \upharpoonright \lambda$ is also $\Pi_1(J_\alpha(\mathbb{R}))$.

Next we will prove (1) by induction on α . As the basis of our induction, we start with $\alpha = 2$. Then $\lambda(\alpha) = \omega$. For $n < \omega$ it is easy to see that $F \upharpoonright n$ is analytically definable, and so $F \upharpoonright n \in J_2(\mathbb{R})$. If α is a limit ordinal, then $\lambda(\alpha) = \sup \{ \lambda(\beta) \mid \beta < \alpha \}$, and so the inductive step of the proof of (1) is trivial for limit α . Suppose $\alpha = \beta + 1$ is a successor ordinal. Then $\lambda(\alpha) = \lambda(\beta) + \omega$. By (2) for β , $F \upharpoonright \lambda(\beta) \in J_\alpha(\mathbb{R})$. We show by induction on n that $F \upharpoonright (\lambda(\beta) + n) \in J_\alpha(\mathbb{R})$. Let $f = F \upharpoonright (\lambda(\beta) + n)$, and $g = F \upharpoonright (\lambda(\beta) + n + 1)$. Suppose $f \in J_\alpha(\mathbb{R})$. Let $S = g(\lambda(\beta) + n)$. Then $S \subset \mathbb{R}$ and, by the previous lemma, S is Σ_0 definable with f and $\mathbb{R}$ as parameters. Thus $S \in J_\alpha(\mathbb{R})$. As $g = f \cup \{ \langle \lambda(\beta) + n, S \rangle \}$, we have that $g \in J_\alpha(\mathbb{R})$. $\square$

Corollary 5.8. *Let $\alpha \geq 2$. Let $\lambda = \lambda(\alpha)$. Let $n \in \omega$. Then the set*

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G^{\lambda+n}(x) \}$$

is $\Pi_{(\alpha, 2n)}$, uniformly in α .

Proof. For $n = 0$, this follows from part (2) of the previous Lemma and part (2) of Lemma 5.6. For $n > 0$ the result follows by induction on n , using part (1) of Lemma 5.6. $\square$

We have defined rank- α iterability for all ordinals α . Before going on we should point out that the definition only has significance for $\alpha \leq \kappa^{\mathbb{R}}$. To explain this we need to define the *length* ω (weak) iteration game on x , $WG(x)$. We turn to this now. Instead of explicitly giving the definition of $WG(x)$, we only describe how $WG(x)$ differs from the *ranked* iteration game described above: The definition of $WG(x)$ is similar to the definition of the ranked iteration game, except that player I does not play any ordinals. So a run of $WG(x)$ can last for ω rounds, in which case we say that II wins. So $WG(x)$ is a game on $\mathbb{R}$ with closed payoff for player II . This completes our description of $WG(x)$. We will also refer to $WG(x)$ as “the length ω weak iteration game on x .”

Note. Our definition of the weak iteration game differs slightly from definitions given in other papers. See for example [St5] and [St2]. The reason for the name “weak” is that there is something else called the strong iteration game. See [MaSt]. We will have no need for the strong iteration game in this paper.

Let $\kappa^{\mathbb{R}}$ be the least ordinal κ such that $J_{\kappa}(\mathbb{R})$ is admissible. The following is a well-known fact about admissible sets and closed games. The result has nothing to do with the particular details of the iteration games we have defined.

Proposition 5.9. *Let x be a mouse code. Then the following are equivalent:*

- (1) *Player II wins $WG(x)$.*
- (2) *For all ordinals α player II wins $G^{\alpha}(x)$.*
- (3) *Player II wins $G^{\kappa^{\mathbb{R}}}(x)$.*

Proof. Clearly $(1) \Rightarrow (2) \Rightarrow (3)$. We will show that $(3) \Rightarrow (1)$. For this it suffices to show that for all mouse codes x such that player II wins $G^{\kappa^{\mathbb{R}}}(x)$, and for all legal first moves y_0 by player I in $WG(x)$, there is an x_0 , a legal response by player II in $WG(x)$, such that player II wins the game $G^{\kappa^{\mathbb{R}}}(x_0)$. Suppose this were false. Let x and y_0 be a counterexample. For each x_0 which is a legal response by player II in $WG(x)$, let $f(x_0)$ be the least ordinal α such that player II does not win $G^{\alpha}(x_0)$. Since player II wins $G^{\kappa^{\mathbb{R}}}(x)$, the range of f is cofinal in $\kappa^{\mathbb{R}}$. But f is Σ_1 definable over $J_{\kappa^{\mathbb{R}}}(\mathbb{R})$. This contradicts the fact that $J_{\kappa^{\mathbb{R}}}(\mathbb{R})$ is admissible. $\square$

Definition 5.10. Let $\mathcal{M}$ be a countable δ -mouse. Then $\mathcal{M}$ is *length- ω iterable above δ* iff for some (equivalently, for all) $x \in \mathbb{R}$ such that $x^* = (\mathcal{M}, \delta)$, there is a winning quasi-strategy for player II in $WG(x)$.

In the case $n = 0$, Corollary 5.8 above tells us that the set

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G^{\lambda(\alpha)}(x) \}$$

is $\Pi_{(\alpha,0)}$ definable. This result is satisfactory for our applications in the sequel. However, if $n \geq 1$, then Corollary 5.8 tells us that the set

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G^{\lambda(\alpha)+n}(x) \}$$

is $\Pi_{(\alpha,2n)}$ definable. This is *not* satisfactory for our applications. Our solution is to replace $G^{\lambda(\alpha)+n}(x)$ with a different game in the case $n > 1$. Let x be a mouse code. Suppose $2 \leq \alpha < \kappa^{\mathbb{R}}$, and let $\lambda = \lambda(\alpha)$. Let $n \in \omega$. As in [St5] we now define a game $G_{(\alpha,n)}(x)$ which is an approximation of the the game $G^{\lambda+n}(x)$. The reason we do this is that under the assumption of $\text{Det}(J_{\alpha+1}(\mathbb{R}))$, the set

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G_{(\alpha,n)}(x) \}$$

will be $\Pi_{(\alpha,n)}$, uniformly in α . This beats the conclusion of Lemma 5.8 by a factor of 2.

As we mentioned above, for $n = 0$ the game $G^\lambda(x)$ is perfectly satisfactory, and so we set $G_{(\alpha,0)}(x) =_{\text{def}} G^\lambda(x)$. For $n > 0$, $G_{(\alpha,n)}(x)$ will not be exactly the same as $G^{\lambda+n}(x)$. The definition of $G_{(\alpha,n)}(x)$ for $n > 0$ depends on whether n is even or odd. The idea is to use the Bounded Quantification Lemma, that is Lemma 2.16 on page 20, in order to reduce the number of quantifiers needed to define the game.

Note. The clever idea of using the Bounded Quantification Lemma to simplify the definition of the iteration game is *not* due to the author. The idea appears in [St5].

First let $n \geq 2$ be even. Then $G_{(\alpha,n)}(x)$ is played in n rounds. At the start of round k , where $1 \leq k \leq n$, we have a mouse code x_{k-1} . (We get started by setting $x_0 = x$.) The rules for round k depend on whether k is even or odd. First suppose that $1 \leq k \leq n-1$ and k is odd. Round k is played as follows: Let $(\mathcal{M}, \delta) = x_{k-1}^*$. Player I must play two

reals, y_k and z_k , such that $y_k^* = \mathcal{T}$ where $\mathcal{T}$ is an ω -maximal, putative iteration tree on $\mathcal{M}$ above δ , and z_k is arbitrary. (We allow player I to play the trivial tree with only one model, so that player I can always meet this obligation.) If $\mathcal{T}$ has a last model which is wellfounded then player I loses. Otherwise, player II must play a real x_k such that x_k is a mouse code and there is some maximal wellfounded branch b of $\mathcal{T}$ so that $x_k^* = \langle Q(b, \mathcal{T} \restriction \sup(b)), \delta(\mathcal{T} \restriction \sup(b)) \rangle$. If there is no maximal wellfounded branch of $\mathcal{T}$ then player II loses. This completes round k in the case that k is odd.

Now suppose that $2 \leq k \leq n$ and k is even. Round k is played as follows: Let $(\mathcal{N}, \gamma) = x_{k-1}^*$. Player I must play a real, y_k such that:

- (i) $y_k^* = \mathcal{U}$ where $\mathcal{U}$ is an ω -maximal, putative iteration tree on $\mathcal{N}$ above γ . (We allow player I to play the trivial tree with only one model, so that player I can always meet this obligation.) And,
- (ii) $y_k \in \Delta_{(\alpha, n-k+1)}(\langle x_{k-1}, z_{k-1} \rangle)$.

If there is no such real y_k , or if $\mathcal{U}$ has a last model which is wellfounded, then player I loses. Otherwise, player II must play a real x_k such that x_k is a mouse code and there is some maximal wellfounded branch c of $\mathcal{U}$ so that $x_k^* = \langle Q(c, \mathcal{U} \restriction \sup(c)), \delta(\mathcal{U} \restriction \sup(c)) \rangle$. If there is no maximal wellfounded branch of $\mathcal{U}$ then player II loses. This completes round k in the case that k is even.

After n rounds of play, if neither player has lost then the play will have produced a real x_n . Then player II wins this run of $G_{(\alpha, n)}(x)$ iff player II has a winning quasi-strategy in the rank- λ iteration game on x_n , $G^\lambda(x_n)$, where $\lambda = \lambda(\alpha)$. This completes the description of $G_{(\alpha, n)}(x)$ in the case that $n \geq 2$ is even.

Notice that for even $n \geq 2$, the game $G_{(\alpha, n)}(x)$ demands more of Player I than does the game $G^{\lambda+n}(x)$. Thus we have the following.

Proposition 5.11. *Let $n \geq 2$ be even. If II wins $G^{\lambda+n}(x)$ then II wins $G_{(\alpha, n)}(x)$.*

Furthermore we have the following.

Lemma 5.12. *Suppose $2 \leq \alpha < \kappa^{\mathbb{R}}$. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \geq 2$ be even. Then the set*

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G_{(\alpha, n)}(x) \}$$

is $\Pi_{(\alpha,n)}$, uniformly in α .

Proof. There is a relation $R \in \Pi_1(J_\alpha(\mathbb{R}))$ such that if x is a mouse code, then II wins $G_{(\alpha,n)}(x)$ iff

$$(\forall y_1, z_1)(\exists x_1)(\forall y_2 \in \Delta_{(\alpha,n-1)}(\langle x_1, z_1 \rangle)(\exists x_2) \cdots \\ (\forall y_{n-1}, z_{n-1})(\exists x_{n-1})(\forall y_n \in \Delta_{(\alpha,1)}(\langle x_{n-1}, z_{n-1} \rangle)(\exists x_n)R(x, x_1, y_1, \dots, x_n, y_n)).$$

Now $R \in \Pi_{(\alpha,0)}$, and for odd $k \geq 1$, $\Pi_{(\alpha,0)} \subseteq \Sigma_{(\alpha,k)}$. So $R \in \Sigma_{(\alpha,k)}$ for all odd k . Using the Bounded Quantification Lemma, it is easy to see that our claim is true. $\square$

For even $n \geq 2$, one should think of the game $G_{(\alpha,n)}(x)$ defined above as an approximation of the game $G^{\lambda+n}(x)$, where $\lambda = \lambda(\alpha)$. Notice that the game $G^\lambda(x)$ is employed in the definition of $G_{(\alpha,n)}(x)$. The relation “ II wins $G^\lambda(x)$ ” is a $\Pi_{(\alpha,0)}$ relation of x , and since n is even $\Pi_{(\alpha,0)} \subseteq \Pi_{(\alpha,n)}$.

Now suppose that $n \geq 1$ is odd. Then, recall that we do not necessarily have that $\Pi_{(\alpha,0)} \subseteq \Pi_{(\alpha,n)}$. This means that we can not employ the game $G^\lambda(x)$ in our definition of the game $G_{(\alpha,n)}(x)$. But yet we still want to have that the game $G_{(\alpha,n)}(x)$ is an “approximation” of the game $G^{\lambda+n}(x)$. So we must use a slightly different strategy. We continue with our definition of $G_{(\alpha,n)}(x)$ below.

Let $n \geq 1$ be odd. Then $G_{(\alpha,n)}(x)$ is played in $n + 1$ rounds. At the start of round k , where $1 \leq k \leq n + 1$, we have a mouse code x_{k-1} . (We get started by setting $x_0 = x$.) Round k is played as follows: Let $(\mathcal{M}, \delta) = x_{k-1}^*$. Player I must play a real y_k such that $y_k^* = \mathcal{T}$ where $\mathcal{T}$ is an ω -maximal, putative iteration tree on $\mathcal{M}$ above δ . (We allow player I to play the trivial tree with only one model, so that player I can always meet this obligation.) If $\mathcal{T}$ has a last model which is wellfounded then player I loses. Otherwise, player II must play a real x_k such that:

- (i) x_k is a mouse code and there is some maximal wellfounded branch b of $\mathcal{T}$ so that $x_k^* = \langle Q(b, \mathcal{T} \upharpoonright \sup(b)), \delta(\mathcal{T} \upharpoonright \sup(b)) \rangle$. And,
- (ii) $x_k \in \Delta_{(\alpha,n)}(y_k)$.

If there is no such x_k then player II loses. This completes round k .

After $n + 1$ rounds of play, if neither player has lost then the play will have produced a real x_{n+1} . Then player II wins this run of $G_{(\alpha,n)}(x)$ iff $(\exists \sigma \in J_\alpha(\mathbb{R}))$ such that σ is a winning quasi-strategy for player II in the *length* ω (weak) iteration game on x_{n+1} , $WG(x_{n+1})$. This completes the description of $G_{(\alpha,n)}(x)$ in the case that n is odd.

Notice that for odd $n \geq 1$, the game $G_{(\alpha,n)}(x)$ demands more of Player II than does the game $G^{\lambda+n}(x)$. In fact, it demands more of Player II than does the game $WG(x)$. Thus we have the following.

Proposition 5.13. *Let $n \geq 1$ be odd. If II wins $G_{(\alpha,n)}(x)$ then II wins $WG(x)$, and so in particular II wins $G^{\lambda+n}(x)$.*

Furthermore we have the following.

Lemma 5.14. *Suppose $2 \leq \alpha < \kappa^{\mathbb{R}}$. Assume $\text{Det}(J_{\alpha+1}(\mathbb{R}))$. Let $n \geq 1$ be odd. Then the set*

$$\{ x \mid x \text{ is a mouse code} \wedge II \text{ wins } G_{(\alpha,n)}(x) \}$$

is $\Pi_{(\alpha,n)}$, uniformly in α .

Proof. There is a relation $R \in \Sigma_1(J_\alpha(\mathbb{R}))$ such that if x is a mouse code, then II wins $G_{(\alpha,n)}(x)$ iff

$$(\forall y_1)(\exists x_1 \in \Delta_{(\alpha,n)}(y_1)) \cdots (\forall y_{n+1})(\exists x_{n+1} \in \Delta_{(\alpha,n)}(y_{n+1})) R(x, x_1, y_1, \dots, x_{n+1}, y_{n+1}).$$

Since n is odd, $R \in \Pi_{(\alpha,n)}$. Our claim then follows from the Bounded Quantification Lemma. $\square$

Remark 5.15. The reader may wonder about our formulation of the game $G_{(\alpha,n)}(x)$ in the case that n is odd. In particular it may seem surprising that the game is played in $n + 1$ rounds instead of n . The explanation is that we want to be able to show that player II can win $G_{(\alpha,n)}(x)$ whenever x is a mouse code and, writing $x^* = (\mathcal{M}, \delta)$, we have that $\mathcal{M}$ is $(\alpha, n + 1)$ -small above δ , and $\mathcal{M}$ is fully iterable. This fact will be needed in the proof of Theorem 7.16. In order to be able to show that player II can win $G_{(\alpha,n)}(x)$ in this situation, we need to have the game played for $n + 1$ rounds. Notice that we still get Lemma 5.14 above even though $G_{(\alpha,n)}(x)$ is played in $n + 1$ rounds.

Definition 5.16. Let $\mathcal{M}$ be a countable δ -mouse. Let $\alpha \geq 2$. Let $n \in \omega$. Then $\mathcal{M}$ is $\Pi_{(\alpha,n)}$ *iterable above δ* iff for some (equivalently, for all) $x \in \mathbb{R}$ such that $x^* = (\mathcal{M}, \delta)$, there is a winning quasi-strategy for player II in $G_{(\alpha,n)}(x)$.

For completeness we would like to define $\Pi_{(\alpha,n)}$ iterability in the case $\alpha = 1$. We will say that $\mathcal{M}$ is $\Pi_{(1,n)}$ iterable above δ iff $\mathcal{M}$ is Π_{n-1} iterable above δ in the sense of [St5].

6. PETITE PREMICE

Recall that our goal in this paper is to prove that $A_{(\alpha,n)}$ is a mouse set. In the previous section we defined the notion “ $\mathcal{M}$ is $\Pi_{(\alpha,n)}$ -iterable.” At this point, we would have liked to prove a certain comparison theorem:

Conjecture 6.1. Suppose that $\mathcal{M}$ and $\mathcal{N}$ are countable premice. Suppose that $\mathcal{N}$ is $\Pi_{(\alpha,n)}$ -iterable and $(\alpha, n+1)$ -small, and $\mathcal{M}$ is fully iterable and $(\alpha, n+1)$ -small. Then $\mathcal{M}$ and $\mathcal{N}$ can be compared.

Suppose that $\mathcal{M}$ is a fully iterable, countable premouse, and that $\mathcal{M}$ is $(\alpha, n+1)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\alpha, n+1)$ -small. If true, the above conjecture would imply that $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha,n+1)}$. The results of Section 4 imply that $A_{(\alpha,n+1)} \subseteq \mathbb{R} \cap \mathcal{M}$. So we would have that $A_{(\alpha,n+1)} = \mathbb{R} \cap \mathcal{M}$, and so $A_{(\alpha,n+1)}$ is a mouse set.

Unfortunately we are not able to prove this conjecture—at least, not for every (α, n) . So we are forced to retreat. First, we will define the notion “ $\mathcal{M}$ is (α, n) -petite.” Then, in the next section, we will prove a version of the conjecture above, with “petite” in place of “small”.

Our goal in this section then is to define the notion “ $\mathcal{M}$ is (α, n) -petite.” This will only be a new notion for pairs (α, n) such that $\text{cof}(\alpha) \leq \omega$ and $n \geq 1$. For other pairs, we will define “ (α, n) -petite” to be synonymous with “ (α, n) -small”. But for pairs (α, n) such that $\text{cof}(\alpha) \leq \omega$ and $n \geq 1$, the condition of being (α, n) -petite will be a little more restrictive than the condition of being (α, n) -small. In other words, if $\mathcal{M}$ is (α, n) -petite then $\mathcal{M}$ is (α, n) -small, but the converse does not hold. We believe that the notion of

(α, n) -small is the “natural” notion, and that our notion of (α, n) -petite is an ad-hoc notion with which we are unfortunately forced to deal.

Recall that a mouse $\mathcal{M}$ is (α, n) -small iff $\mathcal{M}$ is not (α, n) -big. To parallel this situation, we will first define the notion “weakly (α, n) -big”, and then we will say that $\mathcal{M}$ is (α, n) -petite iff $\mathcal{M}$ is not weakly (α, n) -big. Furthermore, recall that a mouse $\mathcal{M}$ is (α, n) -big iff $\mathcal{M}$ is explicitly (α', n') -big for some (α', n') with $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$. To parallel this situation, we will first define the notion “weakly, explicitly (α, n) -big”, and then we will say that $\mathcal{M}$ is weakly (α, n) -big iff $\mathcal{M}$ is weakly, explicitly (α', n') -big for some (α', n') with $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$. The name “weakly, explicitly (α, n) -big” is, of course, very awkward. We choose to tolerate this awkwardness, among other reasons, because we believe that the notion itself is ad-hoc and should be dispensable.

If $\text{cof}(\alpha) > \omega$, or if $\alpha = 1$, or if $\text{cof}(\alpha) \leq \omega$ and $n = 0$, then $\mathcal{M}$ is *weakly, explicitly (α, n) -big above β* iff $\mathcal{M}$ is (strongly) explicitly (α, n) -big above β . So we only need give our definition in the case $\text{cof}(\alpha) \leq \omega$ and $n \geq 1$. We start with the case $n = 1$. Compare the following definition with Definition 3.8 on page 38. The main difference between the two definitions is that in the definition below we do not demand that the δ ’s remain cardinals up to the next admissible.

Definition 6.2. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$, and suppose that α is a successor ordinal, or a limit ordinal of cofinality ω . Then $\mathcal{M}$ is *weakly, explicitly $(\alpha, 1)$ -big above β* iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$ with $\beta \in \mathcal{N}$, such that for cofinally many $\alpha' < \alpha$, and all $n \in \omega$, and cofinally many $\delta < \text{OR} \cap \mathcal{N}$, we have that:

- (a) δ is a cardinal of $\mathcal{N}$, and
- (b) δ is locally (α', n) -Woodin in $\mathcal{N}$.

In Figure 1 on page 39, and in the remark immediately following that figure, we explored the definition of (strongly) explicitly $(\alpha, 1)$ -big in the case that α is a successor ordinal. Similarly, in Figure 2 and Remark 6.3 below, we explore the definition of weakly, explicitly $(\alpha, 1)$ -big in the case that α is a successor ordinal.

Remark 6.3. Suppose $\alpha = \alpha' + 1$ with $\alpha' \geq 1$. Suppose that $\mathcal{M}$ is a countable premouse and $\mathcal{M}$ is weakly, explicitly $(\alpha, 1)$ -big (above 0) but no proper initial segment of $\mathcal{M}$ is.




FIGURE 2. $\alpha \geq 2$ is a successor and $\mathcal{M}$ is weakly, explicitly $(\alpha, 1)$ -big

We will examine what $\mathcal{M}$ looks like. (See Figure 2.) There is a sequence of ordinals $\delta_0 < \gamma_0 < \delta_1 < \gamma_1 < \delta_2 < \dots$ such that each δ_n is a cardinal of $\mathcal{M}$ and a Woodin cardinal of $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$, and $\mathcal{J}_{\gamma_n}^{\mathcal{M}}$ is explicitly (α', n) -big above δ_n . Since no proper initial segment of $\mathcal{M}$ is weakly, explicitly $(\alpha, 1)$ -big, we have that $\sup \{\delta_n\} = \sup \{\gamma_n\} = \text{OR}^{\mathcal{M}}$.

Let $\rho = \text{OR}^{\mathcal{M}}$, and let $\mathcal{M}^*$ be the first admissible structure over $\mathcal{M}$. It is tempting to think that $(\mathcal{M}^*, \rho)$ witnesses that $\mathcal{M}^*$ is (strongly) explicitly $(\alpha, 1)$ -big. Note that in general *this is not true*. The reason is that in general ρ will not be a cardinal in $\mathcal{M}^*$.

We now complete the inductive definition of weakly, explicitly (α, n) -big.

Definition 6.4. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$, and suppose that α is a successor ordinal, or a limit ordinal of cofinality ω . Let $n \geq 2$. Then $\mathcal{M}$ is *weakly, explicitly (α, n) -big above β* iff there is an initial segment $\mathcal{N} \trianglelefteq \mathcal{M}$, and an ordinal $\delta \in \mathcal{N}$ with $\beta < \delta$, such that δ is a Woodin cardinal of $\mathcal{N}$, and $\mathcal{N}$ is weakly, explicitly $(\alpha, n-1)$ -big above δ .

It is easy to see that if $\mathcal{M}$ is explicitly (α, n) -big then $\mathcal{M}$ is weakly, explicitly (α, n) -big. Conversely we have the following.

Lemma 6.5. *Let $\mathcal{M}$ be a countable, realizable premouse, and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha < \omega_1^{\omega_1}$, and suppose that α is a successor ordinal, or a limit ordinal of cofinality ω . Let*

$n \geq 0$. Suppose that $\mathcal{M}$ is weakly, explicitly $(\alpha, n+1)$ -big above β . Furthermore, suppose that $\max(\text{code}(\alpha)) < \beta$. Then $\mathcal{M}$ is (strongly) explicitly (α, n) -big above β .

Proof. First consider the case $n = 0$. Suppose that $\mathcal{M}$ is weakly, explicitly $(\alpha, 1)$ -big above β . Then clearly $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β . This does not use our hypothesis that $\max(\text{code}(\alpha)) < \beta$.

Next consider the case $n = 1$. Suppose $\mathcal{M}$ is weakly, explicitly $(\alpha, 2)$ -big above β , and we will show that $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . Let $\mathcal{N} \trianglelefteq \mathcal{M}$, and let δ be an ordinal of $\mathcal{N}$ such that δ is Woodin in $\mathcal{N}$ and $\mathcal{N}$ is weakly, explicitly $(\alpha, 1)$ -big above δ . By the previous paragraph, $\mathcal{N}$ is explicitly $(\alpha, 0)$ -big above δ . By our hypothesis, $\max(\text{code}(\alpha)) < \delta$. By Lemma 3.24 on page 48, $\mathcal{J}_\delta^{\mathcal{N}}$ is explicitly $(\alpha, 1)$ -big above β .

The result now follows trivially for $n \geq 2$. $\square$

Finally, we get to the definition of (α, n) -petite.

Definition 6.6. Let $\mathcal{M}$ be a premouse and $\beta \in \text{OR}^{\mathcal{M}}$. Let $2 \leq \alpha \leq \omega_1^{\omega_1}$, and let $n \in \omega$. Then $\mathcal{M}$ is *weakly* (α, n) -big above β iff $\mathcal{M}$ is weakly, explicitly (α', n') -big above β for some (α', n') with $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$. Finally, $\mathcal{M}$ is (α, n) -petite above β iff $\mathcal{M}$ is not weakly (α, n) -big above β .

In the remainder of this section we will prove a few lemmas concerning our new notion of (α, n) -petite. We mentioned above that this is only a new notion for certain pairs (α, n) . Let us begin by verifying that this is so:

Lemma 6.7. Suppose that $\text{cof}(\alpha) > \omega$, or $n = 0$. Then (α, n) -petite is synonymous with (α, n) -small.

Proof. Let (α, n) be as in the statement of the lemma. If $\mathcal{M}$ is not (α, n) -small above β , then it follows easily that $\mathcal{M}$ is not (α, n) -petite above β . Conversely, suppose that $\mathcal{M}$ is not (α, n) -petite above β . Let (α', n') be the lexicographically least pair such that $(\alpha, n) \leq_{\text{lex}} (\alpha', n')$ and $\mathcal{M}$ is weakly, explicitly (α', n') -big above β . If $(\alpha', n') = (\alpha, n)$ then, because of our conditions on (α, n) , we have that by definition $\mathcal{M}$ is *strongly* explicitly (α, n) -big above β , and so $\mathcal{M}$ is not (α, n) -small above β . So suppose that $(\alpha, n) <_{\text{lex}} (\alpha', n')$. Then the proof of Lemma 3.13 shows that $\alpha' > \alpha$ and

$n' = 0$. So again, by definition, $\mathcal{M}$ is *strongly* explicitly (α', n') -big above β . So $\mathcal{M}$ is not (α, n) -small above β . $\square$

The following lemma is fairly obvious, but the proof is a bit convoluted.

Lemma 6.8. *Let $\mathcal{M}$ be a countable, realizable premouse, and $\beta \in \text{OR}^{\mathcal{M}}$. Suppose $2 \leq \alpha \leq \omega_1^{\omega_1}$. Then $\mathcal{M}$ is $(\alpha, 0)$ -petite above β iff $\mathcal{M}$ is (α', n) -petite above β for some pair $(\alpha', n) <_{\text{lex}} (\alpha, 0)$.*

Proof. If $\mathcal{M}$ is (α', n) -petite above β for some pair $(\alpha', n) <_{\text{lex}} (\alpha, 0)$, then of course $\mathcal{M}$ is $(\alpha, 0)$ -petite above β . Conversely, suppose that for all $(\alpha', n) <_{\text{lex}} (\alpha, 0)$, $\mathcal{M}$ is not (α', n) -petite above β . We will show that $\mathcal{M}$ is not $(\alpha, 0)$ -petite above β . Suppose towards a contradiction that $\mathcal{M}$ is $(\alpha, 0)$ -petite above β .

Since $\mathcal{M}$ is not (α', n) -petite above β for all $(\alpha', n) <_{\text{lex}} (\alpha, 0)$, we have that either:

(i) $\mathcal{M}$ is weakly, explicitly (α', n) -big above β for cofinally many $(\alpha', n) <_{\text{lex}} (\alpha, 0)$,

or

(ii) $\mathcal{M}$ is weakly, explicitly (α', n) -big above β for some (α', n) with $(\alpha, 0) \leq_{\text{lex}} (\alpha', n)$.

In case (ii), we have that $\mathcal{M}$ is not $(\alpha, 0)$ -petite above β . So suppose that case (i) holds. First notice that case (i) can only hold if $\text{cof}(\alpha) \leq \omega$. This is because $\mathcal{M}$ can only be weakly, explicitly (α', n) -big above β for countably many pairs (α', n) . Thus we may assume that $\text{cof}(\alpha) \leq \omega$. We will show that $\mathcal{M}$ is weakly, explicitly $(\alpha, 0)$ -big above β . Recall that this is synonymous with strongly, explicitly $(\alpha, 0)$ -big above β . By definition, we must prove the following:

Claim. *For cofinally many $\alpha' < \alpha$, and all n , there is a proper initial segment of $\mathcal{M}$ which is strongly, explicitly (α', n) -big above β .*

Proof of Claim. First suppose that α is a limit ordinal of cofinality ω . From our assumption that (i) above holds, and using the proof of Lemma 3.13, we have that for cofinally many $\alpha' < \alpha$, $\mathcal{M}$ is weakly explicitly $(\alpha', 0)$ -big above β . Let $\alpha_0 < \alpha$ and let $n_0 \in \omega$. We will find an α' with $\alpha_0 \leq \alpha' < \alpha$, and an $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is strongly, explicitly (α', n_0) -big above β .

Let α_1 be least such that $\alpha_0 < \alpha_1 < \alpha$ and $\mathcal{M}$ is weakly, explicitly $(\alpha_1, 0)$ -big above β . By definition, $\mathcal{M}$ is *strongly* explicitly $(\alpha_1, 0)$ -big above β . If $\alpha_1 = \alpha_0 + 1$, then by

definition there is a proper initial segment of $\mathcal{M}$ which is explicitly (α_0, n_0) -big above β . So we are done if we set $\alpha' = \alpha_0$. So suppose that $\alpha_1 > \alpha_0 + 1$. Then it can not be true that α_1 is a successor ordinal or a limit ordinal of cofinality ω , so it must be the case that α_1 is a limit ordinal of cofinality ω_1 . Let $\mathcal{N}_1 \trianglelefteq \mathcal{M}$ be such that $\mathcal{N}_1$ witnesses that $\mathcal{M}$ is explicitly $(\alpha_1, 0)$ -big above β . By definition, $\mathcal{N}_1$ is active. Let κ_1 be the critical point of the last extender from the $\mathcal{N}_1$ -sequence. In review we have:

- (a) $\alpha_0 < \alpha_1 < \alpha$.
- (b) $\text{cof}(\alpha_1) = \omega_1$.
- (c) $\mathcal{N}_1 \trianglelefteq \mathcal{M}$ witnesses that $\mathcal{M}$ is explicitly $(\alpha_1, 0)$ -big above β .
- (d) κ_1 is the critical point of the last extender from the $\mathcal{N}$ -sequence.

By replacing α_1 and $\mathcal{N}_1$ if necessary, we may also assume that κ_1 is least possible so that (a) through (d) above holds.

Now let α_2 be least such that $\alpha_1 < \alpha_2 < \alpha$ and $\mathcal{M}$ is weakly, explicitly $(\alpha_2, 0)$ -big above β . By definition, $\mathcal{M}$ is *strongly* explicitly $(\alpha_2, 0)$ -big above β . If $\alpha_2 = \alpha_1 + 1$, then by definition there is a proper initial segment of $\mathcal{M}$ which is explicitly (α_1, n_0) -big above β . So we are done if we set $\alpha' = \alpha_1$. So suppose that $\alpha_2 > \alpha_1 + 1$. Then it can not be true that α_2 is a successor ordinal or a limit ordinal of cofinality ω , so it must be the case that α_2 is a limit ordinal of cofinality ω_1 . Let $\mathcal{N}_2 \trianglelefteq \mathcal{M}$ be such that $\mathcal{N}_2$ witnesses that $\mathcal{M}$ is explicitly $(\alpha_2, 0)$ -big above β . By definition, $\mathcal{N}_2$ is active. Let κ_2 be the critical point of the last extender from the $\mathcal{N}_2$ -sequence. By our minimality hypothesis on κ_1 , we have that $\kappa_1 \leq \kappa_2$. By definition, $\max(\text{code}(\alpha_1)) < \kappa_1$. Thus $\max(\text{code}(\alpha_1)) < \kappa_2$. Then, since $\alpha_1 < \alpha_2$ and $\max(\text{code}(\alpha_1)) < \kappa_2$, there is a proper initial segment $\mathcal{N} \triangleleft J_{\kappa_2}^{\mathcal{M}}$ such that $\mathcal{N}$ is explicitly (α_1, n_0) -big above β . Thus we have accomplished our task, with $\alpha' = \alpha_1$.

Now we must prove our claim in the case that α is a successor ordinal. Suppose $\alpha = \alpha_0 + 1$. From our assumption that (i) above holds, and using the proof of Lemma 3.13, we have that $\mathcal{M}$ is weakly, explicitly (α_0, n) -big above β for all $n \in \omega$. Fix $n_0 \in \omega$. We must show that there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is *strongly*, explicitly (α_0, n_0) -big above β . Let $\mathcal{N}_0 \trianglelefteq \mathcal{M}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ which is weakly, explicitly $(\alpha_0, 0)$ -big above β . By Proposition 3.25, if $\mathcal{N}_0$ had ω Woodin cardinals

above β , then $\mathcal{N}_0$ would be $(\omega_1^{\omega_1}, 0)$ -big above β . Since we are assuming that $\mathcal{M}$ is $(\alpha, 0)$ -petite above β , $\mathcal{N}_0$ does not have ω Woodin cardinals above β . Let m be such that $0 \leq m < \omega$ and $\mathcal{N}_0$ has m Woodin cardinals, but not $m + 1$ Woodin cardinals above β . Let $n = m + n_0 + 10$, and let $\mathcal{N}_1 \trianglelefteq \mathcal{M}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ which is weakly, explicitly (α_0, n) -big above β . Since $\mathcal{N}_1$ is also weakly, explicitly $(\alpha_0, 0)$ -big above β we have that $\mathcal{N}_0 \trianglelefteq \mathcal{N}_1$. Since $\mathcal{N}_0$ does not have n Woodin cardinals, we have that $\mathcal{N}_0$ is a proper initial segment of $\mathcal{N}_1$. Since $n > m + n_0 + 1$, $\mathcal{N}_1$ is weakly explicitly $(\alpha_0, n_0 + 1)$ -big above $\text{OR}^{\mathcal{N}_0}$. By Lemma 3.18, $\max(\text{code}(\alpha_0)) \leq \text{OR}^{\mathcal{N}_0}$. Then, by the proof of Lemma 6.5, there is a proper initial segment of $\mathcal{N}_1$ which is *strongly* explicitly (α_0, n_0) -big above $\text{OR}^{\mathcal{N}_0}$, and so above β . $\diamond$

This completes the proof of the lemma. $\square$

The following lemma is also fairly obvious. The proof just requires untangling definitions. We will use this lemma in the next section.

Lemma 6.9. *Let $\mathcal{M}$ be a countable, realizable premouse. Let $\beta < \delta$ be ordinals in $\mathcal{M}$, and suppose that δ is a Woodin cardinal of $\mathcal{M}$. Let $\alpha \geq 2$.*

- (a) *Let $n \geq 1$ and suppose that $\mathcal{M}$ is $(\alpha, n + 1)$ -petite above β . Then $\mathcal{M}$ is (α, n) -petite above δ .*
- (b) *Suppose that $\mathcal{M}$ is $(\alpha, 1)$ -petite above β . Then $\mathcal{M}$ is (α', m) -petite above δ for some (α', m) with $1 \leq \alpha' < \alpha$.*

Proof. (a) Suppose $\mathcal{M}$ is not (α, n) petite above δ . Since $\mathcal{M}$ is $(\alpha, n + 1)$ -petite above δ , we must have that $\mathcal{M}$ is weakly, explicitly (α, n) -big above δ . As $\mathcal{M} \models \delta$ is Woodin, $\mathcal{M}$ is weakly, explicitly $(\alpha, n + 1)$ -big above β . This is a contradiction.

(b) Suppose that for all $\alpha' < \alpha$ and all $m \in \omega$, $\mathcal{M}$ is not (α', m) -petite above δ . By the previous lemma, $\mathcal{M}$ is not $(\alpha, 0)$ -petite above δ . Since $\mathcal{M}$ is $(\alpha, 1)$ -petite above δ , we must have that $\mathcal{M}$ is weakly, explicitly $(\alpha, 0)$ -big above δ . Recall that weakly, explicitly $(\alpha, 0)$ -big is synonymous with (strongly) explicitly $(\alpha, 0)$ -big.

First consider the case that $\text{cof}(\alpha) > \omega$. Since $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above δ and $\mathcal{M} \models \delta$ is Woodin, $\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . This is a contradiction.

Next consider the case that $\text{cof}(\alpha) \leq \omega$.

Claim. $\max(\text{code}(\alpha)) < \delta$.

Proof of Claim. Let k be the least measurable cardinal of $\mathcal{M}$ which is greater than β . So we have that $\beta < \kappa < \delta$. Since $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above β but $\mathcal{M}$ is not $(\alpha^*, 0)$ -big above β for any $\alpha^* > \alpha$, Lemma 3.27 on page 51 implies that $\max(\text{code}(\alpha)) < \kappa$. $\diamond$

We may now apply Lemma 3.24 on page 48 to conclude that $\mathcal{J}_\delta^\mathcal{M}$ is explicitly $(\alpha, 1)$ -big above β . This is a contradiction. $\square$

In Section 4, it was essential to our proof that an iterate of a big premouse is big. In fact this requirement was one of the central motivating factors behind our definition of (α, n) -big. Similarly, in the next section it will be essential to our proof that an iterate of a petite premouse is petite. That is the content of the next lemma.

Lemma 6.10. *Let $\mathcal{M}$ be a premouse, and $\beta \in \text{OR}^\mathcal{M}$. Let $\mathcal{T}$ be an iteration tree on $\mathcal{M}$, with $\mathcal{T}$ above β . Suppose that $\mathcal{M}$ is (α, n) -petite above β . Then $\forall \xi < \text{lh}(\mathcal{T})$, $\mathcal{M}_\xi^\mathcal{T}$ is (α, n) -petite above β .*

Proof. By induction on ξ . For $\xi < \text{lh}(\mathcal{T})$ we will write $\mathcal{M}_\xi$ instead of $\mathcal{M}_\xi^\mathcal{T}$, and E_ξ instead of $E_\xi^\mathcal{T}$.

First suppose $\xi = \eta + 1$ is a successor ordinal. Let $\gamma = T\text{-pred}(\xi)$. Let $k = \deg(\xi)$. Then there is an initial segment $\mathcal{M}_\xi^* \trianglelefteq \mathcal{M}_\gamma$ such that $\mathcal{M}_\xi = \text{Ult}_k(\mathcal{M}_\xi^*, E_\eta)$. Let $j : \mathcal{M}_\xi^* \rightarrow \mathcal{M}_\xi$ be the canonical k -embedding. Let $\kappa = \text{crit}(j) = \text{crit}(E_\eta)$.

By induction, $\mathcal{M}_\gamma$ is (α, n) -petite above β , and so in particular, $\mathcal{M}_\xi^*$ is (α, n) -petite above β . Let (α_0, n_0) be the lexicographically greatest pair such that $\mathcal{M}_\xi^*$ is weakly, explicitly (α_0, n_0) -big above β . As in the proof of Theorem 4.6 on page 62, we can see that there is such a pair (α_0, n_0) . We have that $(\alpha_0, n_0) <_{\text{lex}} (\alpha, n)$, and that $\mathcal{M}_\xi^*$ is $(\alpha_0, n_0 + 1)$ -petite above β . We will show that $\mathcal{M}_\xi$ is $(\alpha_0, n_0 + 1)$ -petite above β .

Claim 1. *There is an ordinal $\mu \leq \kappa$ such that μ is the critical point of a total-on- $\mathcal{M}_\xi^*$ extender from the $\mathcal{M}_\xi^*$ sequence.*

Proof of Claim 1. If $\gamma = \eta$, then E_η is on the $\mathcal{M}_\xi^*$ sequence, and so we may take $\mu = \kappa$. So suppose that $\gamma < \eta$. Then we have that $\kappa < \text{lh}(E_\gamma)$, and $\text{lh}(E_\gamma)$ is a cardinal of $\mathcal{M}_\eta$, and $\mathcal{M}_\xi^*$ and $\mathcal{M}_\eta$ agree below $\text{lh}(E_\gamma)$. If $\text{crit}(E_\gamma) \leq \kappa < \text{lh}(E_\gamma)$, then it follows that E_γ

is total-on- $\mathcal{M}_\xi^*$ and so we may take $\mu = \text{crit}(E_\gamma^\mathcal{T})$. So suppose $\kappa < \text{crit } E_\gamma$. Then we have that $(\kappa^{++})^{\mathcal{M}_\eta} < \text{crit}(E_\gamma)$. By the initial segment condition on E_η , there is an initial segment of E_η on the $\mathcal{M}_\eta$ sequence below $\text{crit}(E_\gamma)$, and so on the M_ξ^* sequence. So we may take $\mu = \kappa$. $\diamond$

Claim 2. $\max(\text{code}(\alpha_0)) < \kappa$.

Proof of Claim 2. Let $\mu < \kappa$ be as in Claim 1. By the proof of Lemma 3.27 on page 51, since $\mathcal{M}_\xi^*$ is not explicitly $(\alpha^*, 0)$ -big for any $\alpha^* > \alpha_0$, $\max(\text{code}(\alpha_0)) < \mu$. $\diamond$

Now, suppose towards a contradiction that $\mathcal{M}_\xi$ were not $(\alpha_0, n_0 + 1)$ -petite above β . Then there is some (α', n') with $(\alpha_0, n_0) <_{\text{lex}} (\alpha', n')$, such that $\mathcal{M}_\xi$ is weakly, explicitly (α', n') -big above β . Let $s' = \text{code}(\alpha')$. Let $s_0 = \text{code}(\alpha_0)$. Then $\mathcal{M}_\xi$ satisfies the statement:

There exists an s' such that $(|s_0|, n_0) <_{\text{lex}} (|s'|, n')$, and I am weakly, explicitly $(|s'|, n_0)$ -big above β .

Since $\beta < \text{crit}(j)$, and $\max(s_0) < \text{crit}(j)$, and j is either Σ_2 -elementary, or Σ_1 -elementary and cofinal, it follows that $\mathcal{M}_\xi^*$ satisfies the same statement. Since this is a contradiction, we have that $\mathcal{M}_\xi$ is $(\alpha_0, n_0 + 1)$ -petite above β . (There is one subtle, technical point here. If we were using the notion of “ (α, n) -small” instead of the notion of “ (α, n) -petite” in this lemma, then at this point, in order to take care of one particular case, we would need to show that if $\mathcal{M}_\xi$ were admissible, then $\mathcal{M}_\xi^*$ would be admissible too. The contrapositive of this would require us to show that *non-admissibility* is preserved by ultrapowers. Recall that earlier in this paper we were required to show that admissibility was preserved by ultrapowers. We do in fact have a proof that non-admissibility is preserved by ultrapowers. However, since we are using the notion of “ (α, n) -petite” instead of the notion of “ (α, n) -small” in this lemma, we will not need this proof here.)

Finally, we must consider the case that ξ is a limit ordinal. Let $\gamma <_T \xi$ be such that $i_{\gamma, \xi}^\mathcal{T}$ is defined. Let $j = i_{\gamma, \xi}^\mathcal{T}$ and let $\kappa = \text{crit}(j)$. We may assume that γ is greatest so that $\gamma <_T \xi$ and $\kappa = \text{crit}(i_{\gamma, \xi}^\mathcal{T})$. As in Claim 1 we have that there is an ordinal $\mu \leq \kappa$ such that μ is the critical point of a total-on- $\mathcal{M}_\gamma$ extender from the $\mathcal{M}_\gamma$ sequence. Let (α_0, n_0) be the lexicographically greatest pair such that $\mathcal{M}_\gamma$ is weakly, explicitly (α_0, n_0) -big above

β . As in Claim 2 we have that $\max(\text{code}(\alpha_0)) < \kappa$. The rest of the proof is the same as the successor case. $\square$

7. A COMPARISON THEOREM

In this section we prove the following comparison theorem: If $\mathcal{M}$ and $\mathcal{N}$ are ω -mice, and $\mathcal{N}$ is $\Pi_{(\alpha,n)}$ -iterable, and $\mathcal{M}$ is fully iterable and $(\alpha, n+1)$ -petite, then $\mathcal{N} \leq \mathcal{M}$ or $\mathcal{M} \leq \mathcal{N}$. (See part (3) Theorem 7.16 below.) Our theorem is an extension of Lemma 2.2 from [St5]. We employ the same architecture here as in the proof of that lemma. Consequently, we will be borrowing many definitions and ideas from the paper [St5]. We will attempt to make this section self-contained by giving definitions (at least informally) for all of the terms we use. Having a copy of the paper [St5] will be useful to the reader, but should not be absolutely necessary. It may also prove useful to the reader to have some familiarity with the paper [St2]. We work in the theory ZFC.

We begin with a short discussion about our method of coding hereditarily countable sets by reals. Recall that if $x \in \mathbb{R}$ then x^* is the hereditarily countable set coded by x . Suppose that $\mathcal{T}$ is a countable iteration tree on a countable premouse, and $x \in \mathbb{R}$ and $x^* = \mathcal{T}$. Suppose that b is a cofinal, wellfounded branch of $\mathcal{T}$, and that b is the unique cofinal branch of $\mathcal{T}$ with some given property. Thus $\mathcal{M}_b^{\mathcal{T}}$ is definable from $\mathcal{T}$. Consider the following question. Is there a real y such that $y^* = \mathcal{M}_b^{\mathcal{T}}$ and y is definable from x ? The answer is yes. This is because, given $\mathcal{T}$ and $\mathcal{M}_b^{\mathcal{T}}$, the coding of $\mathcal{T}$ by x induces, in a simple way, a coding of $\mathcal{M}_b^{\mathcal{T}}$. Below we make this idea precise.

Definition 7.1. Let $P \subset \text{HC} \times \text{HC}$ be a binary relation on HC. Let $f : P \times \mathbb{R} \rightarrow \mathbb{R}$. Then we say that f is a *code transformation for P* iff for all $\langle a, b, x \rangle \in \text{dom}(f)$, if $x^* = a$ then $(f(a, b, x))^* = b$.

Definition 7.2. Let $P \subset \text{HC} \times \text{HC}$ be a binary relation on HC. Suppose that P is definable over HC. Then we say that P *admits a definable code transformation* iff there is a code transformation f for P such that f is definable over HC.

For example, suppose that $P(a, b) \Leftrightarrow b \subseteq a$. Then P admits a simple definable code transformation, namely, given a wellfounded tree coding a , give the subtree which codes b .

More to the point here is the following proposition.

Proposition 7.3. *Let P be the binary relation on HC defined by $P(\mathcal{T}, \langle b, \mathcal{M} \rangle) \Leftrightarrow \mathcal{T}$ is an iteration tree and b is a cofinal wellfounded branch of $\mathcal{T}$ and $\mathcal{M} = \mathcal{M}_b^{\mathcal{T}}$. Then P admits a definable code transformation.*

Proof. Suppose that $P(\mathcal{T}, \langle b, \mathcal{M} \rangle)$ holds, and $x \in \mathbb{R}$ and $x^* = \mathcal{T}$. We must define a real y such that $y^* = \langle b, \mathcal{M} \rangle$. Our definition is allowed to employ x , $\mathcal{T}$, $\mathcal{M}$, and b as parameters. We will define two reals, y_1 and y_2 , such that $y_1^* = b$ and $y_2^* = \mathcal{M}$. From y_1 and y_2 it is easy to define y .

To begin with, there is some ambiguity about exactly which hereditarily countable set $\mathcal{T}$ is. For concreteness, let us define an iteration tree $\mathcal{T}$ to be an ordered 8-tuple. That is $\mathcal{T} = \langle \alpha, T, \vec{E}, \vec{M}, \vec{M}^*, \vec{i}, D, \deg \rangle$ where $\alpha = \text{lh}(\mathcal{T})$, T is the binary relation on α which is the tree ordering associated with $\mathcal{T}$, $D \subseteq \alpha$ is the drop set associated with $\mathcal{T}$, $\deg : \alpha \rightarrow \omega + 1$ is the degree function associated with $\mathcal{T}$, $\vec{E}$ is the sequence of extenders used on $\mathcal{T}$, $\vec{M}$ is the sequence of models on $\mathcal{T}$, $\vec{M}^*$ is the sequence of models which are dropped to, and $\vec{i}$ is the sequence of embeddings associated with $\mathcal{T}$. (This definition of an iteration tree differs slightly from the one in [MiSt]. In that paper an iteration tree is defined as an ordered 5-tuple. The extra three components of $\mathcal{T}$ which we have here, α , $\vec{M}$, and $\vec{i}$, are definable from the other 5 components. Thus they are redundant. We have included them here to simplify our discussion of definable code transformations.)

Now, since $b \subseteq \alpha$ and x gives us a coding of α , it is easy to see how to define y_1 . So we concentrate on y_2 . To define a real y_2 such that $y_2^* = \mathcal{M}$, it suffices to define an enumeration $f : \omega \xrightarrow[\text{onto}]{1-1} \mathcal{M}$. Since the transitive collapse map is definable over HC, it suffices to define an enumeration of the pre-collapsed version of the direct limit model $\mathcal{M}_b^{\mathcal{T}}$. Thus it suffices to define an enumeration of the set $\bigcup_{\gamma \in b} \{\gamma\} \times \mathcal{M}_\gamma^{\mathcal{T}}$. It is easy to see that we can define such an enumeration over HC, given x , $\mathcal{T}$, and b as parameters. $\square$

We will need to borrow several notions from the paper [St5]. The first such notion is that of a *coiteration* of two premice. If $\mathcal{M}$ and $\mathcal{N}$ are premice, then a coiteration of $\mathcal{M}$ and $\mathcal{N}$ is the sequence of iteration trees that results from comparing $\mathcal{M}$ with $\mathcal{N}$ for some number of steps. The precise definition below is taken word-for-word from [St5].

Definition 7.4. Let $\mathcal{M}$ and $\mathcal{N}$ be premice. A *coiteration* of $\mathcal{M}$ and $\mathcal{N}$ is a sequence $\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \theta \rangle$ such that

- (1) $\mathcal{T}_\alpha$ and $\mathcal{U}_\alpha$ are ω -maximal iteration trees on $\mathcal{M}$ and $\mathcal{N}$ respectively.
- (2) $\alpha < \beta \implies \mathcal{T}_\beta$ extends $\mathcal{T}_\alpha$ and $\mathcal{U}_\beta$ extends $\mathcal{U}_\alpha$.
- (3) α limit $\implies \left(\mathcal{T}_\alpha = \bigcup_{\beta < \alpha} \mathcal{T}_\beta \text{ and } \mathcal{U}_\alpha = \bigcup_{\beta < \alpha} \mathcal{U}_\beta \right)$.
- (4) $\text{lh}(\mathcal{T}_\alpha)$ a limit $\implies \text{lh}(\mathcal{T}_{\alpha+1}) = \text{lh}(\mathcal{T}_\alpha) + 1$ and $\text{lh}(\mathcal{U}_\alpha)$ a limit $\implies \text{lh}(\mathcal{U}_{\alpha+1}) = \text{lh}(\mathcal{U}_\alpha) + 1$.
- (5) $\left(\text{lh}(\mathcal{T}_\alpha) \text{ a limit and } \text{lh}(\mathcal{U}_\alpha) \text{ a successor} \right) \implies \text{lh}(\mathcal{U}_{\alpha+1}) = \text{lh}(\mathcal{U}_\alpha)$, and
 $\left(\text{lh}(\mathcal{U}_\alpha) \text{ a limit and } \text{lh}(\mathcal{T}_\alpha) \text{ a successor} \right) \implies \text{lh}(\mathcal{T}_{\alpha+1}) = \text{lh}(\mathcal{T}_\alpha)$.
- (6) If $\text{lh}(\mathcal{T}_\alpha)$ and $\text{lh}(\mathcal{U}_\alpha)$ are both successor ordinals, and $\alpha + 1 < \theta$, then $\mathcal{T}_{\alpha+1}$ and $\mathcal{U}_{\alpha+1}$ are determined by “iterating the least disagreement” between the last models in $\mathcal{T}_\alpha$ and $\mathcal{U}_\alpha$, and by the rules for ω -maximal iteration trees. (See §7 of [MiSt].)

A coiteration of $\mathcal{M}$ and $\mathcal{N}$ is determined by the cofinal wellfounded branches of $\mathcal{T}_\alpha$ or $\mathcal{U}_\alpha$ used to produce $\mathcal{T}_{\alpha+1}$ or $\mathcal{U}_{\alpha+1}$ in the case $\text{lh}(\mathcal{T}_\alpha)$ or $\text{lh}(\mathcal{U}_\alpha)$ is a limit. This means that $\bigcup_{\alpha < \theta} \mathcal{T}_\alpha$ and $\bigcup_{\alpha < \theta} \mathcal{U}_\alpha$ determine $\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \theta \rangle$, so we can identify the two, and speak of an appropriate pair $(\mathcal{T}, \mathcal{U})$ of iteration trees on $\mathcal{M}$ and $\mathcal{N}$ respectively as a coiteration.

Definition 7.5. Let $(\mathcal{T}, \mathcal{U})$ be a coiteration of $\mathcal{M}$ and $\mathcal{N}$. We say that $(\mathcal{T}, \mathcal{U})$ is *terminal* iff either

- (1) $\text{lh}(\mathcal{T})$ is a limit and $\mathcal{T}$ has no cofinal wellfounded branch, or $\text{lh}(\mathcal{U})$ is a limit and $\mathcal{U}$ has no cofinal wellfounded branch, or
- (2) $\mathcal{T}$ and $\mathcal{U}$ have last models $\mathcal{P}$ and $\mathcal{Q}$, $\mathcal{P} \not\leq \mathcal{Q}$ and $\mathcal{Q} \not\leq \mathcal{P}$, and one of the ultrapowers determined by iterating the least disagreement between $\mathcal{P}$ and $\mathcal{Q}$, according to the rules for ω -maximal trees, is illfounded, or
- (3) $\mathcal{T}$ and $\mathcal{U}$ have last models $\mathcal{P}$ and $\mathcal{Q}$, and $\mathcal{P} \leq \mathcal{Q}$ or $\mathcal{Q} \leq \mathcal{P}$.

We shall call a coiteration *successful* just in case it is terminal and case (3) holds above.

The main theorems in this section and the next section include a large cardinal hypothesis—we will assume that there exists infinitely many Woodin cardinals. We will use this large cardinal hypothesis in two ways. Firstly, the hypothesis will give us the determinacy of sufficiently many games in $L(\mathbb{R})$. This determinacy is needed in order to be able to apply some of the results from earlier sections of this paper. The other way in which the large cardinal hypothesis will be useful to us is that it will give us a certain amount of *generic absoluteness*. We need this generic absoluteness in order to show that our coiterations terminate in countably many steps. (See Lemma 7.9 below.)

Both applications of our large cardinal hypothesis—determinacy and generic absoluteness—are proved using the machinery of Woodin’s *stationary tower forcing*. See Chapter 9 of [Ma2] for a thorough treatment of this machinery. Below we state two facts from the theory of stationary tower forcing. Then we will use these facts to derive the specific results we shall need.

Proposition 7.6 (Woodin). *Let $\langle \delta_n \mid n \in \omega \rangle$ be a strictly increasing sequence of Woodin cardinals. Let $\lambda = \sup_n \kappa_n$. Let $\gamma > \kappa$ be any ordinal. Then there is a generic extension of the universe, $V[G]$, such that in $V[G]$ there is an elementary embedding $j : V \rightarrow N$ such that*

- (i) γ is in the wellfounded part of N ,
- (ii) $\text{crit}(j) = \omega_1$, and $j(\omega_1) = \lambda$, and
- (iii) letting $\mathbb{R}^*$ be the reals of N , we have that $\mathbb{R}^*$ is also the set of reals in a symmetric collapse of V up to λ .

See section 9.5 of [Ma2] for a proof of this proposition.

Proposition 7.7 (Woodin). *Let $\langle \delta_n \mid n \in \omega \rangle$ be a strictly increasing sequence of Woodin cardinals. Let $\lambda = \sup_n \kappa_n$. Let $\mathbb{R}^*$ be the reals of a symmetric collapse of V up to λ . Then $L(\mathbb{R}^*) \models \text{AD}$.*

See section 9.6 of [Ma2] for a proof of this proposition.

From the existence of infinitely many Woodin cardinals, one can not prove $\text{AD}^{L(\mathbb{R})}$. However, the above two propositions do yield that *some* of the games in $L(\mathbb{R})$ are determined. For our purposes in this paper, it will be more than sufficient to know that every game in $J_\kappa(\mathbb{R})$ is determined. We give that argument next.

Corollary 7.8. *Let $\langle \delta_n \mid n \in \omega \rangle$ be a strictly increasing sequence of Woodin cardinals. Let k be the least ordinal such that $J_\kappa(\mathbb{R})$ is admissible. Then every game in $J_\kappa(\mathbb{R})$ is determined.*

Proof. Let $\lambda = \sup_n \kappa_n$. Let $\mathbb{R}'$ be the set of reals in some symmetric collapse of V up to λ . Then there is some ordinal α such that $J_\alpha(\mathbb{R}')$ is admissible. Let α_0 be the least such ordinal. Since the collapse forcing is homogeneous, α_0 does not depend on our particular choice of $\mathbb{R}'$.

Let $\gamma > \alpha_0$ be any ordinal. Let $j : V \rightarrow N$ be as in Proposition 7.6. Let $\mathbb{R}^*$ be the reals of N . By Proposition 7.6, $\mathbb{R}^*$ is also the set of reals in a symmetric collapse of V up to λ . Thus α_0 is the least α such that $J_\alpha(\mathbb{R}^*)$ is admissible. Now α_0 is in the wellfounded part of N . So $N \models$ “ α_0 is the least α such that $J_\alpha(\mathbb{R})$ is admissible.” Thus $j(\kappa) = \alpha_0$.

By Proposition 7.7 $L(\mathbb{R}^*) \models \text{AD}$, and so in particular $J_{\alpha_0}(\mathbb{R}^*) \models \text{AD}$. By the elementarity of j , $J_\kappa(\mathbb{R}) \models \text{AD}$. So every game in $J_\kappa(\mathbb{R})$ is determined. $\square$

The next lemma will tell us that our coiterations terminate in countably many steps.

Lemma 7.9. *Assume that there exists infinitely many Woodin cardinals. Let κ be the least ordinal such that $J_\kappa(\mathbb{R})$ is admissible. Let $\mathcal{M}$ and $\mathcal{N}$ be countable premice. Let $\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \theta \rangle$ be a coiteration of $\mathcal{M}$ and $\mathcal{N}$ which is in $J_\kappa(\mathbb{R})$. Then $\theta < \omega_1$.*

Proof. Suppose otherwise. Let $\mathcal{T}$ on $\mathcal{M}$ and $\mathcal{U}$ on $\mathcal{N}$ be the iteration trees of length ω_1 associated with our coiteration. For a contradiction it is enough to show that both $\mathcal{T}$ and $\mathcal{U}$ have cofinal branches. (For then we can get the usual contradiction involving stationary subsets of ω_1 . See the proof of the Comparison Theorem in [MiSt].)

As in the proof of Corollary 7.8 above, there is a generic elementary embedding $j : J_\kappa(\mathbb{R}) \rightarrow J_{\alpha_0}(\mathbb{R}^*)$ with $\text{crit}(j) = \omega_1$. As κ begins a Σ_1 -gap, $\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \theta \rangle$ is Σ_1 definable in $J_\kappa(\mathbb{R})$ from some real x .

Now $j(\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \omega_1 \rangle)$ is a coiteration of $\mathcal{M}$ and $\mathcal{N}$ in $J_{\alpha_0}(\mathbb{R}^*)$ which properly extends $\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \omega_1 \rangle$. Furthermore $j(\langle (\mathcal{T}_\alpha, \mathcal{U}_\alpha) \mid \alpha < \omega_1 \rangle)$ is Σ_1 definable in $J_{\alpha_0}(\mathbb{R}^*)$ from the real x . Thus there are cofinal branches b of $\mathcal{T}$ and c of $\mathcal{U}$ such that both b and c are definable in $L(\mathbb{R}^*)$ from x and α_0 . As the collapse forcing is homogeneous and $x \in V$, b and c are in V . $\square$

We will need to borrow a few more ideas from [St5]. First a definition.

Definition 7.10. Let $\mathcal{M}$ be a premouse and $\delta \in \text{OR}^{\mathcal{M}}$. We say that δ is a *cutpoint* of $\mathcal{M}$ iff for no extender E on the $\mathcal{M}$ -sequence do we have that $\text{crit}(E) < \delta \leq \text{lh}(E)$.

The following definition comes from [St2].

Definition 7.11. Let $\mathcal{M}$ be premouse. Then $\mathcal{M}$ is *tame* iff for all initial segments $\mathcal{N} \trianglelefteq \mathcal{M}$, and all ordinals $\delta \in \text{OR} \cap \mathcal{N}$, if δ is a Woodin cardinal of $\mathcal{N}$, then for no extender E on the $\mathcal{M}$ sequence do we have that $\text{crit}(E) \leq \delta \leq \text{lh}(E)$.

In particular, if $\mathcal{M}$ is tame and δ is a Woodin cardinal of $\mathcal{M}$, then δ is a cutpoint of $\mathcal{M}$.

Recall that in Section 1 we gave the definition of a *meek* premouse. An easy reflection argument shows that a meek premouse is tame. In particular, if $\mathcal{M}$ is premouse with the property that whenever $\mathcal{N} \trianglelefteq \mathcal{M}$, $\mathcal{N}$ does not have ω Woodin cardinals, then $\mathcal{M}$ is tame. So all of the premice we are working with in this paper are tame. The reason that we brought up the notion of tameness is that we want to highlight the following fact:

Proposition 7.12. *Suppose $\mathcal{M}$ is a tame premouse and $\mathcal{T}$ is an iteration tree on $\mathcal{M}$ of limit length and b is a cofinal wellfounded branch of $\mathcal{T}$. Let $\delta = \delta(\mathcal{T})$ and $\mathcal{Q} = Q(b, \mathcal{T})$. Suppose that $\delta \in \mathcal{Q}$. Then δ is a cutpoint of $\mathcal{Q}$.*

We continue with some more ideas from [St5] The following is Lemma 1.10 from that paper. The lemma says that if you can successfully carry out a comparison of two ω -mice $\mathcal{M}$ and $\mathcal{N}$, then in fact the comparison must have ended immediately as we already have that $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$.

Lemma 7.13. *Let $\mathcal{M}$ and $\mathcal{N}$ be δ -mice, where δ is a cutpoint of $\mathcal{M}$ and $\mathcal{N}$. Suppose that $\mathcal{J}_\delta^\mathcal{M} = \mathcal{J}_\delta^\mathcal{N}$. Let $(\mathcal{T}, \mathcal{U})$ be a coiteration of $\mathcal{M}$ and $\mathcal{N}$, with $\mathcal{P}$ the last model of $\mathcal{T}$, and $\mathcal{Q}$ the last model of $\mathcal{U}$. Suppose $\mathcal{P} \trianglelefteq \mathcal{Q}$. Then $\mathcal{M} \trianglelefteq \mathcal{N}$.*

Let us say that two premice $\mathcal{M}$ and $\mathcal{N}$ are *comparable* iff either $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$. If $\mathcal{M}$ and $\mathcal{N}$ are not comparable, then we will say that they are *incomparable*. We will also need the following, which is Lemma 1.12 from [St5].

Lemma 7.14. *Let $\mathcal{M}$ be a δ -mouse, and $\mathcal{T}$ an ω -maximal iteration tree on $\mathcal{M}$ above δ , with $\mathcal{T}$ of limit length. Let b and c be distinct cofinal wellfounded branches of $\mathcal{T}$. Then $\mathcal{M}_b^\mathcal{T}$ and $\mathcal{M}_c^\mathcal{T}$ are incomparable.*

The following lemma and its proof are implicit in [St5].

Lemma 7.15. *Let $\mathcal{M}$ be a δ -mouse, and $\mathcal{T}$ an ω -maximal iteration tree on $\mathcal{M}$ above δ , with $\mathcal{T}$ of limit length. Suppose that b and c are distinct cofinal wellfounded branches of $\mathcal{T}$. Then:*

- (a) $Q(b, \mathcal{T})$ and $Q(c, \mathcal{T})$ are incomparable.
- (b) Let $\delta' = \delta(\mathcal{T})$. Let $\mathcal{P}$ be any premouse such that $\mathcal{J}_{\delta'}^\mathcal{P} = \mathcal{J}_{\delta'}^{\mathcal{M}_b^\mathcal{T}} = \mathcal{J}_{\delta'}^{\mathcal{M}_c^\mathcal{T}}$. Let $\mathcal{R} = \mathcal{J}_\alpha^\mathcal{P}$ where α is largest so that $\delta' = \alpha$ or δ' is Woodin in $\mathcal{J}_\alpha^\mathcal{P}$. Suppose that $\mathcal{R}$ is comparable with both $Q(b, \mathcal{T})$ and $Q(c, \mathcal{T})$. Then $\mathcal{P}$ is a proper initial segment of both $Q(b, \mathcal{T})$ and $Q(c, \mathcal{T})$.

Proof. (a) Suppose for example that $Q(b, \mathcal{T}) \trianglelefteq Q(c, \mathcal{T})$. If $Q(b, \mathcal{T}) \triangleleft Q(c, \mathcal{T})$, then since $\delta(\mathcal{T})$ is Woodin in $Q(c, \mathcal{T})$, $Q(b, \mathcal{T}) = \mathcal{M}_b^\mathcal{T}$ and so $\mathcal{M}_b^\mathcal{T} \triangleleft \mathcal{M}_c^\mathcal{T}$. This contradicts Lemma 7.14 above. So we must have $Q(b, \mathcal{T}) = Q(c, \mathcal{T})$ and $Q(b, \mathcal{T}) \neq \mathcal{M}_b^\mathcal{T}$ and $Q(c, \mathcal{T}) \neq \mathcal{M}_c^\mathcal{T}$. Then $\delta(\mathcal{T})$ is Woodin in $\mathcal{M}_b^\mathcal{T}$ with respect to functions definable over $Q(b, \mathcal{T})$ (since such functions are in $\mathcal{M}_b^\mathcal{T} \cap \mathcal{M}_c^\mathcal{T}$.) But this contradicts the definition of $Q(b, \mathcal{T})$.

(b) Since $Q(b, \mathcal{T})$ and $Q(c, \mathcal{T})$ are incomparable, we must have that $\mathcal{R}$ is a proper initial segment of both $Q(b, \mathcal{T})$ and $Q(c, \mathcal{T})$. But then $\mathcal{R}$ cannot define a function witnessing the failure of the Woodiness of δ' . So $\mathcal{R} = \mathcal{P}$. $\square$

Finally, we get to the main content of this section. The notion of a k -realizable pre-mouse, and a k -realizable branch of an iteration tree are mentioned in the statement of the next theorem. See [St2] for these definitions.

Theorem 7.16. *Assume that there exists ω Woodin cardinals. Let (α, n) be such that $2 \leq \alpha < \omega_1^{\omega_1}$, and $n \geq 0$.*

- (1) *Let $\mathcal{M}$ be a countable δ -mouse which is $(\alpha, n+1)$ -petite above δ , where δ is a cutpoint of $\mathcal{M}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ -realizable via π . Let $\mathcal{T}$ be an ω -maximal iteration tree on $\mathcal{M}$ of countable limit length, with $\mathcal{T}$ above δ . Suppose $\mathcal{M}_\xi^\mathcal{T}$ is $(\pi, \mathcal{T})$ -realizable for all $\xi < \text{lh}(\mathcal{T})$. Let b be the unique cofinal $(\pi, \mathcal{T})$ -realizable branch of $\mathcal{T}$. If $n \geq 1$ then b is the unique cofinal branch c of $\mathcal{T}$ such that $\mathcal{M}_c^\mathcal{T}$ is wellfounded and $Q(c, \mathcal{T})$ is $\Pi_{(\alpha, n-1)}$ -iterable above $\delta(\mathcal{T})$. If $n = 0$ then b is the unique cofinal branch c of $\mathcal{T}$ such that $\mathcal{M}_c^\mathcal{T}$ is wellfounded and $Q(c, \mathcal{T})$ is $\Pi_{(\alpha, 0)}$ iterable above $\delta(\mathcal{T})$.*
- (2) *Let $\mathcal{M}$ be a countable δ -mouse which is $(\alpha, n+1)$ -petite above δ , where δ is a cutpoint of $\mathcal{M}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ realizable. Then $\mathcal{M}$ is $\Pi_{(\alpha, n)}$ iterable above δ .*
- (3) *Suppose $\mathcal{M}$ and $\mathcal{N}$ are countable δ -mice, where δ is a cutpoint of $\mathcal{M}$ and $\mathcal{N}$. Suppose $\mathcal{J}_\delta^\mathcal{M} = \mathcal{J}_\delta^\mathcal{N}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ realizable and $(\alpha, n+1)$ -petite above δ , and that $\mathcal{N}$ is $\Pi_{(\alpha, n)}$ iterable above δ . Then $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$.*

Note. We conjecture that the above theorem is also true if the word “petite” is replaced everywhere in the statement of the theorem with the word “small.” Unfortunately, we are unable to prove this. During the following proof, we will point out exactly where the proof of the stronger version of the theorem—the version with “small” instead of “petite”—breaks down. A proof of the stronger version of the theorem would be a very nice result. In particular, it would imply that $A_{(\alpha, n)}$ is a mouse set in the case $\text{cof}(\alpha) \leq \omega$ and $n > 0$. Currently, we are not even able to show that $A_{(2, 1)}$ is a mouse set.

Proof of the theorem. The theorem is proved by induction on (α, n) . We begin with (1) for (α, n) . Let $\mathcal{M}, \delta, \pi$ and $\mathcal{T}$ be as in (1). Let b be a cofinal $(\pi, \mathcal{T})$ realizable branch of $\mathcal{T}$; it is shown in [St2] that there is a unique such branch. First we will show that b has the

iterability property stated in (1). Then we will show that b is the unique branch with this property.

First suppose $n = 0$. Since $Q(b, \mathcal{T})$ is realizable, $Q(b, \mathcal{T})$ is length- ω iterable above $\delta(\mathcal{T})$. (See Corollary 1.9 from [St2].) This trivially implies that $Q(b, \mathcal{T})$ is $\Pi_{(\alpha,0)}$ iterable above $\delta(\mathcal{T})$.

Now suppose that $n \geq 1$. By Lemma 6.10, $\mathcal{M}_b^{\mathcal{T}}$ is $(\alpha, n+1)$ -petite above δ . If $\delta(\mathcal{T}) \notin Q(b, \mathcal{T})$ then $Q(b, \mathcal{T})$ is trivially iterable above $\delta(\mathcal{T})$. So suppose that $\delta(\mathcal{T}) \in Q(b, \mathcal{T})$. Then $Q(b, \mathcal{T}) \models \delta(\mathcal{T})$ is Woodin. By part (a) of Lemma 6.9 on page 97, $Q(b, \mathcal{T})$ is (α, n) petite above $\delta(\mathcal{T})$. So by part (2) of the theorem we are proving, for $(\alpha, n-1)$, $Q(b, \mathcal{T})$ is $\Pi_{(\alpha, n-1)}$ iterable above $\delta(\mathcal{T})$.

Now we show that b is unique. Suppose towards a contradiction that there is a second cofinal branch c of $\mathcal{T}$ such that $\mathcal{M}_c^{\mathcal{T}}$ is wellfounded, and in the case $n \geq 1$, $Q(c, \mathcal{T})$ is $\Pi_{(\alpha, n-1)}$ -iterable above $\delta(\mathcal{T})$, and in the case $n = 0$, $Q(c, \mathcal{T})$ is $\Pi_{(\alpha,0)}$ -iterable above $\delta(\mathcal{T})$.

First suppose $n \geq 1$. As we pointed out above, $Q(b, \mathcal{T})$ is (α, n) petite above $\delta(\mathcal{T})$. By part (3) of the theorem we are proving, for $(\alpha, n-1)$, we have that $Q(b, \mathcal{T}) \leq Q(c, \mathcal{T})$ or $Q(c, \mathcal{T}) \leq Q(b, \mathcal{T})$. This contradicts part (a) of Lemma 7.15.

Now suppose $n = 0$ and there is a second cofinal branch c of $\mathcal{T}$ such that $\mathcal{M}_c^{\mathcal{T}}$ is wellfounded and $Q(c, \mathcal{T})$ is $\Pi_{(\alpha,0)}$ -iterable above $\delta(\mathcal{T})$. Again we will derive a contradiction by showing that $Q(b, \mathcal{T}) \leq Q(c, \mathcal{T})$ or $Q(c, \mathcal{T}) \leq Q(b, \mathcal{T})$. If $\delta(\mathcal{T}) \notin Q(b, \mathcal{T})$ then $Q(b, \mathcal{T}) \leq Q(c, \mathcal{T})$. So suppose that $\delta(\mathcal{T}) \in Q(b, \mathcal{T})$. Then $Q(b, \mathcal{T}) \models \delta(\mathcal{T})$ is Woodin. By part (b) of Lemma 6.9 on page 97, there is some $\alpha' < \alpha$ and some $m \in \omega$ such that $Q(b, \mathcal{T})$ is $(\alpha', m+1)$ -petite above $\delta(\mathcal{T})$. Fix such an α' and an m with m even. Since m is even, $Q(c, \mathcal{T})$ is $\Pi_{(\alpha', m)}$ -iterable above $\delta(\mathcal{T})$. By part (3) of the theorem we are proving, for (α', m) , (or by [St5] in the case $\alpha' = 1$) we have that $Q(b, \mathcal{T}) \leq Q(c, \mathcal{T})$ or $Q(c, \mathcal{T}) \leq Q(b, \mathcal{T})$. Again, this contradicts part (a) of Lemma 7.15. This completes the proof of (1) for (α, n) .

Next we prove (2) for (α, n) . Let $\mathcal{M}, \delta$ be as in (2). Since $\mathcal{M}$ is realizable, $\mathcal{M}$ is length- ω iterable above δ . (See Corollary 1.9 from [St2].) If n is even this implies that $\mathcal{M}$ is $\Pi_{(\alpha, n)}$ -iterable above δ . So suppose $n \geq 1$ is odd. Let $x_0 \in \mathbb{R}$ be a mouse code for $(\mathcal{M}, \delta)$. We will now describe a winning strategy for player *II* in $G_{(\alpha, n)}(x_0)$.

Let $1 \leq k \leq n + 1$. At round k of the game we have a real x_{k-1} such that x_{k-1} is a mouse code for some $(\mathcal{M}', \delta')$. (If $k = 1$ then $(\mathcal{M}', \delta') = (\mathcal{M}, \delta)$.) We maintain by induction on k the following hypothesis, which we shall refer to using the symbol $\dagger$:

$$(\dagger)_{k-1} \quad \begin{array}{l} \mathcal{M}' \text{ is } (\alpha, n + 2 - k)\text{-petite above } \delta', \\ \text{and } \delta' \text{ is a cutpoint of } \mathcal{M}' \text{ and } \mathcal{M}' \text{ is } k(\mathcal{M}', \delta') \text{ realizable.} \end{array}$$

Notice that with $k = 1$ we do have that $(\dagger)_0$ holds.

At the start of round k , player I will play a real y_k so that y_k^* is an ω -maximal putative iteration tree $\mathcal{T}$ on $\mathcal{M}'$ above δ' . We will define a real x_k which player II should play in response. If $\mathcal{T}$ has a last wellfounded model then player I loses. So we assume otherwise. Let π be such that $\mathcal{M}'$ is $k(\mathcal{M}', \delta')$ -realizable via π . Then $\mathcal{T}$ has a maximal, wellfounded, $(\pi, \mathcal{T})$ -realizable branch. (This is proved in [St2].) Let b be such a branch of minimal sup. Let $\bar{\mathcal{T}} = \mathcal{T} \upharpoonright \sup(b)$. Then b is the unique cofinal $(\pi, \bar{\mathcal{T}})$ -realizable branch of $\bar{\mathcal{T}}$ and $\mathcal{M}_\xi^{\bar{\mathcal{T}}}$ is $(\pi, \bar{\mathcal{T}})$ -realizable for all $\xi < \text{lh}(\bar{\mathcal{T}})$. So $\mathcal{M}', \delta', \bar{\mathcal{T}}, b$ satisfy the hypotheses of part (1) of this theorem for (α, n) . So b is the unique cofinal branch c of $\bar{\mathcal{T}}$ such that $\mathcal{M}_c^{\bar{\mathcal{T}}}$ is wellfounded and $Q(c, \bar{\mathcal{T}})$ is $\Pi_{(\alpha, n-1)}$ -iterable above $\delta(\bar{\mathcal{T}})$. (Notice that $n \geq 1$ since we are assuming that n is odd.)

Let $\mathcal{Q} = Q(b, \bar{\mathcal{T}})$ and $\gamma = \delta(\bar{\mathcal{T}})$. Since $\mathcal{M}'$ is $(\alpha, n+2-k)$ -petite above δ' , by Lemma 6.10, $\mathcal{M}_b^{\bar{\mathcal{T}}}$ is $(\alpha, n+2-k)$ -petite above δ' . Then by Lemma 6.9, $\mathcal{Q}$ is $(\alpha, n+2-(k+1))$ -petite above γ . We will have player II play some x_k such that x_k is a mouse code for $(\mathcal{Q}, \gamma)$. Notice that if we can do this then we will have maintained our inductive hypotheses $(\dagger)_k$ about x_k . The main problem now is to show that there is such an x_k with $x_k \in \Delta_{(\alpha, n)}(y_k)$.

Let $\bar{y}$ be a real such that $\bar{y}^* = \bar{\mathcal{T}}$ and $\bar{y}$ is recursive in y_k . Let P be the binary relation on HC given by $P(\mathcal{T}', \langle \mathcal{Q}', \delta' \rangle) \Leftrightarrow \mathcal{T}'$ is an iteration tree of limit length, and $\delta' = \delta(\mathcal{T}')$, and there exists exactly one cofinal wellfounded branch b' of $\mathcal{T}'$ such that $\mathcal{Q}' = Q(b', \mathcal{T}')$. Then P admits a definable code transformation. (See the proof of Proposition 7.3.) Let f be a definable code transformation for P . Notice that $P(\bar{\mathcal{T}}, \langle \mathcal{Q}, \gamma \rangle)$ holds. Let $x_k = f(\bar{\mathcal{T}}, \langle \mathcal{Q}, \gamma \rangle, \bar{y})$. Then $x_k^* = (\mathcal{Q}, \gamma)$. Furthermore, x_k is the unique real x such that

$(\exists z, w, r \in \mathbb{R})$ s.t. z^* is a cofinal wellfounded branch of $\bar{y}^*$, and $w^* = Q(z^*, \bar{y}^*)$ and $r^* = \delta(\bar{y}^*)$ and $x^* = (w^*, r^*)$ and player II wins $G_{(\alpha, n-1)}(x)$ and $x = f(\bar{y}^*, x^*, \bar{y})$.

Since the relation “player II wins $G_{(\alpha, n-1)}(x)$ ” is $\Pi_{(\alpha, n-1)}$, we have that $x_k \in \Delta_{(\alpha, n)}(\bar{y})$.

Clearly, if player II plays this x_k he will not lose $G_{(\alpha, n)}(x_0)$ at round k .

Playing in this way for $n + 1$ rounds we eventually produce a real x_{n+1} . We must now show that $\exists \sigma \in J_\alpha(\mathbb{R})$ such that σ is a winning quasi-strategy for player II in the length- ω iteration game $WG(x_{n+1})$. By our induction hypothesis $(\dagger)_n$, x_n is a mouse code for some $(\mathcal{M}', \delta')$ such that $\mathcal{M}'$ is $(\alpha, 1)$ -petite above δ' . Thus x_{n+1} is a mouse code for some $(\mathcal{M}'', \delta'')$ such that $\mathcal{M}''$ is $(\alpha, 1)$ -petite above δ' , and so by Lemma 6.9, for some $\beta < \alpha$ and some $m \in \omega$, $\mathcal{M}''$ is $(\beta, m + 1)$ petite above δ'' . Let σ be the quasi-strategy for player II in the game $WG(x_{n+1})$ which dictates that whenever player I plays a tree, player II should play the unique maximal realizable branch of minimal sup. This is a winning quasi-strategy for player II , and by (1) for (β, m) , σ is definable over $J_\beta(\mathbb{R})$ and so $\sigma \in J_\alpha(\mathbb{R})$.

Finally we prove (3) for (α, n) . There are three cases: $n = 0$, $n \geq 1$ odd, and $n \geq 2$ even.

CASE 1 $n = 0$. Suppose $\mathcal{M}$ and $\mathcal{N}$ are countable δ -mice, where δ is a cutpoint of $\mathcal{M}$ and $\mathcal{N}$. Suppose $\mathcal{J}_\delta^\mathcal{M} = \mathcal{J}_\delta^\mathcal{N}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ realizable and $(\alpha, 1)$ -petite above δ , and that $\mathcal{N}$ is $\Pi_{(\alpha, 0)}$ iterable above δ . Recall that this means that $\mathcal{N}$ is rank- $\lambda(\alpha)$ iterable above δ where

$$\lambda(\alpha) = \begin{cases} \omega\alpha & \text{if } \alpha \geq \omega, \\ \omega(\alpha - 1) & \text{if } \alpha < \omega. \end{cases}$$

Suppose that $\mathcal{M} \not\leq \mathcal{N}$ and $\mathcal{N} \not\leq \mathcal{M}$. We will derive a contradiction.

Note. It is interesting to note that the proof we are giving works in both the case $\text{cof}(\alpha) \leq \omega$ and the case $\text{cof}(\alpha) > \omega$.

We define a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \theta \rangle$ of $\mathcal{M}$ and $\mathcal{N}$ by induction on γ . For this it suffices to define $\mathcal{T}_{\gamma+1}$ in the case $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal, and similarly with $\mathcal{U}_{\gamma+1}$. Fix

a $k(\mathcal{M}, \delta)$ realization π of $\mathcal{M}$. If $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal then let b be the unique cofinal $(\pi, \mathcal{T})$ realizable branch of $\mathcal{T}_\gamma$. It is shown in [St2] that there always is such a b . Let $\mathcal{T}_{\gamma+1}$ be the iteration tree extending $\mathcal{T}_\gamma$ with length $\text{lh}(\mathcal{T}_\gamma) + 1$ and such that $b = [0, \text{lh}(\mathcal{T}_\gamma)]_{\mathcal{T}_{\gamma+1}}$.

If $\text{lh}(\mathcal{U}_\gamma)$ is a limit ordinal then let c be the unique cofinal wellfounded branch of $\mathcal{U}_\gamma$ such that $Q(c, \mathcal{U}_\gamma)$ is rank- $\lambda(\alpha)$ iterable above $\delta(\mathcal{U}_\gamma)$, if there is a unique such branch. If c is defined then let $\mathcal{U}_{\gamma+1}$ be the iteration tree extending $\mathcal{U}_\gamma$ with length $\text{lh}(\mathcal{U}_\gamma) + 1$ and such that $c = [0, \text{lh}(\mathcal{U}_\gamma)]_{\mathcal{U}_{\gamma+1}}$. If c is not defined then stop the construction.

Suppose this construction produces a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \omega_1 \rangle$ of length ω_1 . By (1) for $(\alpha, 0)$ the branch b of $\mathcal{T}_\gamma$ chosen when $\text{lh}(\mathcal{T}_\gamma)$ is a limit is $\Pi_1(J_\alpha(\mathbb{R}), \{\mathcal{T}_\gamma\})$ uniformly in γ . Clearly the branch c of $\mathcal{U}_\gamma$ chosen when $\text{lh}(\mathcal{U}_\gamma)$ is a limit is $\Pi_1(J_\alpha(\mathbb{R}), \{\mathcal{U}_\gamma\})$ uniformly in γ . It follows that $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \omega_1 \rangle \in J_{\alpha+\omega}(\mathbb{R})$. This contradicts Lemma 7.9. Therefore the construction must produce a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ with $\theta < \omega_1$, such that either the construction stops at stage θ for the reason described above, or $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ is terminal. If the construction produces a terminal coiteration, and if this coiteration is successful, then by Lemma 7.13, $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$. This is a contradiction. So we assume that either the construction produces a terminal, non-successful coiteration, or the construction stops for the reason described above. This leaves us with three possibilities:

- (a) $\mathcal{U}_\theta$ has successor length and the coiteration is terminal because the next ultrapower which is prescribed by the definition of a coiteration in order to define $\mathcal{U}_{\theta+1}$ is illfounded. Or,
- (b) $\mathcal{U}_\theta$ has limit length but has no cofinal wellfounded branch c such that $Q(c, \mathcal{U}_\theta)$ is rank- $\lambda(\alpha)$ iterable above $\delta(\mathcal{U}_\theta)$. Or,
- (c) $\mathcal{U}_\theta$ has limit length and has two cofinal wellfounded branches c_1 and c_2 such that both $Q(c_1, \mathcal{U}_\theta)$ and $Q(c_2, \mathcal{U}_\theta)$ are rank- $\lambda(\alpha)$ iterable above $\delta(\mathcal{U}_\theta)$.

We will show that each of these possibilities leads to a contradiction. We begin with the easiest, which is the third.

Suppose towards a contradiction that possibility (c) holds. Let $\mathcal{Q}_1 = Q(c_1, \mathcal{U}_\theta)$ and $\mathcal{Q}_2 = Q(c_2, \mathcal{U}_\theta)$. Let $\mathcal{R}$ be the corresponding realizable Q -structure from $\mathcal{T}_\theta$. (Below we explain what we mean by this. After that we will feel free to use the phrase “the

corresponding realizable Q -structure" without further explanation.) That is, let $\mathcal{P}$ be the last model from $\mathcal{T}_\theta$, if $\mathcal{T}_\theta$ has successor length, otherwise let $\mathcal{P}$ be the direct limit model of the unique cofinal realizable branch of $\mathcal{T}_\theta$. Let $\gamma = \delta(\mathcal{U}_\theta)$. Notice that $\gamma = \delta(\mathcal{T}_\theta)$ if $\mathcal{T}_\theta$ has limit length. In any case we know that $\mathcal{J}_\gamma^{\mathcal{Q}_1} = \mathcal{J}_\gamma^{\mathcal{Q}_2} = \mathcal{J}_\gamma^{\mathcal{P}}$. Finally let $\mathcal{R} = \mathcal{J}_\xi^{\mathcal{P}}$ where ξ is largest such that $\xi = \gamma$ or $\mathcal{J}_\xi^{\mathcal{P}} \models \gamma$ is Woodin. This is what we mean by the corresponding realizable Q -structure from $\mathcal{T}_\theta$. Notice that if $\mathcal{T}_\theta$ has limit length and if b is the unique cofinal realizable branch of $\mathcal{T}_\theta$ then $\mathcal{R} = Q(b, \mathcal{T}_\theta)$. Now, since $\mathcal{M}$ is $(\alpha, 1)$ -petite above δ , by Lemma 6.10, $\mathcal{P}$ is $(\alpha, 1)$ -petite above δ , and so, by Lemma 6.9, there is an α' with $1 \leq \alpha' < \alpha$, and an $m \in \omega$, such that $\mathcal{R}$ is (α', m) -petite above γ . Fix such a pair (α', m) with m even. Since m is even and $\mathcal{Q}_1$ and $\mathcal{Q}_2$ are rank- $\lambda(\alpha)$ iterable above γ , we have that $\mathcal{Q}_1$ and $\mathcal{Q}_2$ are $\Pi_{(\alpha', m)}$ iterable above γ . So by part (3) of this theorem for (α', m) we have that $\mathcal{Q}_1 \leq \mathcal{R}$ or $\mathcal{R} \leq \mathcal{Q}_1$ and also that $\mathcal{Q}_2 \leq \mathcal{R}$ or $\mathcal{R} \leq \mathcal{Q}_2$. By part (b) of Lemma 7.15, we have that $\mathcal{P} \triangleleft \mathcal{Q}_1$. But then by Lemma 7.13, $\mathcal{M} \leq \mathcal{N}$. This is a contradiction.

Now we assume towards a contradiction that either possibility (a) or possibility (b) has occurred with our coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$. Let x_0 be a mouse code for $(\mathcal{N}, \delta)$. Since $\mathcal{N}$ is rank $\lambda(\alpha)$ iterable above δ , player *II* wins the rank- $\lambda(\alpha)$ iteration game $G^{\lambda(\alpha)}(x_0)$. Fix a winning strategy σ for player *II* in this game. If possibility (a) holds then let $\mathcal{U}$ be the putative iteration tree extending $\mathcal{U}_\theta$ whose last model is illfounded and which results from $\mathcal{U}_\theta$ by taking the ultrapower which is mentioned in possibility (a). If possibility (b) holds then let $\mathcal{U} = \mathcal{U}_\theta$. In either case let y_1 be a real such that $y_1^* = \mathcal{U}$. Notice that for each β with $1 \leq \beta < \alpha$, and each $m \in \omega$, $(\lambda(\beta) + m, y_1)$ is a legal first move for player *I* in $G^{\lambda(\alpha)}(x_0)$, and that if player *I* plays this move he does not immediately lose. For each such pair (β, m) , let $x_1^{(\beta, m)}$ be σ 's response when player *I* plays $(\lambda(\beta) + m, y_1)$ as his first move in the game $G^{\lambda(\alpha)}(x_0)$. Since σ is a winning strategy, for each pair (β, m) there is some maximal wellfounded branch c of $\mathcal{U}$ so that $(x_1^{(\beta, m)})^* = \langle Q(c, \mathcal{U} \upharpoonright \sup(c)), \delta(\mathcal{U} \upharpoonright \sup(c)) \rangle$. Let $c_{(\beta, m)}$ be such a maximal wellfounded branch. Let $\mathcal{Q}_{(\beta, m)}, \delta_{(\beta, m)}$ be such that $(x_1^{(\beta, m)})^* = \langle \mathcal{Q}_{(\beta, m)}, \delta_{(\beta, m)} \rangle$. Since σ is a winning strategy, $\mathcal{Q}_{(\beta, m)}$ is rank- $(\lambda(\beta) + m)$ iterable above $\delta_{(\beta, m)}$. In summary, for each (β, m) such that $(1, 0) \leq_{\text{lex}} (\beta, m) <_{\text{lex}} (\alpha, 0)$, we have the following:

- (i) $c_{(\beta, m)}$ is a maximal wellfounded branch of $\mathcal{U}$.

- (ii) $\mathcal{Q}_{(\beta,m)} = Q(c_{(\beta,m)}, \mathcal{U} \upharpoonright \sup(c_{(\beta,m)}))$
- (iii) $\delta_{(\beta,m)} = \delta(\mathcal{U} \upharpoonright \sup(c_{(\beta,m)}))$.
- (iv) $\mathcal{Q}_{(\beta,m)}$ is rank- $(\lambda(\beta) + m)$ iterable above $\delta_{(\beta,m)}$.

By discarding some of the $c_{(\beta,m)}$ if necessary, and then re-indexing, we may maintain (i) through (iv) above and in addition we may assume

- (v) If $(\beta, m) <_{\text{lex}} (\beta', m')$ then $\sup(c_{(\beta,m)}) \leq \sup(c_{(\beta',m')})$.

It follows from this that if $(\beta, m) <_{\text{lex}} (\beta', m')$ then $\delta_{(\beta,m)} \leq \delta_{(\beta',m')}$.

Now, for each (β, m) as above, let $\gamma_{(\beta,m)} \leq \theta$ be that stage of the coiteration construction such that $c_{(\beta,m)}$ is cofinal in $\mathcal{U}_{\gamma_{(\beta,m)}}$. By property (v) above, if $(\beta, m) <_{\text{lex}} (\beta', m')$ then $\gamma_{(\beta,m)} \leq \gamma_{(\beta',m')}$.

Fix (β, m) for a moment. Notice that

$$\delta_{(\beta,m)} = \delta(\mathcal{U}_{\gamma_{(\beta,m)}})$$

and

$$\mathcal{Q}_{(\beta,m)} = Q(c_{(\beta,m)}, \mathcal{U}_{\gamma_{(\beta,m)}}).$$

Now let $\mathcal{R}_{(\beta,m)}$ be the corresponding realizable Q -structure from $\mathcal{T}_{\gamma_{(\beta,m)}}$. (We explained what we meant by this phrase in the proof above that possibility (c) cannot occur.)

Claim 1. *Suppose $1 \leq \beta < \alpha$ and $m \in \omega$ and m is even. Then $\mathcal{R}_{(\beta,m)}$ is not $(\beta, m+1)$ -petite above $\delta_{(\beta,m)}$.*

Proof of Claim 1. Suppose towards a contradiction that $\mathcal{R}_{(\beta,m)}$ is $(\beta, m+1)$ -petite above $\delta_{(\beta,m)}$. Now, since $\mathcal{Q}_{(\beta,m)}$ is rank- $(\lambda(\beta) + m)$ iterable above $\delta_{(\beta,m)}$ and m is even, $\mathcal{Q}_{(\beta,m)}$ is $\Pi_{(\beta,m)}$ iterable above $\delta_{(\beta,m)}$. By part (3) of this theorem for (β, m) , $\mathcal{R}_{(\beta,m)} \trianglelefteq \mathcal{Q}_{(\beta,m)}$ or vice-versa. Now, $c_{(\beta,m)}$ is a cofinal branch of $\mathcal{U}_{\gamma_{(\beta,m)}}$, and $\mathcal{Q}_{(\beta,m)} = Q(c_{(\beta,m)}, \mathcal{U}_{\gamma_{(\beta,m)}})$. To derive our contradiction we will show that there is a second cofinal branch c of $\mathcal{U}_{\gamma_{(\beta,m)}}$ such that $Q(c, \mathcal{U}_{\gamma_{(\beta,m)}})$ is also rank- $(\lambda(\beta) + m)$ iterable above $\delta_{(\beta,m)}$. To see that such a c exists consider two cases. First suppose that $\gamma_{(\beta,m)} < \theta$. Then $c_{(\beta,m)}$ is a maximal, non-cofinal branch of $\mathcal{U}_\theta$. In this case let c be the unique cofinal branch of $\mathcal{U}_{\gamma_{(\beta,m)}}$ which is a non-maximal branch of $\mathcal{U}$. That is, let c be the branch which was chosen by our construction. Then by the definition of our construction, $Q(c, \mathcal{U}_{\gamma_{(\beta,m)}})$ is rank- $\lambda(\alpha)$ iterable above $\delta_{(\beta,m)}$. Next consider the case that $\gamma_{(\beta,m)} = \theta$. Then $c_{(\beta,m)}$ is a cofinal branch of $\mathcal{U}_\theta$ and so it

must be possibility (b) which occurred. That is, $\mathcal{U}_\theta$ has limit length but has no cofinal wellfounded branch c such that $Q(c, \mathcal{U}_\theta)$ is rank- $\lambda(\alpha)$ iterable above $\delta(\mathcal{U}_\theta)$. Now we also have that $\gamma_{(\beta', m')} = \theta$ for all (β', m') with $(\beta, m) <_{\text{lex}} (\beta', m') <_{\text{lex}} (\alpha, 0)$. So in this case we can let $c = c_{(\beta', m')}$ where (β', m') is lexicographically least such that $(\beta, m) <_{\text{lex}} (\beta', m')$ and $c_{(\beta', m')} \neq c_{(\beta, m)}$. There is such a (β', m') because otherwise $\mathcal{Q}_{(\beta, m)}$ is rank- $\lambda(\alpha)$ iterable above $\delta_{(\beta, m)} = \delta(\mathcal{U})$, and this violates the fact that possibility (b) has occurred. So in either case we have seen that there is a second cofinal branch c of $\mathcal{U}_{\gamma_{(\beta, m)}}$ such that $Q(c, \mathcal{U}_{\gamma_{(\beta, m)}})$ is rank- $(\lambda(\beta) + m)$ iterable above $\delta_{(\beta, m)}$. So exactly as with $\mathcal{Q}_{(\beta, m)}$ above, we see that $\mathcal{R}_{(\beta, m)} \trianglelefteq Q(c, \mathcal{U}_{\gamma_{(\beta, m)}})$, or vice-versa. Now, as in the proof above that possibility (c) cannot occur, we see that part (b) of Lemma 7.15 implies that at stage $\gamma_{(\beta, m)}$ we attain a successful coiteration of $\mathcal{M}$ and $\mathcal{N}$, and so by Lemma 7.13, $\mathcal{M} \trianglelefteq \mathcal{N}$. This is a contradiction. $\diamond$

If $\mathcal{T}_\theta$ has successor length, let $\mathcal{P}$ be the last model of $\mathcal{T}_\theta$. Otherwise, let $\mathcal{P}$ be the direct limit model of the unique cofinal realizable branch of $\mathcal{T}_\theta$. We will derive our contradiction by showing that $\mathcal{P}$ is not $(\alpha, 1)$ -petite above δ . As $\mathcal{M}$ is $(\alpha, 1)$ -petite above δ , this will contradict Lemma 6.10.

Claim 2. *Suppose $1 \leq \beta < \alpha$ and $m \in \omega$. Then $\mathcal{R}_{(\beta, m)} \trianglelefteq \mathcal{P}$.*

Proof of Claim 2. Let $\mathcal{W}$ be the last model model from $\mathcal{T}_{\gamma_{(\beta, m)}}$, if $\mathcal{T}_{\gamma_{(\beta, m)}}$ has successor length, otherwise, let $\mathcal{W}$ be the direct limit model of the unique cofinal realizable branch of $\mathcal{T}_{\gamma_{(\beta, m)}}$. By definition of $\mathcal{R}_{(\beta, m)}$, $\mathcal{R}_{(\beta, m)} \trianglelefteq \mathcal{W}$. If $\gamma_{(\beta, m)} = \theta$ then $\mathcal{W} = \mathcal{P}$ and so $\mathcal{R}_{(\beta, m)} \trianglelefteq \mathcal{P}$. So suppose that $\gamma_{(\beta, m)} < \theta$. Then $\mathcal{W}$ is a model on the tree $\mathcal{T}_\theta$.

Now, let c be the unique cofinal branch of $\mathcal{U}_{\gamma_{(\beta, m)}}$ which is a non-maximal branch of $\mathcal{U}_\theta$. That is, let c be the branch which was chosen by our construction. Then by the definition of our construction, $Q(c, \mathcal{U}_{\gamma_{(\beta, m)}})$ is rank- $\lambda(\alpha)$ iterable above $\delta_{(\beta, m)}$. Let $\mathcal{S} = Q(c, \mathcal{U}_{\gamma_{(\beta, m)}})$. Since $\mathcal{R}_{(\beta, m)}$ is $(\alpha, 1)$ -petite above δ , by Lemma 6.9, there is some $\beta' < \alpha$ and some $m' \in \omega$ such that $\mathcal{R}_{(\beta, m)}$ is $(\beta', m' + 1)$ -petite above $\delta_{(\beta, m)}$. Fix such a (β', m') with m' even. Since $\mathcal{S}$ is rank- $\lambda(\alpha)$ iterable above $\delta_{(\beta, m)}$ and m' is even, $\mathcal{S}$ is $\Pi_{(\beta', m')}$ iterable above $\delta_{(\beta, m)}$. Then by part (3) of this theorem for (β', m') , $\mathcal{R}_{(\beta, m)} \trianglelefteq \mathcal{S}$ or $\mathcal{S} \trianglelefteq \mathcal{R}_{(\beta, m)}$. Suppose for a moment that $\mathcal{R}_{(\beta, m)} \triangleleft \mathcal{S}$. If $\mathcal{R}_{(\beta, m)} \triangleleft \mathcal{W}$, then $\delta_{(\beta, m)}$ is Woodin in $\mathcal{W}$ with respect to functions definable over $\mathcal{R}_{(\beta, m)}$. This contradicts the definition of $\mathcal{R}_{(\beta, m)}$, and so we must

have that $\mathcal{R}_{(\beta,m)} = \mathcal{W}$. So $\mathcal{W} \triangleleft \mathcal{S}$, and so by Lemma 7.13, $\mathcal{M} \trianglelefteq \mathcal{N}$. This is a contradiction. So $\mathcal{R}_{(\beta,m)} \not\triangleleft \mathcal{S}$. Similarly, $\mathcal{S} \not\triangleleft \mathcal{R}_{(\beta,m)}$. Thus $\mathcal{R}_{(\beta,m)} = \mathcal{S}$.

Now let ξ be such that $\mathcal{W} = \mathcal{M}_\xi^{\mathcal{T}_\theta}$. If $\mathcal{W}$ is the last model of $\mathcal{T}_\theta$, then $\mathcal{W} = \mathcal{P}$ and so $\mathcal{R}_{(\beta,m)} \trianglelefteq \mathcal{P}$. So suppose that $\mathcal{W}$ is not the last model of $\mathcal{T}_\theta$. Then the extender $E_\xi^{\mathcal{T}_\theta}$ is defined. By the definition of a coiteration, $\text{lh}(E_\xi^{\mathcal{T}_\theta})$ is the least ordinal η such that $\mathcal{J}_\eta^{\mathcal{W}} \neq \mathcal{J}_\eta^{\mathcal{N}_c}$. Since $\mathcal{R}_{(\beta,m)} = \mathcal{S}$, $\text{lh}(E_\xi^{\mathcal{T}_\theta}) > \text{OR} \cap \mathcal{R}_{(\beta,m)}$. By the properties of an iteration tree, $\mathcal{W}$ and $\mathcal{P}$ agree below the length of this extender. Thus $\mathcal{R}_{(\beta,m)} \triangleleft \mathcal{P}$. $\diamond$

One consequence of Claim 2 is that for all $(\beta, m), (\beta', m')$, we have that $\mathcal{R}_{(\beta,m)} \trianglelefteq \mathcal{R}_{(\beta',m')}$ or $\mathcal{R}_{(\beta',m')} \trianglelefteq \mathcal{R}_{(\beta,m)}$. If $\gamma_{(\beta,m)} = \gamma_{(\beta',m')}$, then by definition $\delta_{(\beta,m)} = \delta_{(\beta',m')}$ and $\mathcal{R}_{(\beta,m)} = \mathcal{R}_{(\beta',m')}$. We also have the following.

Claim 3. *Suppose (β, m) and (β', m') are such that $\gamma_{(\beta,m)} < \gamma_{(\beta',m')} < \theta$. Then $\mathcal{R}_{(\beta,m)} \triangleleft \mathcal{R}_{(\beta',m')}$ and $\text{OR} \cap \mathcal{R}_{(\beta,m)} < \delta_{(\beta',m')}$ and $\delta_{(\beta,m)}$ is a cardinal in $\mathcal{R}_{(\beta',m')}$.*

Proof of Claim 3. Fix $(\beta, m), (\beta', m')$ as in the statement of the claim. Let c be the unique cofinal branch of $\mathcal{U}_{\gamma_{(\beta,m)}}$ which is a non-maximal branch of $\mathcal{U}_\theta$. That is, let c be the branch which was chosen by our construction in order to define $\mathcal{U}_{\gamma_{(\beta,m)}+1}$. Let $\mathcal{S} = Q(c, \mathcal{U}_{\gamma_{(\beta,m)}})$. Let $\mathcal{H} = \mathcal{N}_c$. Then $\mathcal{H}$ is the last model of $\mathcal{U}_{\gamma_{(\beta,m)}+1}$. Also $\mathcal{H}$ is on the tree $\mathcal{U}_\theta$. Let ξ be such that $\mathcal{H} = \mathcal{N}_\xi^{\mathcal{U}_\theta}$. $\mathcal{H}$ is not the last model of $\mathcal{U}_\theta$ and so the extender $E_\xi^{\mathcal{U}_\theta}$ is defined. As in the proof of Claim 2 we see that $\mathcal{R}_{(\beta,m)} = \mathcal{S}$, and so $\text{lh}(E_\xi^{\mathcal{U}_\theta}) > \text{OR} \cap \mathcal{S}$. Let $\lambda = \text{lh}(E_\xi^{\mathcal{U}_\theta})$.

Now let c' be the unique cofinal branch of $\mathcal{U}_{\gamma_{(\beta',m')}}$ which is a non-maximal branch of $\mathcal{U}_\theta$. That is, let c' be the branch which was chosen by our construction in order to define $\mathcal{U}_{\gamma_{(\beta',m')}+1}$. Let $\mathcal{S}' = Q(c', \mathcal{U}_{\gamma_{(\beta',m')}})$. Let $\mathcal{H}' = \mathcal{N}_{c'}$. Then $\mathcal{H}'$ is the last model of $\mathcal{U}_{\gamma_{(\beta',m')}+1}$. Also $\mathcal{H}'$ is on the tree $\mathcal{U}_\theta$. As in the proof of Claim 2 we see that $\mathcal{R}_{(\beta',m')} = \mathcal{S}'$.

Let ξ' be such that $\mathcal{H}' = \mathcal{N}_{\xi'}^{\mathcal{U}_\theta}$. Clearly $\xi < \xi'$. So by the properties of an iteration tree we have that $J_\lambda^{\mathcal{H}} = J_\lambda^{\mathcal{H}'}$ and λ is a cardinal in $\mathcal{H}'$. Also $\lambda < \delta_{(\beta',m')}$ and so $J_\lambda^{\mathcal{H}} \triangleleft \mathcal{S}'$. So we have $\mathcal{R}_{(\beta,m)} = \mathcal{S} \triangleleft J_\lambda^{\mathcal{H}} \triangleleft \mathcal{S}' = \mathcal{R}_{(\beta',m')}$. Also, λ is a cardinal in $\mathcal{S}' = \mathcal{R}_{(\beta',m')}$ and $\delta_{(\beta,m)}$ is a cardinal in $\mathcal{H}$, so $\delta_{(\beta,m)}$ is a cardinal in $J_\lambda^{\mathcal{H}} = J_\lambda^{\mathcal{S}'}$ so $\delta_{(\beta,m)}$ is a cardinal in $\mathcal{R}_{(\beta',m')}$. $\diamond$

To obtain our contradiction, we will show that $\mathcal{P}$ is not $(\alpha, 1)$ -petite above δ . Our proof of this fact has two cases.

Suppose first that there is some (β_0, m_0) such that for all (β, m) with $(\beta_0, m_0) \leq_{\text{lex}} (\beta, m) <_{\text{lex}} (\alpha, 0)$, we have that $\gamma_{(\beta, m)} = \gamma_{(\beta_0, m_0)}$. That is, suppose that all but boundedly many of our branches are cofinal in the same initial segment of $\mathcal{U}_\theta$. (Notice that this first case *must* be true if $\text{cof}(\alpha) > \omega$. This is because θ is a countable ordinal.) Then for all sufficiently large (β, m) , $\delta_{(\beta_0, m_0)} = \delta_{(\beta, m)}$, and $\mathcal{R}_{(\beta_0, m_0)} = \mathcal{R}_{(\beta, m)}$. Write $\mathcal{R} = \mathcal{R}_{(\beta_0, m_0)}$ and $\delta' = \delta_{(\beta_0, m_0)}$. Then δ' is Woodin in $\mathcal{R}$ and, by Claim 1, for all (β, m) with $(\beta_0, m_0) \leq_{\text{lex}} (\beta, m) <_{\text{lex}} (\alpha, 0)$, $\mathcal{R}$ is not (β, m) -petite above δ' . By part (b) of Lemma 6.9, $\mathcal{R}$ is not $(\alpha, 1)$ -petite above δ . By Claim 2, $\mathcal{R} \leq \mathcal{P}$. Thus $\mathcal{P}$ is not $(\alpha, 1)$ -petite above δ . This is a contradiction.

Note. Notice that in the situation described in the previous paragraph, if $\text{cof}(\alpha) \leq \omega$ then we have something stronger than the fact that $\mathcal{R}$ is not $(\alpha, 1)$ -petite above δ . We actually have that $\mathcal{R}$ is not $(\alpha, 1)$ -small above δ . This is relevant because we conjecture that the theorem we are currently proving is also true if the word “petite” is replaced by the word “small” in the statement of the theorem. Up to this point in the proof, we have not made any use of the distinction between “small” and “petite.” We shall do so in the next paragraph.

Next suppose that there is no such pair (β_0, m_0) as above. As we pointed out above, this case can only occur if $\text{cof}(\alpha) \leq \omega$. Then we can choose an ω -sequence of pairs $(\beta_0, m_0) <_{\text{lex}} (\beta_1, m_1) <_{\text{lex}} (\beta_2, m_2) <_{\text{lex}} \cdots$ such that $\sup \{ (\beta_i, m_i) \mid i \in \omega \} = (\alpha, 0)$, and such that $\gamma_{(\beta_i, m_i)} < \gamma_{(\beta_{i+1}, m_{i+1})}$. Let $\eta = \sup_i \delta_{(\beta_i, m_i)}$. It follows from Claims 2 and 3 that for each i , $\delta_{(\beta_i, m_i)}$ is a cardinal in $J_\eta^\mathcal{P}$, and $\mathcal{R}_{(\beta_i, m_i)}$ is a proper initial segment of $J_\eta^\mathcal{P}$. Let us write $\mathcal{R}_i = \mathcal{R}_{(\beta_i, m_i)}$ and $\delta_i = \delta_{(\beta_i, m_i)}$. If there is some i such that $\mathcal{R}_i$ is not $(\alpha, 1)$ -petite above δ_i , then we have our contradiction. So suppose otherwise. Then, since $\mathcal{R}_i$ is not (β_i, m_i) -petite above δ_i , there must be some (β'_i, m'_i) with $(\beta_i, m_i) \leq_{\text{lex}} (\beta'_i, m'_i) <_{\text{lex}} (\alpha, 0)$ such that $\mathcal{R}_i$ is weakly, explicitly (β'_i, m'_i) -big above δ_i . Let κ be the least measurable cardinal of $J_\eta^\mathcal{P}$ greater than δ . Then $\delta < \kappa < \delta_0$. (In fact δ_0 is a limit of measurable cardinals in $J_\eta^\mathcal{P}$ since δ_0 is a cardinal in $J_\eta^\mathcal{P}$ and δ_0 is Woodin in $\mathcal{R}_0$.) By the proof of Lemma 3.27, we may assume that $\max(\text{code}(\beta'_i)) < \kappa$ for all i . Then, by Lemma 6.5, we have that $\mathcal{R}_{i+1}$ is *strongly* explicitly (β'_i, m'_i) -big above δ_{i+1} . Thus, δ_{i+1} is locally (β'_i, m'_i) -Woodin in $J_\eta^\mathcal{P}$. Since the δ_i are cofinal in the ordinals of $J_\eta^\mathcal{P}$ and the (β'_i, m'_i) are

cofinal in $(\alpha, 0)$, it follows that $J_\eta^{\mathcal{P}}$ is weakly, explicitly $(\alpha, 1)$ -big above δ . Thus $\mathcal{P}$ is not $(\alpha, 1)$ -petite above δ . As this is a contradiction, this completes the proof of (3) for (α, n) in the case that $n = 0$.

Note. Let us consider for a moment how we might prove the current theorem if it were stated with the word “small” instead of the word “petite.” The previous paragraph is the only place in the proof where we use the distinction between “small” and “petite”. In the previous paragraph we showed that $J_\eta^{\mathcal{P}}$ is not $(\alpha, 1)$ -petite above δ , and so $\mathcal{P}$ is not $(\alpha, 1)$ -petite above δ . This is a contradiction since we assumed that $\mathcal{M}$ is $(\alpha, 1)$ -petite above δ . We did not show that $\mathcal{P}$ is not $(\alpha, 1)$ -small above δ , and so if we were only assuming that $\mathcal{M}$ was $(\alpha, 1)$ -small above δ , we would not yet have a contradiction. One possible strategy for proving the stronger version of the current theorem—the version with “small” instead of “petite”—is to now get a contradiction by showing that our coiteration is successful. For a moment, suppose that we are assuming only that $\mathcal{M}$ is $(\alpha, 1)$ -small above δ , rather than making the stronger assumption that $\mathcal{M}$ is $(\alpha, 1)$ -petite above δ . Using the notation from the previous paragraph, let $\gamma = \sup \{ \gamma_{(\beta_i, m_i)} \mid i \in \omega \}$. Then $\gamma \leq \theta$, and $\text{lh}(\mathcal{U}_\gamma)$ is a limit ordinal. Notice that $\sup \{ \delta_i \mid i \in \omega \} = \text{OR} \cap J_\eta^{\mathcal{P}} = \eta = \delta(\mathcal{U}_\gamma)$. Suppose that $\mathcal{U}_\gamma$ has a cofinal, wellfounded branch, c . (This is true if $\gamma < \theta$.) Let $\mathcal{H} = \mathcal{N}_c^{\mathcal{U}_\gamma}$. Notice that each δ_i is a cardinal in $\mathcal{H}$. Notice also that $\mathcal{J}_\eta^{\mathcal{P}} = \mathcal{J}_\eta^{\mathcal{H}}$. If $\mathcal{H}$ is at least as tall as the first admissible structure over $J_\eta^{\mathcal{P}}$, then since $\mathcal{P}$ is $(\alpha, 1)$ -small above δ , we must have that $\mathcal{P} \triangleleft \mathcal{H}$, and so our coiteration is successful. If $\mathcal{H}$ is not as tall as the first admissible structure over $J_\eta^{\mathcal{P}}$, then $\text{OR} \cap \mathcal{H}$ is less than the least $\xi > \eta$ such that ξ is the index of an extender on the $\mathcal{P}$ -sequence. So again we have $\mathcal{H} \trianglelefteq \mathcal{P}$ or $\mathcal{P} \trianglelefteq \mathcal{H}$. So our coiteration is successful, and this is a contradiction.

Notice that in the previous paragraph, we don’t need that $\mathcal{H}$ is fully wellfounded, but only that η is in the wellfounded part of $\mathcal{H}$. For then if $\mathcal{H}$ is not wellfounded, then the wellfounded part of $\mathcal{H}$ is at least as tall as the first admissible over $J_\eta^{\mathcal{P}}$, and so $\mathcal{P} \triangleleft \mathcal{H}$.

So our inability to prove the stronger version of the current theorem—the version with “small” instead of “petite”—boils down to our inability to prove that if $\gamma = \theta$, then $\mathcal{U}_\theta$ has a cofinal branch c , that this branch c has only boundedly many drops so that the direct limit model $\mathcal{N}_c^{\mathcal{U}_\theta}$ is defined, and that η is in the wellfounded part of $\mathcal{N}_c^{\mathcal{U}_\theta}$. Actually,

it is not too hard to see that $\mathcal{U}_\theta$ does have a cofinal branch c . If c has only boundedly many drops so that the direct limit model $\mathcal{N}_c^{\mathcal{U}_\theta}$ is defined, then it is not too hard to see that η is in the wellfounded part of $\mathcal{N}_c^{\mathcal{U}_\theta}$. So the problem boils down to showing that c has only boundedly many drops. This is what we are unable to do!

CASE 2 $n \geq 1$ is odd. Suppose $\mathcal{M}$ and $\mathcal{N}$ are countable δ -mice, where δ is a cutpoint of $\mathcal{M}$ and $\mathcal{N}$. Suppose $\mathcal{J}_\delta^{\mathcal{M}} = \mathcal{J}_\delta^{\mathcal{N}}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ realizable and $(\alpha, n+1)$ -petite above δ , and that $\mathcal{N}$ is $\Pi_{(\alpha, n)}$ iterable above δ . Suppose that $\mathcal{M} \not\leq \mathcal{N}$ and $\mathcal{N} \not\leq \mathcal{M}$. We will derive a contradiction.

Since $\mathcal{N}$ is $\Pi_{(\alpha, n)}$ iterable above δ and n is odd, we actually have that $\mathcal{N}$ is length- ω iterable above δ .

Again we define a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \theta \rangle$ of $\mathcal{M}$ and $\mathcal{N}$ by induction on γ . For this it suffices to define $\mathcal{T}_{\gamma+1}$ in the case $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal, and similarly with $\mathcal{U}_{\gamma+1}$. Fix a $k(\mathcal{M}, \delta)$ realization π of $\mathcal{M}$. If $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal then let b be the unique cofinal $(\pi, \mathcal{T})$ realizable branch of $\mathcal{T}_\gamma$. It is shown in [St2] that there always is such a b . Let $\mathcal{T}_{\gamma+1}$ be the iteration tree extending $\mathcal{T}_\gamma$ with length $\text{lh}(\mathcal{T}_\gamma) + 1$ and such that $b = [0, \text{lh}(\mathcal{T}_\gamma)]_{\mathcal{T}_{\gamma+1}}$.

If $\text{lh}(\mathcal{U}_\gamma)$ is a limit ordinal then let c be the unique cofinal wellfounded branch of $\mathcal{U}_\gamma$ such that $Q(c, \mathcal{U}_\gamma)$ is $\Pi_{(\alpha, n-1)}$ iterable above $\delta(\mathcal{U}_\gamma)$, if there is a unique such branch. If c is defined then let $\mathcal{U}_{\gamma+1}$ be the iteration tree extending $\mathcal{U}_\gamma$ with length $\text{lh}(\mathcal{U}_\gamma) + 1$ and such that $c = [0, \text{lh}(\mathcal{U}_\gamma)]_{\mathcal{U}_{\gamma+1}}$. If c is not defined then stop the construction. Note that there always is at least one cofinal wellfounded branch c' of $\mathcal{U}_\gamma$ such that $Q(c', \mathcal{U}_\gamma)$ is $\Pi_{(\alpha, n-1)}$ iterable above $\delta(\mathcal{U}_\gamma)$. (Proof: We are assuming that $\mathcal{N}$ is length- ω iterable above δ . As $n-1$ is even, length- ω iterability implies $\Pi_{(\alpha, n-1)}$ iterability. This implies that for all limit $\gamma' < \gamma$, we actually chose the unique length- ω iterable branch of $\mathcal{U}_{\gamma'}$. This implies that there is a cofinal branch c' of $\mathcal{U}_\gamma$ such that $Q(c', \mathcal{U}_\gamma)$ is length- ω iterable above $\delta(\mathcal{U}_\gamma)$.) So the only way that the construction can stop is if there are two such branches.

The construction cannot produce a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \omega_1 \rangle$ of length ω_1 . This is because, by part (a) of this theorem for (α, n) , such a coiteration would be definable over

$J_\alpha(\mathbb{R})$, contradicting Lemma 7.9. Therefore the construction must produce a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ with $\theta < \omega_1$, such that either the construction stops at stage θ for the reason described above, or $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ is terminal. If the construction produces a terminal coiteration, and if this coiteration is successful, then by Lemma 7.13, $\mathcal{M} \trianglelefteq \mathcal{N}$ or $\mathcal{N} \trianglelefteq \mathcal{M}$. This is a contradiction. So we assume that the construction does not produce a successful coiteration. But our construction makes it impossible to produce a non-successful, terminal coiteration. So the only possibility left is that the construction stopped for the reason described above. That is: $\mathcal{U}_\theta$ has limit length and has two cofinal wellfounded branches c_1 and c_2 such that both $Q(c_1, \mathcal{U}_\theta)$ and $Q(c_2, \mathcal{U}_\theta)$ are $\Pi_{(\alpha, n-1)}$ iterable above $\delta(\mathcal{U}_\theta)$. Let $\mathcal{Q}_1 = Q(c_1, \mathcal{U}_\theta)$ and $\mathcal{Q}_2 = Q(c_2, \mathcal{U}_\theta)$. Let $\mathcal{R}$ be the corresponding realizable Q -structure from $\mathcal{T}_\theta$. Let $\gamma = \delta(\mathcal{U}_\theta)$. Since $\mathcal{M}$ is (α, n) -petite above δ , $\mathcal{R}$ is $(\alpha, n-1)$ -petite above γ . So by part (3) of this theorem for $(\alpha, n-1)$ we have that $\mathcal{Q}_1 \trianglelefteq \mathcal{R}$ or $\mathcal{R} \trianglelefteq \mathcal{Q}_1$ and also that $\mathcal{Q}_2 \trianglelefteq \mathcal{R}$ or $\mathcal{R} \trianglelefteq \mathcal{Q}_2$. As in the proof that possibility (c) cannot occur, in the proof of Case 1 above, this is a contradiction. This completes the proof of (3) for (α, n) in the case that $n \geq 1$ is odd.

CASE 3 $n \geq 2$ is even. Suppose $\mathcal{M}$ and $\mathcal{N}$ are countable δ -mice, where δ is a cutpoint of $\mathcal{M}$ and $\mathcal{N}$. Suppose $\mathcal{J}_\delta^\mathcal{M} = \mathcal{J}_\delta^\mathcal{N}$. Suppose $\mathcal{M}$ is $k(\mathcal{M}, \delta)$ realizable and $(\alpha, n+1)$ -petite above δ , and that $\mathcal{N}$ is $\Pi_{(\alpha, n)}$ iterable above δ . Suppose that $\mathcal{M} \not\trianglelefteq \mathcal{N}$ and $\mathcal{N} \not\trianglelefteq \mathcal{M}$. We will derive a contradiction.

Fix a mouse code x_0 for $(\mathcal{N}, \delta)$. Again we define a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \theta \rangle$ of $\mathcal{M}$ and $\mathcal{N}$ by induction on γ . For this it suffices to define $\mathcal{T}_{\gamma+1}$ in the case $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal, and similarly with $\mathcal{U}_{\gamma+1}$. Fix a $k(\mathcal{M}, \delta)$ realization π of $\mathcal{M}$. If $\text{lh}(\mathcal{T}_\gamma)$ is a limit ordinal then let b be the unique cofinal $(\pi, \mathcal{T})$ realizable branch of $\mathcal{T}_\gamma$. We know that there always is such a b . Let $\mathcal{T}_{\gamma+1}$ be the iteration tree extending $\mathcal{T}_\gamma$ with length $\text{lh}(\mathcal{T}_\gamma) + 1$ and such that $b = [0, \text{lh}(\mathcal{T}_\gamma)]_{\mathcal{T}_{\gamma+1}}$.

Now suppose that $\text{lh}(\mathcal{U}_\gamma)$ is a limit ordinal. Let $\mathcal{R}_\gamma$ be the corresponding realizable Q -structure from $\mathcal{T}_\gamma$. As $\mathcal{N}$ is $\Pi_{(\alpha, n)}$ iterable above δ , player *II* wins the game $G_{(\alpha, n)}(x_0)$. Let A_γ be the set of all reals x_1 such that:

$(\exists y_1, z_1 \in \mathbb{R})$ s.t. $y_1^* = \mathcal{U}_\gamma$ and $z_1^* = \mathcal{R}_\gamma$ and $\langle y_1, z_1, x_1 \rangle$ is a winning position for player *II* in $G_{(\alpha, n)}(x_0)$.

Now let c be the unique cofinal wellfounded branch of $\mathcal{U}_\gamma$ such that $(\exists x_1 \in A_\gamma)$ with $x_1^* = (Q(c, \mathcal{U}_\gamma), \delta(\mathcal{U}_\gamma))$, if there is a unique such branch. If c is defined then let $\mathcal{U}_{\gamma+1}$ be the iteration tree extending $\mathcal{U}_\gamma$ with length $\text{lh}(\mathcal{U}_\gamma) + 1$ and such that $c = [0, \text{lh}(\mathcal{U}_\gamma)]_{\mathcal{U}_{\gamma+1}}$. If c is not defined then stop the construction.

The construction cannot produce a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma < \omega_1 \rangle$ of length ω_1 . This is because, by part (a) of this theorem for (α, n) , such a coiteration would be definable over $J_\alpha(\mathbb{R})$ contradicting Lemma 7.9. Therefore the construction must produce a coiteration $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ with $\theta < \omega_1$, such that either the construction stops at stage θ for the reason described above, or $\langle (\mathcal{T}_\gamma, \mathcal{U}_\gamma) \mid \gamma \leq \theta \rangle$ is terminal. If the construction produces a terminal coiteration, and if this coiteration is successful, then by Lemma 7.13, $\mathcal{M} \leq \mathcal{N}$ or $\mathcal{N} \leq \mathcal{M}$. This is a contradiction. So we assume that either the construction produces a terminal, non-successful coiteration, or the construction stops for the reason described above. This leaves us with three possibilities:

- (a) $\mathcal{U}_\theta$ has successor length and the coiteration is terminal because the next ultrapower which is prescribed by the definition of a coiteration in order to define $\mathcal{U}_{\theta+1}$ is illfounded. Or,
- (b) $\mathcal{U}_\theta$ has limit length but has no cofinal wellfounded branch c such that $(\exists x_1 \in A_\theta)$ with $x_1^* = (Q(c, \mathcal{U}_\theta), \delta(\mathcal{U}_\theta))$. Or,
- (c) $\mathcal{U}_\theta$ has limit length and has two cofinal wellfounded branches c_1 and c_2 such that both c_1 and c_2 satisfy the conditions on c in possibility (b) above.

We will show that each of these possibilities leads to a contradiction.

First suppose that either possibility (a) or possibility (b) holds. If possibility (a) holds then let $\mathcal{U}$ be the putative iteration tree extending $\mathcal{U}_\theta$ whose last model is illfounded and which results from $\mathcal{U}_\theta$ by taking the ultrapower which is mentioned in possibility (a). If possibility (b) holds then let $\mathcal{U} = \mathcal{U}_\theta$. In either case let y_1 be a real such that $y_1^* = \mathcal{U}$.

Now, if $\mathcal{T}_\theta$ has limit length, then let b be the unique cofinal realizable branch of $\mathcal{T}_\theta$ and let $X = Q(b, \mathcal{T}_\theta)$. Otherwise let $X = \emptyset$. In either case let z_1 be any real such that $z_1^* =$ the transitive closure of $\{\mathcal{T}_\theta, X\}$.

Notice that $\langle y_1, z_1 \rangle$ is a legal first move for player I in $G_{(\alpha, n)}(x_0)$, and that if player I plays this move he does not immediately lose. As player II wins $G_{(\alpha, n)}(x_0)$ there is a real x_1 such that $\langle y_1, z_1, x_1 \rangle$ is a winning position for player II . Fix such an x_1 . Then there is some maximal wellfounded branch c of $\mathcal{U}$ so that $x_1^* = \langle Q(c, \mathcal{U} \upharpoonright \sup(c)), \delta(\mathcal{U} \upharpoonright \sup(c)) \rangle$.

First we will show that c is not a cofinal branch of $\mathcal{U}$. Suppose towards a contradiction that c is a cofinal branch of $\mathcal{U}$. Then $\mathcal{U}$ has limit length and so $\mathcal{U} = \mathcal{U}_\theta$. Let $\mathcal{R}_\theta$ be the corresponding realizable Q -structure from $\mathcal{T}_\theta$. Then $\mathcal{R}_\theta$ is in the transitive closure of $\{\mathcal{T}_\theta, X\}$. So there is a real z such that $z^* = \mathcal{R}_\theta$ and z is recursive in z_1 . Examining the rules $G_{(\alpha, n)}(x_0)$ for even n we see that $\langle y_1, z, x_1 \rangle$ is also a winning position for player II . This means that $x_1 \in A_\theta$. But this violates the hypothesis that either possibility (a) or possibility (b) occurred. So c is not a cofinal branch of $\mathcal{U}$.

So let $\gamma < \theta$ be such that c is cofinal in $\mathcal{U}_\gamma$. Then $\mathcal{U}_\gamma$ has limit length. Now $\mathcal{R}_\gamma$ is in the transitive closure of $\{\mathcal{T}_\theta, X\}$. So there is a real z such that $z^* = \mathcal{R}_\gamma$ and z is recursive in z_1 . Fix such a z . Let y be any real such that $y^* = \mathcal{U}_\gamma$. By examining the rules of $G_{(\alpha, n)}(x_0)$ for even n we see that $\langle y, z, x_1 \rangle$ is also a winning position for player II . Thus $x_1 \in A_\gamma$. Now let c' be the unique cofinal branch of $\mathcal{U}_\gamma$ which is a non-maximal branch of $\mathcal{U}_\theta$. Then c and c' witness that we should have stopped the coiteration construction at stage γ because possibility (c) occurred. This is a contradiction. So we have shown that neither possibility (a) nor possibility (b) can occur.

Finally, assume towards a contradiction that possibility (c) has occurred. Let c_1 and c_2 be two cofinal wellfounded branches of $\mathcal{U}_\theta$ such that $(\exists x_1 \in A_\theta)$ with $x_1^* = (Q(c_1, \mathcal{U}_\theta), \delta(\mathcal{U}_\theta))$, and similarly with c_2 . Let $\delta^* = \delta(\mathcal{U}_\theta)$. Let $\mathcal{Q} = Q(c_1, \mathcal{U}_\theta)$, and let $\mathcal{Q}' = Q(c_2, \mathcal{U}_\theta)$. Let $\mathcal{R}$ be the corresponding realizable Q -structure from $\mathcal{T}_\theta$. To obtain a contradiction, we will show that $\mathcal{R}$ is comparable with both $\mathcal{Q}$ and $\mathcal{Q}'$. Below we will show that $\mathcal{R}$ is comparable with $\mathcal{Q}$. The proof that $\mathcal{R}$ is comparable with $\mathcal{Q}'$ is identical.

So suppose towards a contradiction that $\mathcal{R} \not\leq \mathcal{Q}$ and $\mathcal{Q} \not\leq \mathcal{R}$. Now we define a coiteration $\langle (\bar{\mathcal{T}}_\gamma, \bar{\mathcal{U}}_\gamma) \mid \gamma < \bar{\theta} \rangle$ of $\mathcal{R}$ and $\mathcal{Q}$ by induction on γ . For this it suffices to define $\bar{\mathcal{T}}_{\gamma+1}$ in the case $\text{lh}(\bar{\mathcal{T}}_\gamma)$ is a limit ordinal, and similarly with $\bar{\mathcal{U}}_{\gamma+1}$. Fix a $k(\mathcal{R}, \delta)$ realization $\bar{\pi}$ of $\mathcal{R}$. If $\text{lh}(\bar{\mathcal{T}}_\gamma)$ is a limit ordinal then let b be the unique cofinal $(\bar{\pi}, \bar{\mathcal{T}})$ realizable branch of

$\bar{\mathcal{T}}_\gamma$. We know that there always is such a b . Let $\bar{\mathcal{T}}_{\gamma+1}$ be the iteration tree extending $\bar{\mathcal{T}}_\gamma$ with length $\text{lh}(\bar{\mathcal{T}}_\gamma) + 1$ and such that $b = [0, \text{lh}(\bar{\mathcal{T}}_\gamma)]_{\bar{\mathcal{T}}_{\gamma+1}}$.

If $\text{lh}(\bar{\mathcal{U}}_\gamma)$ is a limit ordinal then let c be the unique cofinal wellfounded branch of $\bar{\mathcal{U}}_\gamma$ such that $Q(c, \bar{\mathcal{U}}_\gamma)$ is $\Pi_{(\alpha, n-2)}$ iterable above $\delta(\bar{\mathcal{U}}_\gamma)$, if there is a unique such branch. If c is defined then let $\bar{\mathcal{U}}_{\gamma+1}$ be the iteration tree extending $\bar{\mathcal{U}}_\gamma$ with length $\text{lh}(\bar{\mathcal{U}}_\gamma) + 1$ and such that $c = [0, \text{lh}(\bar{\mathcal{U}}_\gamma)]_{\bar{\mathcal{U}}_{\gamma+1}}$. If c is not defined then stop the construction.

The construction cannot produce a coiteration $\langle (\bar{\mathcal{T}}_\gamma, \bar{\mathcal{U}}_\gamma) \mid \gamma < \omega_1 \rangle$ of length ω_1 . This is because, by part (a) of this theorem for $(\alpha, n-1)$, such a coiteration would be definable over $J_\alpha(\mathbb{R})$, contradicting Lemma 7.9. Therefore the construction must produce a coiteration $\langle (\bar{\mathcal{T}}_\gamma, \bar{\mathcal{U}}_\gamma) \mid \gamma \leq \bar{\theta} \rangle$ with $\bar{\theta} < \omega_1$, such that either the construction stops at stage $\bar{\theta}$ for the reason described above, or $\langle (\bar{\mathcal{T}}_\gamma, \bar{\mathcal{U}}_\gamma) \mid \gamma \leq \bar{\theta} \rangle$ is terminal. If the construction produces a terminal coiteration, and if this coiteration is successful, then by Lemma 7.13, $\mathcal{R} \leq \mathcal{Q}$ or $\mathcal{Q} \leq \mathcal{R}$. This is a contradiction. So we assume that either the construction produces a terminal, non-successful coiteration, or the construction stops for the reason described above. This leaves us with three possibilities:

- (i) $\bar{\mathcal{U}}_{\bar{\theta}}$ has successor length and the coiteration is terminal because the next ultrapower which is prescribed by the definition of a coiteration in order to define $\bar{\mathcal{U}}_{\bar{\theta}+1}$ is illfounded. Or,
- (ii) $\bar{\mathcal{U}}_{\bar{\theta}}$ has limit length but has no cofinal wellfounded branch c such that $Q(c, \bar{\mathcal{U}}_{\bar{\theta}})$ is $\Pi_{(\alpha, n-2)}$ iterable above $\delta(\bar{\mathcal{U}}_{\bar{\theta}})$. Or,
- (iii) $\bar{\mathcal{U}}_{\bar{\theta}}$ has limit length and has two cofinal wellfounded branches c and c' such that both $Q(c, \bar{\mathcal{U}}_{\bar{\theta}})$ and $Q(c', \bar{\mathcal{U}}_{\bar{\theta}})$ are $\Pi_{(\alpha, n-2)}$ iterable above $\delta(\bar{\mathcal{U}}_{\bar{\theta}})$

We will show that each of these possibilities leads to a contradiction. We begin with the easiest, which is the third.

Suppose towards a contradiction that possibility (iii) holds. Let $\bar{\delta} = \delta(\bar{\mathcal{U}}_{\bar{\theta}})$. Let $\mathcal{Q}_1 = Q(c, \bar{\mathcal{U}}_{\bar{\theta}})$ and $\mathcal{Q}_2 = Q(c', \bar{\mathcal{U}}_{\bar{\theta}})$. Let $\bar{\mathcal{R}}$ be the corresponding realizable Q -structure from $\bar{\mathcal{T}}_{\bar{\theta}}$. Since $\mathcal{M}$ is $(\alpha, n+1)$ -petite above δ , $\mathcal{R}$ is $(\alpha, n-1)$ -petite and above δ^* , and so $\bar{\mathcal{R}}$ is $(\alpha, n-2)$ -petite and above $\bar{\delta}$. Also $\mathcal{Q}_1$ and $\mathcal{Q}_2$ are $\Pi_{(\alpha, n-2)}$ iterable above $\bar{\delta}$. So by part (3) of this theorem for $(\alpha, n-2)$ we have that $\mathcal{Q}_1 \leq \bar{\mathcal{R}}$ or $\bar{\mathcal{R}} \leq \mathcal{Q}_1$ and also that $\mathcal{Q}_2 \leq \bar{\mathcal{R}}$

or $\bar{\mathcal{R}} \leq \mathcal{Q}_2$. This is a contradiction. (See the proof that possibility (c) cannot occur in the proof of Case 1 above.)

Now we assume towards a contradiction that either possibility (i) or possibility (ii) has occurred. Recall that by our current case hypotheses we have that $(\exists x_1 \in A_\theta)$ with $x_1^* = (\mathcal{Q}, \delta^*)$. Fix such an x_1 . By the definition of A_θ , $(\exists y_1, z_1 \in \mathbb{R})$ s.t. $y_1^* = \mathcal{U}_\theta$ and $z_1^* = \mathcal{R}$, and $\langle y_1, z_1, x_1 \rangle$ is a winning position for player *II* in $G_{(\alpha, n)}(x_0)$. (Recall that x_0 is our fixed mouse code for $(\mathcal{N}, \delta)$.) Fix such reals y_1, z_1 .

If possibility (i) holds then let $\mathcal{U}$ be the putative iteration tree extending $\bar{\mathcal{U}}_{\bar{\theta}}$ whose last model is illfounded and which results from $\bar{\mathcal{U}}_{\bar{\theta}}$ by taking the ultrapower which is mentioned in possibility (i). If possibility (ii) holds then let $\mathcal{U} = \bar{\mathcal{U}}_{\bar{\theta}}$.

Claim. *There is a real y_2 such that $y_2^* = \mathcal{U}$ and $y_2 \in \Delta_{(\alpha, n-1)}(x_1, z_1)$.*

Proof of Claim. We will prove the claim in the case that possibility (ii) has occurred, that is assuming that $\mathcal{U} = \bar{\mathcal{U}}_{\bar{\theta}}$. The proof in the case that possibility (i) has occurred is similar.

Let $\rho = \text{lh}(\mathcal{U})$. First we will show that there is a real $w \in \text{WO}$ such that $|w| = \rho$ and $w \in \Delta_{(\alpha, n-1)}(x_1, z_1)$. For $y, w \in \mathbb{R}$ put $S(y, w)$ iff y^* is an iteration tree on $\mathcal{Q}_1$, and $w \in \text{WO}$ and $|w|$ is equal to the length of the tree, and the tree results from a comparison of $\mathcal{Q}_1$ with $\mathcal{R}$ in which all branches chosen are $\Pi_{(\alpha, n-2)}$ iterable. More precisely, $S(y, w)$ iff

$w \in \text{WO}$ and there is some real v such that v^* is some coiteration $\langle (\mathcal{T}'_\gamma, \mathcal{U}'_\gamma) \mid \gamma \leq \mu \rangle$ of some two premice $\mathcal{R}', \mathcal{Q}'$ and $(x_1^*)_0 = \mathcal{Q}'$ and $z_1^* = \mathcal{R}'$ and $y^* = \mathcal{U}'_\mu$ and $|w| = \text{lh}(y^*)$ and $(\forall a, b, c, d, e \in \mathbb{R})$ if there is some $\gamma < \mu$ s.t. (either $a^* = \mathcal{T}'_\gamma$, or $a^* = \mathcal{U}'_\gamma$) and a^* has limit length and b^* is the unique cofinal branch of a^* which was chosen by the construction, and $c^* = \mathcal{Q}(b^*, a^*)$ and $d^* = \delta(a^*)$ and $e^* = (c^*, d^*)$ then player *II* wins the game $G_{(\alpha, n-2)}(e)$.

Then $S \in \Pi_{(\alpha, n-2)}(x_1, z_1)$. Now set $R(w) \Leftrightarrow (\exists y)S(y, w)$. Then $R \in \Sigma_{(\alpha, n-1)}(x_1, z_1)$ and ρ is the largest element of the set $\{|w| : R(w)\}$. Since $n-1$ is odd, we may apply Kechris's Boundedness Theorem for Subsets of ω_1 , Lemma 2.17 on page 20, to conclude that there is a $w \in \text{WO}$ such that $|w| = \rho$ and $w \in \Delta_{(\alpha, n-1)}(x_1, z_1)$. Fix such a w .

Now, let P be the binary relation on HC given by $P(\langle \mathcal{M}', \rho' \rangle, \mathcal{T}') \Leftrightarrow \mathcal{M}'$ is a premouse and $\mathcal{T}'$ is an iteration tree on $\mathcal{M}'$ of length ρ' . Then P admits a definable code transformation, f . [Proof: Suppose $P(\langle \mathcal{M}', \rho' \rangle, \mathcal{T}')$ holds. Suppose $(x')^* = \mathcal{M}'$ and $w' \in \text{WO}$ and $|w'| = \rho'$. As in the proof of Proposition 7.3, it suffices to see that we can define (over HC, using $\mathcal{M}', \rho', \mathcal{T}', x'$ and w' as parameters) a bijection from ω onto $\mathcal{T}'$. Since x' gives us a bijection from ω on $\mathcal{M}'$, and w' gives us a bijection from ω onto ρ' , it is easy to see how to define a bijection from ω onto $\mathcal{T}'$.]

Now $P(\langle \mathcal{Q}_1, \rho \rangle, \mathcal{U})$ holds. Let $y_2 = f(\langle \mathcal{Q}_1, \rho \rangle, \mathcal{U}, \langle x_1, w \rangle)$. Then $y_2^* = \mathcal{U}$ and y_2 is the unique real y such that:

$$S(y, w) \text{ and } y = f(\langle x_1^*, |w| \rangle, y^*, \langle x_1, w \rangle).$$

As $w \in \Delta_{(\alpha, n-1)}(x_1, z_1)$, we have that $y_2 \in \Delta_{(\alpha, n-1)}(x_1, z_1)$. ◇

Fix a real y_2 as in the claim. Notice that $\langle y_1, z_1, x_1, y_2 \rangle$ is a legal position for player I in $G_{(\alpha, n)}(x_0)$ which is not an immediate loss for player I . Since $\langle y_1, z_1, x_1 \rangle$ is a winning position for player II , there is a real x_2 such that $\langle y_1, z_1, x_1, y_2, x_2 \rangle$ is a winning position for player II . By examining the rules of $G_{(\alpha, n)}(x_0)$ we see that this means that there is a maximal branch c of $\mathcal{U}$ such that $x_2^* = \langle Q(c, \mathcal{U} \upharpoonright \sup(c)), \delta(\mathcal{U} \upharpoonright \sup(c)) \rangle$ and also such that player II wins the game $G_{(\alpha, n-2)}(x_2)$. The branch c can not be cofinal in $\mathcal{U}$ because this would violate our hypothesis that either possibility (i) or possibility (ii) occurred. So let $\gamma < \bar{\theta}$ be such that c is cofinal in $\bar{\mathcal{U}}_\gamma$. Then at stage γ of our coiteration we should have stopped the construction because possibility (iii) occurred. This contradiction completes the proof of Theorem 7.16. □

We conclude this section with the one and only application of the previous theorem that we shall need.

Corollary 7.17. *Assume that there exists ω Woodin cardinals. Suppose $2 \leq \alpha < \omega_1^{\omega_1}$, and $n \in \omega$. Let $\mathcal{M}$ be a countable, realizable, sound premouse. Suppose that $\mathcal{M}$ is $(\alpha, n+1)$ -petite. Then $\mathbb{R} \cap \mathcal{M} \subseteq A_{(\alpha, n+1)}$.*

Proof. Recall that $(\alpha, n+1)$ -petite means $(\alpha, n+1)$ -petite above 0, or equivalently above ω . Let $x \in \mathbb{R} \cap \mathcal{M}$. Let ξ be the rank of x in the order of construction of $\mathcal{M}$. Let $\mathcal{N} \trianglelefteq \mathcal{M}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ containing x . Then $\mathcal{N}$ projects to ω , and since $\mathcal{N}$ is sound, $\mathcal{N}$ is an ω -mouse. By part (3) of Theorem 7.16, if $\mathcal{N}'$ is any ω -mouse, and $\mathcal{N}'$ is $\Pi_{(\alpha, n)}$ iterable, then $\mathcal{N} \trianglelefteq \mathcal{N}'$ or $\mathcal{N}' \trianglelefteq \mathcal{N}$. (Notice that the requirements that $\mathcal{N}$ and $\mathcal{N}'$ agree through ω , and that ω be a cutpoint of $\mathcal{N}$ and $\mathcal{N}'$ are met trivially.) Thus, x is the unique real such that there is some $\Pi_{(\alpha, n)}$ iterable ω -mouse $\mathcal{N}'$ such that x is the ξ^{th} real in the order of construction of $\mathcal{N}'$. Let $w \in \text{WO}$ with $|w| = \xi$. Then x is the unique real x' such that:

$(\exists y, z \in \mathbb{R})$ s.t. z is a mouse code and player II wins the game $G_{(\alpha, n)}(z)$
 and $(z^*)_1 = \omega$ and $y^* = (z^*)_0$ and $x' \in y^*$ and the rank of x' in the order of
 construction of y^* is $|w|$.

Thus $x \in \Delta_{(\alpha, n+1)}(w)$. Since w was arbitrary, $x \in \Delta_{(\alpha, n+1)}(\xi)$. So $x \in A_{(\alpha, n+1)}$. $\square$

8. CONCLUSIONS

In this section we put together the results of Section 4 and Section 7 to obtain our main results.

Theorem 8.1. *Assume that there exists ω Woodin cardinals. Suppose that $(2, 0) \leq_{\text{lex}} (\alpha, n) \leq_{\text{lex}} (\omega_1^{\omega_1}, 0)$, and either $\text{cof}(\alpha) = \omega_1$, or $n = 0$. Then $A_{(\alpha, n)}$ is a mouse set.*

Proof. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is (α, n) -big, but every proper initial segment of $\mathcal{M}$ is (α, n) -small. We will show that $A_{(\alpha, n)} = \mathbb{R} \cap \mathcal{M}$.

First we will show that there exists a premouse $\mathcal{M}$ as in the previous paragraph. In fact we will show that there exists a “canonical” such $\mathcal{M}$. It is shown in [St2] that our large cardinal hypothesis implies that there is a proper class premouse $L[\vec{E}]$ such that $L[\vec{E}] \models “\exists \omega \text{ Woodin cardinals}.”$ Let $\mathcal{M}^*$ be the $\trianglelefteq$ -least initial segment of $L[\vec{E}]$ such that $\mathcal{M}^*$ has ω Woodin cardinals which are cofinal in the ordinals of $\mathcal{M}^*$. It is easy to see that $\mathcal{M}^*$ projects to ω . (This uses a simple condensation argument, similar to the proof of

Corollary 1.11.) Thus $\mathcal{M}^*$ is an ω -mouse. It follows that $\mathcal{M}^*$ is the unique $\mathcal{M}'$ such that $\mathcal{M}'$ is a realizable, ω -mouse, $\mathcal{M}' \models$ “There are ω Woodin cardinals cofinal in the ordinals,” and no proper initial segment of $\mathcal{M}'$ models this sentence. (Use Lemma 7.13 on page 105 to see this.) Thus we may think of $\mathcal{M}^*$ as the canonical inner model for the theory “There are ω Woodin cardinals cofinal in the ordinals.” By Proposition 3.25 on page 48, $\mathcal{M}^*$ is $(\omega_1^{\omega_1}, 0)$ -big. So in particular, $\mathcal{M}^*$ is (α, n) -big. Let $\mathcal{M}$ be the $\leq$ -least initial segment of $\mathcal{M}^*$ which is (α, n) -big. We may think of $\mathcal{M}$ as the “canonical” (α, n) -big mouse.

Our large cardinal hypothesis implies that every game in $J_\kappa(\mathbb{R})$ is determined, where κ is the least $\mathbb{R}$ -admissible ordinal. (See Corollary 7.8.) In particular we have that $\text{Det}(\Sigma_1(J_{\omega_1^{\omega_1}}(\mathbb{R})))$ holds. So we can apply our main correctness result, Corollary 4.7 on page 62, to conclude that $A_{(\alpha, n)} \subseteq \mathbb{R} \cap \mathcal{M}$.

Conversely, let $x \in \mathbb{R} \cap \mathcal{M}$ and we will show that $x \in A_{(\alpha, n)}$. Let $\mathcal{N}$ be the $\leq$ -least initial segment of $\mathcal{M}$ such that $x \in \mathcal{N}$. Then $\mathcal{N}$ is a proper initial segment of $\mathcal{M}$, so by hypothesis $\mathcal{N}$ is (α, n) -small.

First suppose that $n \geq 1$. Then by hypothesis $\text{cof}(\alpha) > \omega$. By definition (α, n) -petite is synonymous with (α, n) -small, so we have that $\mathcal{N}$ is (α, n) -petite. By the corollary to our main comparison theorem, Corollary 7.17 on page 124, $\mathbb{R} \cap \mathcal{N} \subseteq A_{(\alpha, n)}$. Thus $x \in A_{(\alpha, n)}$.

Now suppose that $n = 0$. Recall that

$$A_{(\alpha, 0)} = \left\{ z \in \mathbb{R} \mid (\exists \beta < \alpha) z \in \text{OD}^{J_\beta(\mathbb{R})} \right\}.$$

So we must show that $x \in \text{OD}^{J_\beta(\mathbb{R})}$ for some $\beta < \alpha$. We have that $\mathcal{N}$ is $(\alpha, 0)$ -small. By definition this is synonymous with $(\alpha, 0)$ -petite, so $\mathcal{N}$ is $(\alpha, 0)$ -petite. It follows that $\mathcal{N}$ is (β, m) -petite for some $(\beta, m) <_{\text{lex}} (\alpha, 0)$. (See Lemma 6.8 on page 95.) Fix such a (β, m) with $m \geq 1$. As in the previous paragraph we have that $\mathbb{R} \cap \mathcal{N} \subseteq A_{(\beta, m)}$. Thus $x \in A_{(\beta, m)}$. In particular, $x \in \text{OD}^{J_\beta(\mathbb{R})}$. $\square$

Remark 8.2. Let $\mathcal{M}^*$ be as in the proof above. We believe that $\mathbb{R} \cap \mathcal{M}^* = A_{(\kappa, 0)}$, where $\kappa = \kappa^{\mathbb{R}}$ is the least $\mathbb{R}$ -admissible ordinal. It would take us too far afield to prove this here, so we will save this for another paper. $A_{(\kappa, 0)}$ is an important set in descriptive set

theory. It is the set of reals which are hyperprojective in a countable ordinal. It is also the largest countable inductive set of reals.

Remark 8.3. One specific consequence of Theorem 8.1 is that for $\alpha < \omega_1^{\omega_1}$, $\text{OD}^{J_\alpha(\mathbb{R})}$ is a mouse set. This is because $\text{OD}^{J_\alpha(\mathbb{R})} = A_{(\alpha+1,0)}$.

Recall that on page 22 we discussed the relationship between the sets $A_{(\alpha,n)}$ and $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$. As promised we can now answer Questions 1 and 2 from that discussion affirmatively, at least for certain pairs (α, n) . We begin with Question 1.

Theorem 8.4. *Assume that there exists ω Woodin cardinals. Suppose that $2 < \alpha < \omega_1^{\omega_1}$, $\text{cof}(\alpha) = \omega_1$, and $n \geq 1$. Then $A_{(\alpha,n)} = \text{OD}_{n+1}^{J_\alpha(\mathbb{R})}$.*

Proof. The proof of Theorem 8.1 actually showed this, since our main correctness result, Corollary 4.7, not only gives us that $A_{(\alpha,n)} \subseteq \mathbb{R} \cap \mathcal{M}$, but actually that $\text{OD}_{n+1}^{J_\alpha(\mathbb{R})} \subseteq \mathbb{R} \cap \mathcal{M}$. $\square$

The next theorem answers Question 2 from our discussion on page 22, at least for ordinals $\alpha < \omega_1^{\omega_1}$.

Theorem 8.5. *Assume that there exists ω Woodin cardinals. Suppose that $2 \leq \alpha < \omega_1^{\omega_1}$. Then $\bigcup_n A_{(\alpha,n)} = \text{OD}^{J_\alpha(\mathbb{R})}$.*

Proof. Fix α . Notice that $\text{OD}^{J_\alpha(\mathbb{R})} = A_{(\alpha+1,0)}$. Let $\mathcal{M}$ be a countable, realize premouse such that $\mathcal{M}$ is $(\alpha+1, 0)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\alpha+1, 0)$ -small. The proof of Theorem 8.1 actually showed that $A_{(\alpha+1,0)} \subseteq \mathbb{R} \cap \mathcal{M}$, and $\mathbb{R} \cap \mathcal{M} \subseteq \bigcup_n A_{(\alpha,n)}$. $\square$

By showing that $A_{(\alpha,n)}$ is a mouse set, we have demonstrated a connection between inner model theory and descriptive set theory. Connections between two different areas of mathematics are useful, because they allow us to use the tools from one area to tackle problems from another area. In the rest of this section we will explore in a little more detail the connection between mice and reals which are ordinal definable in $L(\mathbb{R})$.

If $\mathcal{M}$ is a mouse and $A_{(\alpha,n)} = \mathbb{R} \cap \mathcal{M}$, then certain aspects of the structure of $\mathcal{M}$ can give us information about $A_{(\alpha,n)}$. For example, we can learn something about $A_{(\alpha,n)}$ by looking at the ordinal $\omega_1^{\mathcal{M}}$. We turn to this idea now.

Suppose $\alpha \leq \omega_1^{\omega_1}$ and $\text{cof}(\alpha) > \omega$. Under the hypothesis that there are ω Woodin cardinals, we have enough determinacy to conclude that the set

$$A_{(\alpha,0)} = \bigcup_{\beta < \alpha} \text{OD}^{J_\beta(\mathbb{R})}$$

is countable. This implies in particular that $A_{(\alpha,0)} = A_{(\beta,0)}$ for some $\beta < \alpha$. Let β be the least such ordinal. Then we will say that β is the OD *cutoff ordinal* below α .

Proposition 8.6. *Assume that there exists ω Woodin cardinals. Suppose that $\alpha \leq \omega_1^{\omega_1}$ and $\text{cof}(\alpha) = \omega_1$. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is $(\alpha, 0)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\alpha, 0)$ -small. Then $\mathcal{M} \models \text{“}\omega_1 \text{ exists.”}$ Furthermore, letting $\gamma = \omega_1^{\mathcal{M}}$ and letting*

$$\beta = \sup \{ \alpha' < \alpha \mid \max(\text{code}(\alpha')) < \gamma \},$$

we have that β is the OD cutoff ordinal below α .

Proof. Let $\mathcal{M}$ be as in the statement of the proposition. It is shown in the proof of Theorem 8.1 above that $A_{(\alpha,0)} = \mathbb{R} \cap \mathcal{M}$. If $\mathcal{M}$ is *explicitly* $(\alpha, 0)$ -big then, since no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big, we have that $\mathcal{M} \models \text{“}\omega_1 \text{ exists.”}$ Note also that in this case $\mathcal{M}$ is explicitly $(\alpha, 0)$ -big above $\omega_1^{\mathcal{M}}$. Now suppose that $\mathcal{M}$ is not explicitly $(\alpha, 0)$ -big, and let $\alpha' > \alpha$ be least such that $\mathcal{M}$ is explicitly $(\alpha', 0)$ -big. Then α' cannot be a successor ordinal or a limit ordinal of cofinality ω , so α' must be a limit ordinal of cofinality ω_1 . So, as no proper initial segment of $\mathcal{M}$ is $(\alpha', 0)$ -big, $\mathcal{M} \models \text{“}\omega_1 \text{ exists.”}$ Note also that in this case $\mathcal{M}$ is explicitly $(\alpha', 0)$ -big above $\omega_1^{\mathcal{M}}$. In any case we have that $\mathcal{M} \models \text{“}\omega_1 \text{ exists.”}$

Now let $\gamma = \omega_1^{\mathcal{M}}$ and let

$$\beta = \sup \{ \alpha' < \alpha \mid \max(\text{code}(\alpha')) < \gamma \}.$$

It is easy to see that $\max(\text{code}(\beta)) = \gamma$, and that β is a limit ordinal of cofinality ω . Let δ be the OD cutoff ordinal below α . We will show that $\beta = \delta$.

First we will show that $\delta \leq \beta$. Let $\alpha' < \delta$, and we will show that $\alpha' < \beta$. Since $\alpha' < \delta$, there is an $x \in \mathbb{R}$ such that $x \in A_{(\alpha,0)}$ but $x \notin A_{(\alpha',0)}$. Since $A_{(\alpha,0)} = \mathbb{R} \cap \mathcal{M}$, there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $x \in \mathbb{R} \cap \mathcal{N}$ and $\text{OR}^{\mathcal{N}} < \gamma$. Fix such an $\mathcal{N}$. We claim that $\mathcal{N}$ is $(\beta, 0)$ -petite. For suppose $\mathcal{N}$ were not $(\beta, 0)$ -petite. Since $\mathcal{N}$ is $(\alpha, 0)$ -petite, there must be some (α_0, n_0) with $(\beta, 0) \leq_{\text{lex}} (\alpha_0, n_0) <_{\text{lex}} (\alpha, 0)$ such that $\mathcal{N}$ is explicitly (α_0, n_0) -big. Since $\beta \leq \alpha_0 < \alpha$, we have that $\max(\text{code}(\alpha_0)) \geq \gamma$. But this is impossible because according to Lemma 3.18, $\max(\text{code}(\alpha_0)) \leq \text{OR}^{\mathcal{N}}$. Thus $\mathcal{N}$ is $(\beta, 0)$ -petite. Then, as in the proof of Theorem 8.1 above, it follows that $x \in A_{(\beta,0)}$. Since $x \notin A_{(\alpha',0)}$ it follows that $\alpha' < \beta$.

Now we will show that $\beta \leq \delta$. Let $\alpha' < \beta$ and we will show that $\alpha' < \delta$. Since $\alpha' < \beta$, there is an ordinal ξ with $\alpha' \leq \xi < \alpha$ such that $\max(\text{code}(\xi)) < \gamma$. Fix such a ξ . As we pointed out in the first paragraph of this proof, $\mathcal{M}$ is explicitly $(\alpha_0, 0)$ -big above γ , for some $\alpha_0 \geq \alpha$. Since $\xi < \alpha$ and $\max(\text{code}(\xi)) < \gamma$, we have by Lemma 3.26 that for all $n \in \omega$, there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is explicitly (ξ, n) -big above γ . In particular there is a proper initial segment $\mathcal{N} \triangleleft \mathcal{M}$ such that $\mathcal{N}$ is explicitly $(\xi, 1)$ -big (above 0.) Let $\mathcal{N}$ be the $\leq$ -least such initial segment. By our second corollary to the Condensation Theorem, Corollary 1.12 on page 14, $\text{OR}^{\mathcal{N}} < \gamma$. Thus there is a real x such that $x \in \mathcal{M}$ but $x \notin \mathcal{N}$. Fix such a real x . Since $x \in \mathcal{M}$, $x \in A_{(\alpha,0)}$. Since $x \notin \mathcal{N}$, $x \notin A_{(\xi,0)}$. Thus $\xi < \delta$. Since $\alpha' \leq \xi$, we have shown that $\alpha' < \delta$. $\square$

Below we highlight two particular concrete examples of the previous proposition: $\alpha = \omega_1$, and $\alpha = \omega_1^{\omega_1}$.

Corollary 8.7. *Assume that there exists ω Woodin cardinals. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is $(\omega_1, 0)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\omega_1, 0)$ -small. Let $\gamma = \omega_1^{\mathcal{M}}$. Then γ is the OD cutoff ordinal below ω_1 .*

Proof. Let

$$\beta = \sup \{ \alpha' < \omega_1 \mid \max(\text{code}(\alpha')) < \gamma \}.$$

If $\alpha' < \omega_1$ then $\text{code}(\alpha') = \{(0, \alpha')\}$, and so $\max(\text{code}(\alpha')) = \alpha'$. Thus $\beta = \gamma$. $\square$

Corollary 8.8. *Assume that there exists ω Woodin cardinals. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is $(\omega_1^{\omega_1}, 0)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\omega_1^{\omega_1}, 0)$ -small. Let $\gamma = \omega_1^{\mathcal{M}}$. Then ω_1^γ is the OD cutoff ordinal below $\omega_1^{\omega_1}$.*

Proof. Let

$$\beta = \sup \{ \alpha' < \omega_1^{\omega_1} \mid \max(\text{code}(\alpha')) < \gamma \}.$$

$\text{code}(\omega_1^\gamma) = \{(\gamma, 1)\}$. If $\max(\text{code}(\alpha')) < \gamma$ then it follows that $\alpha' < \omega_1^\gamma$. Furthermore, there are cofinally many $\alpha' < \omega_1^\gamma$ such that $\max(\text{code}(\alpha')) < \gamma$. Thus $\beta = \omega_1^\gamma$. $\square$

We continue with our program of exploring the connections between the structure of certain mice, and certain sets of ordinal definable reals. In this paper we have defined two distinct notions: the notion of (α, n) -big, and the notion of *explicitly* (α, n) -big. One might ask, what is the significance of a mouse $\mathcal{M}$ which is (α, n) -big but not explicitly (α, n) -big? The next proposition sheds some light on this question. The proposition says that α must be in the midst of a long gap in which no new reals are ordinal defined in $L(\mathbb{R})$.

Proposition 8.9. *Assume that there exists ω Woodin cardinals. Suppose $(2, 0) \leq_{\text{lex}} (\alpha, n) <_{\text{lex}} (\omega_1^{\omega_1}, 0)$. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is (α, n) -big, but every proper initial segment of $\mathcal{M}$ is (α, n) -small. Suppose that $\mathcal{M}$ is not explicitly (α, n) -big. Then there is an ordinal $\alpha^* > \alpha$ such that $\text{cof}(\alpha^*) = \omega_1$, and such that α is greater than the OD cutoff ordinal below α^* .*

Proof. Let (α^*, n^*) be the lexicographically least pair such that $(\alpha, n) \leq_{\text{lex}} (\alpha^*, n^*)$, and $\mathcal{M}$ is explicitly (α^*, n^*) -big. We are assuming that $(\alpha, n) <_{\text{lex}} (\alpha^*, n^*)$. It follows that $n^* = 0$ and $\alpha^* > \alpha$. Then α^* must be a limit ordinal of cofinality ω_1 . By definition, $\mathcal{M}$ is active. Let κ be the critical point of the last extender on the $\mathcal{M}$ sequence. Since $\mathcal{M}$ is not explicitly (α, n) -big, we must have that $\max(\text{code}(\alpha)) \geq \kappa$. In particular $\max(\text{code}(\alpha)) > (\omega_1)^{\mathcal{M}}$. Now $\mathcal{M}$ is the $\leq$ -least initial segment of $\mathcal{M}$ which is $(\alpha^*, 0)$ -big. So by Proposition 8.6 above, α is greater than the OD cutoff ordinal below α^* . $\square$

We will mention one last result about the connection between the structure of mice and ordinal definable reals. As we have pointed out, assuming determinacy there are

only countably many reals which are ordinal definable in $L(\mathbb{R})$. So there will be many pairs (α, n) so that $A_{(\alpha, n)} = A_{(\alpha, n+1)}$. If $A_{(\alpha, n)} \neq A_{(\alpha, n+1)}$, let us say that $A_{(\alpha, n+1)}$ is *non-trivial*. Let $\mathcal{M}_1$ be the least mouse which is (α, n) -big, and let $\mathcal{M}_2$ be the least mouse which is $(\alpha, n+1)$ -big. Of course we can tell whether or not $A_{(\alpha, n+1)}$ is trivial by checking whether or not $\mathbb{R} \cap \mathcal{M}_2 \subseteq \mathcal{M}_1$. Our last proposition says that we can also check whether or not $A_{(\alpha, n+1)}$ is trivial by comparing $\text{code}(\alpha)$ with $\omega_1^{\mathcal{M}_2}$.

Proposition 8.10. *Assume that there exists ω Woodin cardinals. Suppose that $\alpha < \omega_1^{\omega_1}$, $\text{cof}(\alpha) = \omega_1$, and $n \geq 0$. Let $\mathcal{M}$ be a countable, realizable premouse such that $\mathcal{M}$ is $(\alpha, n+1)$ -big, but every proper initial segment of $\mathcal{M}$ is $(\alpha, n+1)$ -small. Then $A_{(\alpha, n+1)}$ is non-trivial, if and only if $\max(\text{code}(\alpha)) < \omega_1^{\mathcal{M}}$.*

Proof. First suppose that $\max(\text{code}(\alpha)) < \omega_1^{\mathcal{M}}$. We claim that $\mathcal{M}$ is *explicitly* $(\alpha, n+1)$ -big. For otherwise, as in the proof of the previous proposition, we must have that $\mathcal{M}$ is active and $\max(\text{code}(\alpha))$ is greater than the critical point of the last extender on the $\mathcal{M}$ sequence. But this contradicts our assumption that $\max(\text{code}(\alpha)) < \omega_1^{\mathcal{M}}$. Thus $\mathcal{M}$ is explicitly $(\alpha, n+1)$ -big. Since no proper initial segment of $\mathcal{M}$ is explicitly $(\alpha, n+1)$ -big, $\mathcal{M}$ is explicitly $(\alpha, n+1)$ -big above $\omega_1^{\mathcal{M}}$. Since $\max(\text{code}(\alpha)) < \omega_1^{\mathcal{M}}$, Lemma 3.26 tells us that there is a *proper* initial segment of $\mathcal{M}$ which is explicitly (α, n) -big above $\omega_1^{\mathcal{M}}$. Let $\mathcal{N}$ be the $\trianglelefteq$ -least initial segment of $\mathcal{M}$ which is explicitly (α, n) -big (above 0.) Our second corollary to the Condensation Theorem, Corollary 1.12, implies that $\text{OR}^{\mathcal{N}} < \omega_1^{\mathcal{M}}$. So there is a real x such that $x \in \mathcal{M}$ but $x \notin \mathcal{N}$. Since $x \in \mathcal{M}$, $x \in A_{(\alpha, n+1)}$. Since $x \notin \mathcal{N}$, $x \notin A_{(\alpha, n)}$. So $A_{(\alpha, n+1)}$ is non-trivial.

Conversely, suppose that $A_{(\alpha, n+1)}$ is non-trivial. Let $x \in A_{(\alpha, n+1)} - A_{(\alpha, n)}$. Let $\mathcal{N} \triangleleft \mathcal{M}$ be such that $x \in \mathcal{N}$ and $\text{OR}^{\mathcal{N}} < \omega_1^{\mathcal{M}}$. Then $\mathcal{N}$ is not (α, n) -petite. (If $\mathcal{N}$ were (α, n) -petite then we would have that $x \in A_{(\alpha, n)}$.) Since $\text{cof}(\alpha) = \omega_1$, (α, n) -petite is synonymous with (α, n) -small. So $\mathcal{N}$ is not (α, n) -small. But $\mathcal{N}$ is $(\alpha, n+1)$ -small. Thus $\mathcal{N}$ must be *explicitly* (α, n) -big. By Lemma 3.18, $\max(\text{code}(\alpha)) \leq \text{OR}^{\mathcal{N}}$. Thus $\max(\text{code}(\alpha)) < \omega_1^{\mathcal{M}}$. $\square$

REFERENCES

- [Ke1] A. S. Kechris, *The Theory of Countable Analytical Sets*, **Trans. Amer. Math. Soc.**, vol 202 (1975), pp. 259-297.

- [Ke2] A. S. Kechris, *Countable Ordinals and the Analytical Hierarchy I*, ***Pacific Journal of Mathematics***, vol. 60 (1975), pp. 223-227.
- [Ke3] A. S. Kechris, *Measure and Category in Effective Descriptive Set Theory*, ***Annals of Math. Logic***, vol. 5 (1973), pp. 337-384.
- [Ke4] A. S. Kechris, *Projective Ordinals and Countable Analytical Sets*, Ph. D. Thesis, UCLA, 1972.
- [KeMaSo] A. S. Kechris, D. A. Martin, R. M. Solovay, *Introduction to Q -Theory*, ***Cabal Seminar 79-81, Lecture Notes in Mathematics***, vol. 1019, Springer-Verlag, 1983, pp. 199-281.
- [Ma1] D. A. Martin, *The Largest Countable This, That, and the Other*, ***Cabal Seminar 79-81, Lecture Notes in Mathematics***, vol. 1019, Springer-Verlag, 1983, pp. 97-106.
- [Ma2] D. A. Martin, untitled book on large cardinals and determinacy, in preparation.
- [MaSt] D. A. Martin, J. R. Steel, *Iteration Trees*, ***Journal of the AMS***, vol. 7, Number 1, January, 1994.
- [MiSt] W. J. Mitchell, J. R. Steel, *Fine Structure and Iteration Trees*, ***Lecture Notes in Logic***, vol. 3, Springer-Verlag, 1994.
- [Mo] Y. N. Moschovakis, ***Descriptive Set Theory***, North Holland, 1980.
- [Ru1] M. Rudominer, *Mouse Sets Definable in $L(\mathbb{R})$* , Ph. D Thesis, UCLA, 1995.
- [Ru2] M. Rudominer, *On the Hierarchy of Ordinal Definable Reals in $L(\mathbb{R})$* , forthcoming.
- [St1] J. R. Steel, *Scales in $L(\mathbb{R})$* , ***Cabal Seminar 79-81, Lecture Notes in Mathematics***, vol. 1019, Springer-Verlag, 1983, pp. 107-156.
- [St2] J. R. Steel, *Inner Models with Many Woodin Cardinals*, ***Annals of Pure and Applied Logic***, vol. 65 (1993), pp. 185-209.
- [St3] J. R. Steel, *The Core Model Iterability Problem*, to appear.
- [St4] J. R. Steel, *Core Models with More Woodin Cardinals*, to appear.
- [St5] J. R. Steel, *Projectively Wellordered Inner Models*, ***Annals of Pure and Applied Logic***, vol. 74 (1995), issue 1, pp. 77 -104.

DEPARTMENT OF MATHEMATICS, FLORIDA INTERNATIONAL UNIVERSITY, MIAMI, FL 33199

E-mail address: rudomine@fiu.edu